

FAUNA AQUATICA AUSTRIACA

CILIOPHORA (Ciliates)

Erna Aescht, Wilhelm Foissner, Hubert Blatterer,
Fritz Kohmann & Helmut Berger

Dr. Erna Aescht
Leitung der Sammlung Wirbellose Tiere
Biologiezentrum des Oberösterreichischen Landesmuseums
J.-W.-Klein-Str. 73
A-4040 Linz
e.aescht@landesmuseum.at

Univ. Prof. (i.R.) Dr. Wilhelm Foissner
Universität Salzburg
Hellbrunnerstr. 34
A-5020 Salzburg
wilhelm.foissner@sbg.ac.at

Mag. Dr. Hubert Blatterer
Amt der Oö. Landesregierung
Direktion Umwelt und Wasserwirtschaft, Abteilung
Oberflächengewässerwirtschaft, Gewässerschutz
Kärntnerstraße 12
A-4021 Linz
hubert.blatterer@ooe.gv.at

Dr. Fritz Kohmann
Römerstraße 15
D-56321 Brey
kohmann@t-online.de

Dr. Helmut Berger
Technisches Büro für Ökologie
Radetzkystrasse 10
A-5020 Salzburg
berger.helmut@protozoology.com

Quotation note

Aescht, E., Foissner, W., Blatterer, H., Kohmann, F. & H. Berger
(2017): Ciliophora. In Moog, O. & A. Hartmann (Eds.): Fauna
AquatICA Austriaca, 3. Edition 2017. BMLFUW, Wien.



Austria takes a special status in the international research on ciliates. Wilhelm Foissner, one of the most experienced and productive taxonomists, and his working group made Austria to a hotspot of ciliate-diversity research (Aescht & Berger 2008a, b, Aescht 2008, Berger & Al-Rasheid 2008).

In addition, a natural monument (“Krauthügel”) expressly for ciliated protozoa in the city of Salzburg (Austria) is unique in the world (Foissner et al. 2012, Cotterill et al. 2013). This protected ephemeral pond harbours about 150 ciliate species. Such a conservation status cannot be taken as granted for a group of microorganisms which can be identified only by specialists (Foissner 1993, Foissner & Foissner 1988, Foissner et al. 1991, 1992, 1994, 1995, 1999, Foissner & Berger 1996, Berger et al. 1997, Berger 1999, 2006, 2008, 2011, Aescht 2001, 2008, Blatterer 2002, 2008, Aescht & Berger 2008a, b, Berger & Foissner 2003, Aescht 2012, 2013).

Foissner et al. (1991, 1992, 1994, 1995) are the standard references for the identification and ecological and saprobiological classification of ciliate indicator species (“ciliate-atlas”), while Blatterer (1995, see also ÖNORM M 6118) published a specific saprobic index for ciliates. Unfortunately, the Austrian and European administrations do not support the use of ciliates for water quality assessment since several years and thus no further species have been classified ecologically. In the Austrian guidelines for the saprobiological classification of running waters, the ciliates are only optionally included as additional group for specific issues (Moog et al. 1999). Consequently, only few monitoring programs dealing with ciliates have been realized in recent years (for reviews, see Aescht 2012, 2013) and thus the new edition of the Fauna Aquatica Austriaca contains an updated list of Austrian records, but the chapters dealing with the ecology and saprobiology are identical with previous editions.

The names of the species have been checked carefully. In spite of that, we cannot guarantee that all details are correct. A question mark ahead of a combining author means that this author is uncertain. In *Alinostoma* Jankowski, 1980 (Cyrtophoria) and in two genera of peritrichs (*Pseudocarchesium* Sommer, 1951; *Pallitrichodina* Van As & Basson in Aescht, 2001) important notes on their taxonomy and nomenclature have been made. The classification follows mainly Lynn (2008).

Acknowledgements

Supported by the Austrian Science Fund (FWF): Project P26974-B25 (H. Berger).

References

- Aescht, E. (2001): Catalogue of the generic names of ciliates (Protozoa, Ciliophora). *Denisia* 1: 1–350.
- Aescht, E. (2008): Annotated catalogue of “type material” of ciliates (Ciliophora) and some further protists at the Upper Austrian Museum in Linz (Austria) including a guideline for “typification” of species. *Denisia* 23: 125–234.
- Aescht, E. (2012): Wimperlänge (Protista: Ciliophora) aus Oberösterreichs Gemeinden und 7 weiteren Bundesländern – Daten zur Checkliste der Fauna Österreichs. *Beitr. Naturk. Oberösterreichs* 22: 83–832.
- Aescht, E. (2013): Ciliophora. *Biosystematics and Ecology Series No. 30 / Checklisten der Fauna Österreichs No. 7*: 111 pp.
- Aescht, E. & H. Berger (Sci. Ed.) (2008a): The Wilhelm Foissner Festschrift. A tribute to an outstanding protistologist on the occasion of his 60th birthday. *Denisia* 23: 1–462.
- Aescht, E. & H. Berger (2008b): Univ.-Prof. Dr. Wilhelm Foissner – 60 years: a biographical sketch and bibliography. *Denisia* 23: 15–46.
- Aescht, E., Foissner W., Blatterer H., Kohmann F. & H. Berger (2017): Ciliophora. In Moog, O. & A. Hartmann (Eds.): *Fauna Aquatica Austriaca*, 3. Lieferung 2017. BMLFUW, Wien.
- Albrecht, J. (1984): Zur Autökologie ausgewählter Aufwuchsciliaten des Weser-Flußsystems (Protozoa: Ciliophora). *Decheniana* 137: 132–167.
- Berger, H. (1999): Monograph of the Oxytrichidae (Ciliophora, Hypotrichia). *Monographiae biol.* 78: i–xii, 1–1080.
- Berger, H. (2006): Monograph of the Urostyloidea (Ciliophora, Hypotricha). *Monographiae biol.* 85: i–xvi, 1–1303.
- Berger, H. (2008): Monograph of the Amphisiellidae and Trachelostylidae (Ciliophora, Hypotricha). *Monographiae biol.* 88: i–xvi, 1–737.
- Berger, H. (2011): Monograph of the Gonostomatidae and Kahliellidae (Ciliophora, Hypotricha). *Monographiae biol.* 90: i–xiv, 1–741.
- Berger, H. & K. A. S. Al-Rasheid (2008): Wilhelm Foissner: nomenclatural and taxonomic summary 1967–2007. *Denisia* 23: 65–124.
- Berger, H. & W. Foissner (2003): Illustrated guide and ecological notes to ciliate indicator species (Protozoa, Ciliophora) in running waters, lakes, and sewage plants. *Handbuch Angew. Limnol.* 17. Erg.Lfg. III-2.1: 1–160.
- Berger, H., Foissner, W. & F. Kohmann (1997): Bestimmung und Ökologie der Mikrosaprobien nach DIN 38410. Gustav Fischer, Stuttgart, Jena, Lübeck, Ulm: viii + 291 pp.
- Blatterer, H. (1995): Verbessertes Verfahren zur Berechnung des Saprobienindex mittels Ciliaten (Ciliophora, Protozoa). *Lauterbornia* 20: 23–36.
- Blatterer, H. (2002): Some conditions for the distribution and abundance of ciliates (Protozoa) in running waters – Do we really find every species everywhere? *Verh. Internat. Verein Limnol.* 28: 1046–1049.
- Blatterer, H. (2008): Umfassende Zusammenschau von Freiland-Erkenntnissen über Fließgewässer-Ciliaten (Protozoa, Ciliophora). *Denisia* 23: 337–359.
- Blatterer, H. & W. Foissner (1990): Beiträge zur Ciliatenfauna (Protozoa: Ciliophora) der Amper (Bayern, Bundesrepublik Deutschland). *Arch. Protistenk.* 138: 93–115.
- Cotterill, F. P. D., Augustin, H., Medicus, R. & W. Foissner (2013): Conservation of protists: The Krauthügel pond in Austria. *Diversity* 2013, 5: 374–392.
- Finlay, B. J. (1982): Procedures for the isolation, cultivation and identification of protozoa. In: Burns R. G. & J. H. Slater (Eds.): *Experimental Microbial Ecology*, Blackwell Scientific Publications, Oxford; pp. 44–65.
- Foissner, W. (1993): Colpodea. *Protozoenfauna* 4/1: i–x, 1–798.

- Foissner, W. & H. Berger (1996): A user-friendly guide to the ciliates (Protozoa, Ciliophora) commonly used by hydrobiologists as bioindicators in rivers, lakes, and waste waters, with notes on their ecology. *Freshw. Biol.* 35: 375–482.
- Foissner, W. & I. Foissner (1988): Stamm Ciliophora. *Catalogus Faunae Austriae Ic*: 1–147.
- Foissner, W., Blatterer, H., Berger, H., & F. Kohmann (1991): Taxonomische und ökologische Revision der Ciliaten des Saprobiensystems - Band I: Cytrophorida, Oligotrichida, Hypotrichida, Colpodea. *Informationsberichte des Bayer. Landesamtes für Wasserwirtschaft* 1/91: 478 pp.
- Foissner, W., Berger, H. & F. Kohmann (1992): Taxonomische und ökologische Revision der Ciliaten des Saprobiensystems - Band II: Peritricha, Heterotrichida, Odontostomatida. *Informationsberichte des Bayer. Landesamtes für Wasserwirtschaft* 5/92: 502 pp.
- Foissner, W., Berger, H. & F. Kohmann (1994): Taxonomische und ökologische Revision der Ciliaten des Saprobiensystems - Band III: Hymenostomata, Prostomatida, Nassulida. *Informationsberichte des Bayer. Landesamtes für Wasserwirtschaft* 1/94: 548 pp.
- Foissner, W., Berger, H., Blatterer H. & F. Kohmann (1995): Taxonomische und ökologische Revision der Ciliaten des Saprobiensystems - Band IV: Gymnostomatea, *Loxodes*, Suctorina. *Informationsberichte des Bayer. Landesamtes für Wasserwirtschaft* 1/95: 540 pp.
- Foissner, W., Berger, H. & J. Schaumburg (1999): Identification and ecology of limnetic plankton ciliates. *Informationsberichte des Bayer. Landesamtes für Wasserwirtschaft* 3/99: 793 pp.
- Foissner, W., Chao, A. & L. A. Katz (2008): Diversity and geographic distribution of ciliates (Protista: Ciliophora). *Biodiversity & Conservation* 17: 345–363.
- Foissner, W., Medicus, R. & H. Augustin (2012): Ein Naturdenkmal für Wimpertierchen! *Natur und Land* 98/4: 6–7.
- ICZN (International Commission on Zoological Nomenclature) (1999): International code of zoological nomenclature. International Trust for Zoological Nomenclature, London: 306 pp.
- Jankowski, A. W. (1994): Taxonomy of Ciliophora. 2. New species of *Opercularia*, *Entziella* and *Circolagenophrys* from the Black Sea and Pacific, and taxonomic notes on other peritrichs (Peritricha). *Zoosyst. Rossica* 2 (Jahr 1993): 217–222.
- Lynn, D. H. (2008): The Ciliated Protozoa. Characterization, Classification and Guide to the Literature. Springer; xxxiv + 605 pp.
- Moog, O., Chovanec, A., Hinteregger, J. & A. Römer (1999): Richtlinie zur Bestimmung der saprobiologischen Gewässergüte von Fließgewässern. Bundesministerium für Land- und Forstwirtschaft, Wasserwirtschaftskataster, Wien, ISBN 3-85 174-033-5: 144 pp.
- ÖNORM M 6118 (Ausgabe 2005-12-01): Richtlinien für die ökologische Untersuchung von Fließgewässern. Ciliaten (Protozoa). Guidelines for the ecological study and assessment of rivers - Ciliates (Protozoa). Österreichisches Normungsinstitut, 1020 Wien. (Autoren: Berger, H., Blatterer, H., Petz, W., Schiftner, U., Foissner, W.).
- Precht, H. (1935): Epizoen der Kieler Bucht. *Nova Acta Leopoldina* 3: 405–474.
- Sládeček, V., Zelinka, M., Rothschein, J. & V. Moravcova (1981): Biologicky rozbor povrchové vody. Komentár k CSN 83 0532 - části 6: Stanovení saprobiálního indexu. Vydalo Vydavatelství Úradu pro normalizaci a měření, Praha (in Czech). 186 pp.
- Sommer, G. (1951): Die peritrichen Ciliaten des Großen Plöner Sees. *Arch. Hydrobiol.* 44: 349–440.
- Van As, J. G. & L. Basson (1993): On the biology of *Pallitrichodina rogenae* gen. n., sp. n. and *P. stephani* sp. n. (Ciliophora: Peritrichida), mantle cavity symbionts of the giant African snail *Achatina* in Mauritius and Taiwan. *Acta Protozool.* 32: 47–62.

Species inventory

Subphylum Postciliodesmatophora

Class Karyorelictida

Order Loxodida

Genus *Loxodes* EHRENBERG, 1830

Loxodes magnus STOKES, 1887

Loxodes rostrum (MÜLLER, 1773) EHRENBERG, 1830

Loxodes striatus (ENGELMANN, 1862) PENARD, 1917

Loxodes vorax STOKES, 1884

Class Heterotrichea

Order Heterotrichida

Genus *Blepharisma* PERTY, 1849

Blepharisma bimicronucleatum VILLENEUVE-BRACHON, 1940

Blepharisma elongatum (STOKES, 1884) KAHL, 1926

Blepharisma hyalinum PERTY, 1852

Blepharisma lateritium (EHRENBERG, 1831) STEIN, 1859

Blepharisma ovatum (STOKES, 1884) PENARD, 1922

Blepharisma persicinum PERTY, 1849

Blepharisma steini KAHL, 1932

Blepharisma undulans STEIN, 1867

Genus *Climacostomum* STEIN, 1859

Climacostomum minimum FOISSNER, 1980

Climacostomum patulum (MÜLLER, 1786) KAHL, 1932

Climacostomum virens (EHRENBERG, 1838) STEIN, 1859

Genus *Condylostoma* BORY, 1824

Condylostoma caudatum LAUTERBORN, 1908

Genus *Folliculina* LAMARCK, 1816

Folliculina boltoni KENT, 1881

Genus *Linostomella* AESCHT in FOISSNER, BERGER & SCHAUMBURG, 1999

Linostomella vorticella (EHRENBERG, 1834) AESCHT in FOISSNER, BERGER & SCHAUMBURG, 1999

Genus *Spirostomum* EHRENBERG, 1834

Spirostomum ambiguum (MÜLLER, 1786) EHRENBERG, 1834

Spirostomum caudatum (MÜLLER, 1786) DELPHY, 1939

Spirostomum minus ROUX, 1901

Spirostomum teres CLAPARÈDE & LACHMANN, 1858

Genus *Stentor* OKEN, 1815

Stentor amethystinus LEIDY, 1880

Stentor coeruleus (PALLAS, 1766) EHRENBERG, 1831

Stentor igneus EHRENBERG, 1838

Stentor muelleri EHRENBERG, 1831

Stentor multififormis (MÜLLER, 1786) EHRENBERG, 1838

Stentor niger (MÜLLER, 1773) EHRENBERG, 1831
Stentor pallidus FOISSNER, 1980
Stentor polymorphus (MÜLLER, 1773) EHRENBERG, 1830
Stentor roeselii EHRENBERG, 1835

Subphylum Intramacronucleata

Class Spirotrichea

Subclass Hypotrichia

Genus Chaetospira LACHMANN, 1856

Chaetospira muelleri LACHMANN, 1856

Genus Cladotricha GAJEWSKAJA, 1926

Cladotricha sigmoidea RUINEN, 1938

Genus Deviata EIGNER, 1995

Deviata abbrevescens EIGNER, 1995

Genus Hypotrichidium ILOWAISKY, 1921

Hypotrichidium conicum ILOWAISKY, 1921

Genus Paraholosticha WENZEL, 1953

Paraholosticha herbicola (KAHL, 1932) WENZEL, 1953

Genus Lamtostyla BUITKAMP, 1977

Lamtostyla decorata FOISSNER, AGATHA & BERGER, 2002

Genus Mucotrichidium FOISSNER, OLEKSIV & MÜLLER, 1990

Mucotrichidium hospes (EHRENBERG, 1831) FOISSNER, OLEKSIV & MÜLLER, 1990

Genus Parastrongylidium FLEURY & FRYD-VERSAVEL, 1985

Parastrongylidium oswaldi AESCHT & FOISSNER, 1992

Genus Psilotricha STEIN, 1859

Psilotricha acuminata STEIN, 1859

Psilotricha succisa (MÜLLER, 1786) FOISSNER, 1983

Genus Stichotricha PERTY, 1849

Stichotricha aculeata WRZESNIEWSKI, 1866

Stichotricha secunda PERTY, 1849

Stichotricha socialis GRUBER, 1880

Stichotricha tubicola (GRUBER, 1880) BORROR, 1972

Genus Strongylidium STERKI, 1878

Strongylidium lanceolatum KOWALEWSKI, 1882

Genus Wallackia FOISSNER, 1976

Wallackia schiffmanni FOISSNER, 1976

Family Oxytrichidae

Genus Allotricha STERKI, 1878

Allotricha mollis STERKI, 1878

Genus Australocirrus BLATTERER & FOISSNER, 1988*Australocirrus aspoecki* (FOISSNER, 2004) KUMAR & FOISSNER, 2015**Genus Cyrtohymena** FOISSNER, 1989*Cyrtohymena citrina* (BERGER & FOISSNER, 1987) FOISSNER, 1989**Genus Gastrostyla** ENGELMANN, 1862*Gastrostyla mystacea* (STEIN, 1859) STERKI, 1878*Gastrostyla steinii* ENGELMANN, 1862**Genus Histriculus** CORLISS, 1960*Histriculus complanatus* (STOKES, 1887) CORLISS, 1960*Histriculus histrio* (MÜLLER, 1773) CORLISS, 1960**Genus Kerona** MÜLLER, 1786*Kerona pediculus* (MÜLLER, 1773) BLOCHMANN, 1886**Genus Neokeronopsis** WARREN, FYDA & SONG, 2002*Neokeronopsis spectabilis* (KAHL, 1932) WARREN, FYDA & SONG, 2002**Genus Onychodromus** STEIN, 1859*Onychodromus grandis* STEIN, 1859**Genus Oxytricha** BORY, 1824*Oxytricha acidotolerans* WEISSE, MOSER, SCHEFFEL, STADLER, BERENDONK, WEITHOFF & BERGER, 2013*Oxytricha chlorelligera* KAHL, 1932*Oxytricha fallax* STEIN, 1859*Oxytricha hymenostoma* STOKES, 1887*Oxytricha opisthomuscorum* FOISSNER, BLATTERER, BERGER & KOHMANN, 1991*Oxytricha parallela* ENGELMANN, 1862*Oxytricha saprobia* KAHL, 1932*Oxytricha setigera* STOKES, 1891*Oxytricha similis* ENGELMANN, 1862*Oxytricha siseris* VUXANOVICI, 1963**Genus Paraurostyla** BORROR, 1972*Paraurostyla weissei* (STEIN, 1859) BORROR, 1972**Genus Pleurotricha** STEIN, 1859*Pleurotricha grandis* STEIN, 1859*Pleurotricha lanceolata* (EHRENBERG, 1835) STEIN, 1859**Genus Rubrioxxytricha** BERGER, 1999*Rubrioxxytricha ferruginea* (STEIN, 1859) BERGER, 1999*Rubrioxxytricha haematoplasma* (BLATTERER & FOISSNER, 1990) BERGER, 1999**Genus Steinia** DIESING, 1866*Steinia platystoma* (EHRENBERG, 1831) DIESING, 1866*Steinia sphagnicola* FOISSNER, 1989**Genus Sterkiella** FOISSNER, BLATTERER, BERGER & KOHMANN, 1991*Sterkiella admirabilis* (FOISSNER, 1980) BERGER, 1999*Sterkiella histriomuscorum* (FOISSNER, BLATTERER, BERGER & KOHMANN, 1991) FOISSNER, BLATTERER, BERGER & KOHMANN, 1991

Genus Stylonychia EHRENBERG, 1830*Stylonychia mytilus*-Komplex*Stylonychia putrina* STOKES, 1885**Genus Tachysoma** STOKES, 1887*Tachysoma pellionellum* (MÜLLER, 1773) BORROR, 1972**Genus Tetmemena** EIGNER, 1999*Tetmemena pustulata* (MÜLLER, 1786) EIGNER, 1999**Genus Urosoma** KOWALEWSKI, 1882*Urosoma acuminata* (STOKES, 1887) BÜTSCHLI, 1889*Urosoma caudatum* (EHRENBERG, 1833) BERGER, 1999**Genus Urosomoida** HEMBERGER IN FOISSNER, 1982*Urosomoida agilis* (ENGELMANN, 1862) HEMBERGER IN FOISSNER, 1982**Family Uroleptidae****Genus Uroleptus** EHRENBERG, 1831*Uroleptus caudatus* (STOKES, 1886) BARDELE, 1981*Uroleptus gallina* (MÜLLER, 1786) FOISSNER, BLATTERER, BERGER & KOHMANN, 1991*Uroleptus lamella* EHRENBERG, 1831*Uroleptus musculus* (KAHL, 1932) FOISSNER, BLATTERER, BERGER & KOHMANN, 1991*Uroleptus piscis* (MÜLLER, 1773) EHRENBERG, 1831*Uroleptus violaceus* STEIN, 1859*Uroleptus willii* SONNTAG, STRÜDER-KYPKE & SUMMERER, 2008**Order Urostylida****Genus Anteholosticha** BERGER, 2003*Anteholosticha antecirrata* BERGER, 2006*Anteholosticha intermedia* (BERGH, 1889) BERGER, 2006*Anteholosticha monilata* (KAHL, 1928) BERGER, 2003*Anteholosticha xanthichroma* (WIRNSBERGER & FOISSNER, 1987) BERGER, 2003**Genus Australothrix** BLATTERER & FOISSNER, 1988*Australothrix gibba* (CLAPARÈDE & LACHMANN, 1859) BLATTERER & FOISSNER, 1988**Genus Diaxonella** JANKOWSKI, 1979*Diaxonella pseudorubra pseudorubra* (KALTENBACH, 1960) BERGER, 2006**Genus Holosticha** WRZESNIEWSKI, 1877*Holosticha pullaster* (MÜLLER, 1773) FOISSNER, BLATTERER, BERGER & KOHMANN, 1991**Genus Holostichides** FOISSNER, 1987*Holostichides dumonti* FOISSNER, 2000**Genus Pseudokeronopsis** BORROR & WICKLOW, 1983*Pseudokeronopsis similis* (STOKES, 1886) BORROR & WICKLOW, 1983**Genus Pseudourostyla** BORROR, 1972*Pseudourostyla cristata* (JERKA-DZIADOSZ, 1964) BORROR, 1972**Genus Trichototaxis** STOKES, 1891*Trichototaxis aeruginosa* FOISSNER, 1980

Genus Urostyla EHRENBERG, 1830

- Urostyla grandis* EHRENBERG, 1830
Urostyla chlorelligera FOISSNER, 1980
Urostyla viridis STEIN, 1859

Order Euplotida**Genus Aspidisca** EHRENBERG, 1830

- Aspidisca cicada* (MÜLLER, 1786) CLAPARÈDE & LACHMANN, 1858
Aspidisca lynceus (MÜLLER, 1773) EHRENBERG, 1830
Aspidisca turrita (EHRENBERG, 1831) CLAPARÈDE & LACHMANN, 1858

Genus Euplotes EHRENBERG, 1831

- Euplotes charon* (MÜLLER, 1773) EHRENBERG, 1830
Euplotes harpa STEIN, 1859
Euplotes moebiusi KAHL, 1932
Euplotes parki CURDS, 1974

Genus Euplotoides BORROR & HILL, 1995

- Euplotoides aediculatus* (PIERSON, 1943) BORROR & HILL, 1995
Euplotoides eurystomus (WRZESNIEWSKI, 1870) BORROR & HILL, 1995
Euplotoides patella (MÜLLER, 1773) BORROR & HILL, 1995

Genus Euplotopsis BORROR & HILL, 1995

- Euplotopsis affinis* (DUJARDIN, 1841) BORROR & HILL, 1995
Euplotopsis novemcarinatus (WANG, 1930) BORROR & HILL, 1995

Subclass Halteria**Order Halteriida****Genus Halteria** DUJARDIN, 1841

- Halteria bifurcata* TAMAR, 1968
Halteria chlorelligera KAHL, 1932
Halteria grandinella (MÜLLER, 1773) DUJARDIN, 1841
Halteria minuta MAEDA, 1986

Genus Meseres SCHEWIAKOFF, 1892

- Meseres corlissi* PETZ & FOISSNER, 1992

Genus Pelagohalteria FOISSNER, SKOGSTAD & PRATT, 1988

- Pelagohalteria cirrifera* (KAHL, 1932) FOISSNER, SKOGSTAD & PRATT, 1988
Pelagohalteria viridis (FROMENTEL, 1876) FOISSNER, SKOGSTAD & PRATT, 1988

Subclass Choreotrichia**Order Choreotrichida****Genus Strobilidium** SCHEWIAKOFF, 1892

- Strobilidium caudatum* (FROMENTEL, 1876) FOISSNER, 1987
Strobilidium lacustris FOISSNER, SKOGSTAD & PRATT, 1988

Subclass Oligotrichia**Order Strobiliida****Genus Rimostrombidium** JANKOWSKI, 1978*Rimostrombidium brachykinetum* KRAINER, 1995*Rimostrombidium conicum* (KAHL, 1932) PETZ & FOISSNER, 1992*Rimostrombidium humile* (PENARD, 1922) PETZ & FOISSNER, 1992*Rimostrombidium hyalinum* (MIRABDULLAEV, 1985) PETZ & FOISSNER, 1992*Rimostrombidium lacustris* (FOISSNER, SKOGSTAD & PRATT, 1988) PETZ & FOISSNER, 1992*Rimostrombidium velox* (FAURE-FREMIET, 1924) JANKOWSKI, 1978**Order Strombidiida****Genus Limnostrombidium** KRAINER, 1995*Limnostrombidium pelagicum* (KAHL, 1932) KRAINER, 1995*Limnostrombidium viride* (STEIN, 1867) KRAINER, 1995**Genus Opisthostrombidium** AGATHA, 2011*Opisthostrombidium montagnesi* (XU, SONG & WARREN, 2006) AGATHA, 2011**Genus Pelagostrombidium** KRAINER, 1991*Pelagostrombidium fallax* (ZACHARIAS, 1895) KRAINER, 1991*Pelagostrombidium mirabile* (PENARD, 1916) KRAINER, 1991**Genus Strombidium** CLAPARÈDE & LACHMANN, 1859*Strombidium rehwaldi* PETZ & FOISSNER, 1992*Strombidium turbo* CLAPARÈDE & LACHMANN, 1859**Order Tintinnida****Genus Codonella** HAECKEL, 1873*Codonella cratera* (LEIDY, 1877) IMHOF, 1885**Genus Membranicola** FOISSNER, BERGER & SCHAUMBURG, 1999*Membranicola tamari* FOISSNER, BERGER & SCHAUMBURG, 1999**Genus Tintinnidium** KENT, 1881*Tintinnidium (Semitintinnidium) semiciliatum* (STERKI, 1879) KENT, 1881 in AGATHA & STRÜDER-KYPKE (2007)*Tintinnidium (Tintinnidium) fluviatile* (STEIN, 1863) KENT, 1881*Tintinnidium (Tintinnidium) pusillum* ENTZ, 1909**Genus Tintinnopsis** STEIN, 1867*Tintinnopsis cylindrata* KOFOID & CAMPBELL, 1929**Class Armophorea****Order Armophorida****Genus Caenomorpha** PERTY, 1852*Caenomorpha lauterborni* KAHL, 1927*Caenomorpha medusula* PERTY, 1852*Caenomorpha uniserialis* LEVANDER, 1894**Genus Ludio** PENARD, 1922*Ludio parvulus* PENARD, 1922

Order Metopida**Genus Bothrostoma** STOKES, 1887*Bothrostoma undulans* STOKES, 1887**Genus Brachonella** JANKOWSKI, 1964*Brachonella caduca* (KAHL, 1927) JANKOWSKI, 1964*Brachonella caenomorphoides* FOISSNER, 1980*Brachonella galeata* (KAHL, 1927) JANKOWSKI, 1964*Brachonella spiralis* (SMITH, 1897) JANKOWSKI, 1964**Genus Metopus** CLAPARÈDE & LACHMANN, 1858*Metopus alpestris* FOISSNER, 1980*Metopus bothrostomiformis* FOISSNER, 1980*Metopus contortus* (QUENNERSTEDT, 1867) LAUTERBORN, 1916*Metopus es* (MÜLLER, 1776) LAUTERBORN, 1916*Metopus laminarius* KAHL, 1927*Metopus minor* KAHL, 1927*Metopus rectus* KAHL, 1932*Metopus spinosus* KAHL, 1927*Metopus striatus* MCMURRICH, 1884*Metopus tortus* KAHL, 1927**Genus Tropidoatractus** LEVANDER, 1894*Tropidoatractus acuminatus* LEVANDER, 1894**Class Litostomatea****Order Tracheliida****Genus Trachelius** SCHRANK, 1803*Trachelius anas* (MÜLLER, 1773) EHRENBERG, 1831*Trachelius ovum* (EHRENBERG, 1831) EHRENBERG, 1833**Order Dileptida****Genus Apodileptus** VĎAČNÝ, ORSI, BOURLAND, SHIMANO, EPSTEIN & FOISSNER, 2011*Apodileptus visscheri rhabdoplites* VĎAČNÝ & FOISSNER, 2012*Apodileptus visscheri visscheri* (DRAGESCO, 1963) VĎAČNÝ, ORSI, BOURLAND, SHIMANO, EPSTEIN & FOISSNER, 2011**Genus Dileptus** DUJARDIN, 1841*Dileptus anatinus* GOLINSKA, 1971*Dileptus jonesi* DRAGESCO, 1963*Dileptus margaritifera* (EHRENBERG, 1833) DUJARDIN, 1841*Dileptus viridis* (EHRENBERG, 1834) BUITKAMP, 1977**Genus Monilicaryon** JANKOWSKI, 1967*Monilicaryon monilatum* (STOKES, 1886) JANKOWSKI, 1967**Genus Monomacrocaryon** VĎAČNÝ, ORSI, BOURLAND, SHIMANO, EPSTEIN & FOISSNER, 2011*Monomacrocaryon gigas* (CLAPARÈDE & LACHMANN, 1859) VĎAČNÝ, ORSI, BOURLAND, SHIMANO, EPSTEIN & FOISSNER, 2011

Genus Paradileptus WENRICH, 1929*Paradileptus elephantinus* (SVEC, 1897) KAHL, 1931**Genus Pelagodileptus** FOISSNER, BERGER & SCHAUMBURG, 1999*Pelagodileptus trachelioides* (ZACHARIAS, 1894) FOISSNER, BERGER & SCHAUMBURG, 1999**Genus Pseudomonilicaryon** FOISSNER, 1997*Pseudomonilicaryon anser* (MÜLLER, 1773) VĎAČNÝ & FOISSNER, 2012**Subclass Haptoria****Order Didiniida****Genus Didinium** STEIN, 1859*Didinium nasutum* (MÜLLER, 1773) STEIN, 1859**Genus Monodinium** FABRE-DOMERGUE, 1888*Monodinium alveolatum* (KAHL, 1930) FOISSNER, BERGER & SCHAUMBURG, 1999*Monodinium balbianii balbianii* FABRE-DOMERGUE, 1888*Monodinium balbianii breviprobois* FOISSNER, BERGER & SCHAUMBURG, 1999*Monodinium balbianii rostratum* (KAHL, 1926) FOISSNER, BERGER & SCHAUMBURG, 1999*Monodinium chlorelligerum* KRÄINER, 1995*Monodinium perrieri* DELPHY, 1925**Order Haptorida****Genus Enchelydium** KAHL, 1930*Enchelydium alpinum* FOISSNER, 1980*Enchelydium piliforme* (KAHL, 1930) FOISSNER, 1984*Enchelydium simile* FOISSNER, 1980*Enchelydium trichocystis* FOISSNER, 1980**Genus Fuscheria** FOISSNER, 1983*Fuscheria nodosa nodosa* FOISSNER, 1983*Fuscheria nodosa salisburgensis* FOISSNER & GABILONDO in GABILONDO & FOISSNER, 2009**Genus Lagynophrya** KAHL, 1927*Lagynophrya acuminata* KAHL, 1935**Order Lacrymariida****Genus Lacrymaria** BORY, 1824*Lacrymaria filiformis* (BARAUD in MASKELL, 1886) FOISSNER, 1983*Lacrymaria olor* (MÜLLER, 1786) BORY, 1824*Lacrymaria pumilio* VUXANOVICI, 1962*Lacrymaria pupula* (MÜLLER, 1773) KAHL, 1930*Lacrymaria robusta* VUXANOVICI, 1959*Lacrymaria vaginifera* SONG & WILBERT, 1989*Lacrymaria viridis* (EHRENBERG, 1834) DUJARDIN, 1841**Genus Lagynus** QUENNERSTEDT, 1867*Lagynus cucumis* (PENARD, 1922) FOISSNER, 1987*Lagynus elegans* (ENGELMANN, 1862) QUENNERSTEDT, 1867*Lagynus verrucosus* FOISSNER, 1983

Genus Phialina BORY, 1824

- Phialina jankowskii* FOISSNER, 1984
Phialina macrostoma FOISSNER, 1983
Phialina vermicularis (MÜLLER, 1786) BORY, 1824
Phialina vertens (STOKES, 1885) FOISSNER & ADAM, 1979
Phialina viridis EHRENBERG, 1831

Order Pleurostomatida**Genus Acineria** DUJARDIN, 1841

- Acineria incurvata* DUJARDIN, 1841
Acineria punctata SONG & WILBERT, 1989
Acineria uncinata TUCOLESCO, 1962

Genus Amphileptus EHRENBERG, 1830

- Amphileptus carchesii* STEIN, 1867
Amphileptus falcatus SONG & WILBERT, 1989
Amphileptus meleagris (EHRENBERG, 1835) CLAPARÈDE & LACHMANN, 1859
Amphileptus piger (VUXANOVICI, 1962) SONNTAG & FOISSNER, 2004
Amphileptus pleurosigma (STOKES, 1884) FOISSNER, 1984
Amphileptus plurivacuolatus (FOISSNER, 1978) FOISSNER, 1987
Amphileptus procerus (PENARD, 1922) KAHL, 1943
Amphileptus punctatus (KAHL, 1926) KAHL, 1943

Genus Apoamphileptus LIN & SONG, 2004

- Apoamphileptus claparedii* (STEIN, 1867) LIN & SONG, 2004

Genus Litonotus WRZESNIEWSKI, 1870

- Litonotus alpestris* FOISSNER, 1978
Litonotus anguilloides SRÁMEK-HUSEK, 1957
Litonotus carinatus STOKES, 1885
Litonotus crystallinus VUXANOVICI, 1960
Litonotus cygnus (MÜLLER, 1773) FOISSNER, BERGER, BLATTERER & KOHMANN, 1995
Litonotus fusidens KAHL, 1926
Litonotus lamella (MÜLLER, 1773) SCHEWIAKOFF, 1896
Litonotus obtusus (MAUPAS, 1888) KAHL, 1932
Litonotus trichocystiferus FOISSNER, 1984
Litonotus uninucleatus FOISSNER, 1978
Litonotus varsaviensis (WRZESNIEWSKI, 1866) WRZESNIEWSKI, 1870

Genus Loxophyllum DUJARDIN, 1841

- Loxophyllum helus* (STOKES, 1884) PENARD, 1922
Loxophyllum meleagris (MÜLLER, 1773) DUJARDIN, 1841
Loxophyllum semilunare VUXANOVICI, 1959

Genus Opisthodon STEIN, 1859

- Opisthodon niemeccensis* STEIN, 1859

Genus Pseudoamphileptus FOISSNER, 1983

- Pseudoamphileptus macrostoma* (CHEN, 1955) FOISSNER, 1983

Genus Siroloxophyllum FOISSNER & LEIPE, 1995*Siroloxophyllum utricularium* (PENARD, 1922) FOISSNER & LEIPE, 1995**Order Spathidiida****Genus Acaryophrya** ANDRÉ, 1915*Acaryophrya sphaerica* (GELEI, 1934) DINGFELDER, 1962**Genus Acropisthium** PERTY, 1852*Acropisthium mutabile* PERTY, 1852**Genus Actinobolina** STRAND, 1928*Actinobolina radians* (STEIN, 1867) STRAND, 1928*Actinobolina smalli* HOLT, LYNN & CORLISS, 1973*Actinobolina vorax* (WENRICH, 1929) KAHL, 1930**Genus Actinorhabdos** FOISSNER, 1984*Actinorhabdos trichocystifera* FOISSNER, 1984**Genus Apertospathula** FOISSNER, AGATHA & BERGER, 2002*Apertospathula implicata* (KAHL, 1930) FOISSNER & OERTEL, 2009**Genus Balantidion** EBERHARD, 1862*Balantidion pellucidum* EBERHARD, 1862**Genus Belonophrya** ANDRE, 1914*Belonophrya pelagica* ANDRE, 1914**Genus Berghophrya** FOISSNER, 2003*Berghophrya emmae* (BERGH, 1896) FOISSNER, 2003**Genus Cataphractes** FOISSNER, 2016*Cataphractes austriacus* FOISSNER, 2016**Genus Chaenea** QUENNERSTEDT, 1867*Chaenea limicola* LAUTERBORN, 1901*Chaenea stricta* (DUJARDIN, 1841) FOISSNER, BERGER, BLATTERER & KOHMANN, 1995**Genus Cranotheridium** SCHEWIAKOFF, 1893*Cranotheridium foliosum* (FOISSNER, 1983) WIRNSBERGER, FOISSNER & ADAM, 1984**Genus Enchelyodon** CLAPARÈDE & LACHMANN, 1859*Enchelyodon anulatus* FOISSNER, 1984*Enchelyodon farctus* CLAPARÈDE & LACHMANN, 1859**Genus Enchelys** MÜLLER, 1773*Enchelys arcuata* CLAPARÈDE & LACHMANN, 1859*Enchelys binucleata* FOISSNER, 1983*Enchelys farcimen* MÜLLER, 1773*Enchelys gasterosteus* KAHL, 1926*Enchelys mutans* (MERMUD, 1914) KAHL, 1930*Enchelys nebulosa* MÜLLER, 1773*Enchelys pupa* MÜLLER, 1786**Genus Epitholiolus** FOISSNER, AGATHA & BERGER, 2002*Epitholiolus attenuatus* (FOISSNER, 1983) FOISSNER, AGATHA & BERGER, 2002

Genus Homalozoon STOKES, 1890*Homalozoon vermiculare* (STOKES, 1887) STOKES, 1890**Genus Myriokaryon** JANKOWSKI, 1973*Myriokaryon lieberkuehnii* (BÜTSCHLI, 1889) JANKOWSKI, 1973**Genus Papillorhabdos** FOISSNER, 1984*Papillorhabdos carchesii* FOISSNER, 1984**Genus Perispira** STEIN, 1859*Perispira pyriformis* WIRNSBERGER, FOISSNER & ADAM, 1984**Genus Pithothorax** KAHL, 1926*Pithothorax ovatus* (KAHL, 1926) KAHL, 1927**Genus Semispathidium** FOISSNER, AGATHA & BERGER, 2002*Semispathidium pulchrum* FOISSNER, HESS & AL-RASHEID, 2010**Genus Spathidium** DUJARDIN, 1841*Spathidium ampulliforme minuta* KALTENBACH, 1960*Spathidium anguilla* VUXANOVICI, 1962*Spathidium deforme* KAHL, 1928*Spathidium depressum* KAHL, 1930*Spathidium lieberkuehnii* BÜTSCHLI, 1889*Spathidium liepolti* KALTENBACH, 1960*Spathidium puteolagri* BAUMEISTER in KAHL, 1930*Spathidium spathula* (MÜLLER, 1773) ?DUJARDIN, 1841*Spathidium tortum* FOISSNER, 1980**Genus Teuthophrys** CHATTON & BEAUCHAMP, 1923*Teuthophrys trisulca trisulca* CHATTON & BEAUCHAMP, 1923**Genus Trachelophyllum** CLAPARÈDE & LACHMANN, 1859*Trachelophyllum apiculatum* (PERTY, 1852) CLAPARÈDE & LACHMANN, 1859*Trachelophyllum clavatum* STOKES, 1886*Trachelophyllum hyalinum* FOISSNER, 1983*Trachelophyllum sigmoides* KAHL, 1926*Trachelophyllum valkanovi* (LEPSI, 1959) FOISSNER, 1983*Trachelophyllum vestitum* STOKES, 1884**Subclass Trichostomatia****Order Cyclotrichida****Genus Askenasia** BLOCHMANN, 1895*Askenasia acrostomia* KRÄINER & FOISSNER, 1990*Askenasia chlorelligera* KRÄINER & FOISSNER, 1990*Askenasia volvox* (EICHWALD, 1852) BLOCHMANN, 1895**Genus Cyclotrichium** MEUNIER, 1910*Cyclotrichium viride* GAJEWSKAJA, 1933**Genus Mesodinium** STEIN, 1863*Mesodinium acarus* STEIN, 1867*Mesodinium pulex* (CLAPARÈDE & LACHMANN, 1859) STEIN, 1867

Genus Pelagovasicola JANKOWSKI, 1980*Pelagovasicola cinctus* (VOIGT, 1901) JANKOWSKI, 1980**Genus Rhabdoaskenasia** KRAINER & FOISSNER, 1990*Rhabdoaskenasia minima* KRAINER & FOISSNER, 1990**Order Pseudoholophryida****Genus Ovalorhabdos** FOISSNER, 1984*Ovalorhabdos sapropelica* FOISSNER, 1984**Genus Paraenchelys** FOISSNER, 1983*Paraenchelys spiralis* FOISSNER, 1983**Class Phyllopharyngea****Subclass Cyrtophoria****Order Chlamyodontida****Genus Alinostoma** JANKOWSKI, 1980*Alinostoma burkli* BLATTERER & FOISSNER, 1990

Remarks: In BLATTERER (2002) this species is quoted as *Alinostoma plurivacuolata*, because in some populations (n = 30) several specimens had a characteristic dorsal body which projected the ventral side, a feature used for the separation of *A. plurivacuolata* (DEROUX & DRAGESCO, 1968) JANKOWSKI, 1980 from *A. burkli* (BLATTERER & FOISSNER 1990). *Alinostoma* is very likely a synonym of *Pseudochilodonopsis* FOISSNER, 1979 (AESCHT 2001, p. 20).

Genus Chilodonella STRAND, 1928*Chilodonella cyprini* (MOROFF, 1902) KAHL, 1931*Chilodonella hexasticha* (KIERNIK, 1909) KAHL, 1931*Chilodonella labiata* (STOKES, 1891) KAHL, 1931*Chilodonella schewiakoffi* (SCHOUTEDEN, 1906) KAHL, 1931*Chilodonella uncinata* (EHRENBERG, 1838) STRAND, 1928**Genus Chlamydonella** PETZ, SONG & WILBERT, 1995*Chlamydonella alpestris* (FOISSNER, 1979) PETZ, SONG & WILBERT, 1995*Chlamydonella minuta* (PÄTSCH, 1974) PETZ, SONG & WILBERT, 1995*Chlamydonella rostrata* (VUXANOVICI, 1963) PETZ, SONG & WILBERT, 1995**Genus Chlamydonellopsis** BLATTERER & FOISSNER, 1990*Chlamydonellopsis plurivacuolata* BLATTERER & FOISSNER, 1990*Chlamydonellopsis polonica* (FOISSNER, CZAPIK & WIACKOWSKI, 1981) BLATTERER & FOISSNER, 1990**Genus Gastronauta** ENGELMANN in BÜTSCHLI, 1889*Gastronauta aloisi* OBERSCHMIDLEITNER & AESCHT, 1996*Gastronauta membranaceus* ENGELMANN in BÜTSCHLI, 1889**Genus Odontochlamys** CERTES, 1891*Odontochlamys alpestris* FOISSNER, 1981*Odontochlamys gouraudi* CERTES, 1891**Genus Paragastronauta** FOISSNER, 2001*Paragastronauta clatratus* (DEROUX, 1976) FOISSNER, 2001

Genus Phascolodon STEIN, 1859*Phascolodon vorticella* STEIN, 1859**Genus Pseudochilodonopsis** FOISSNER, 1979*Pseudochilodonopsis algivora* (KAHL, 1931) FOISSNER, 1979*Pseudochilodonopsis caudata* (PERTY, 1852) BLATTERER & FOISSNER, 1990*Pseudochilodonopsis fluviatilis* FOISSNER, 1988*Pseudochilodonopsis kloiberi* FOISSNER, 1979*Pseudochilodonopsis piscatoris* (BLOCHMANN, 1895) FOISSNER, 1979*Pseudochilodonopsis polyvacuolata* FOISSNER & DIDIER, 1981**Genus Thigmogaster** DEROUX, 1976*Thigmogaster oppositevacuolatus* AUGUSTIN & FOISSNER, 1989*Thigmogaster potamophilus* FOISSNER, 1988**Genus Trithigmostoma** JANKOWSKI, 1967*Trithigmostoma alpestris* FOISSNER, 1979*Trithigmostoma cucullulus* (MÜLLER, 1786) JANKOWSKI, 1967*Trithigmostoma marginatus* (SRÁMEK-HUSEK, 1957) FOISSNER, 1987*Trithigmostoma pituitosum* FOISSNER, 1979*Trithigmostoma srameki* FOISSNER, 1988*Trithigmostoma steini* (BLOCHMANN, 1895) FOISSNER, 1988**Genus Wilbertella** GONG & SONG, 2006*Wilbertella distyla* (WILBERT, 1971) GONG & SONG, 2006**Order Dysteriida****Genus Dysteria** HUXLEY, 1857*Dysteria fluviatilis* (STEIN, 1859) BLOCHMANN, 1895*Dysteria navicula* KAHL, 1928*Dysteria scutellum* WILBERT, 1971**Genus Orthotrochilia** SONG, 2003*Orthotrochilia agamalievi* (DEROUX, 1976) SONG, 2003**Genus Trochilia** DUJARDIN, 1841*Trochilia minuta* (ROUX, 1899) KAHL, 1931*Trochilia palustris* STEIN, 1859**Genus Trochilioides** JANKOWSKI, 2007*Trochilioides fimbriata* (FOISSNER, 1984) JANKOWSKI, 2007*Trochilioides recta* (KAHL, 1928) JANKOWSKI, 2007**Subclass Suctoria****Order Exogemmida****Genus Spirochona** STEIN, 1852*Spirochona gemmipara* STEIN, 1852**Order Exogenida****Genus Loricophrya** MATTHES, 1956*Loricophrya lauterborni* (SONDHEIM, 1929) CURDS, 1987

Genus Metacineta BÜTSCHLI, 1889*Metacineta mystacina* (EHRENBERG, 1831) BÜTSCHLI, 1889**Genus Mistarcon** JANKOWSKI, 1997*Mistarcon parasitica* (NOZAWA, 1939) JANKOWSKI, 1997**Genus Multifasciculatum** GOODRICH & JAHN, 1943*Multifasciculatum elongatum* (CLAPARÈDE & LACHMANN, 1859) JANKOWSKI, 1981**Genus Parapodophrya** KAHL, 1931*Parapodophrya soliformis* (LAUTERBORN, 1908) KAHL, 1931**Genus Podophrya** EHRENBERG, 1833*Podophrya fixa* (MÜLLER, 1786) EHRENBERG, 1833*Podophrya libera* PERTY, 1852*Podophrya niphargi* STROUHAL, 1939*Podophrya stylonychia* (KENT, 1882) MATTHES, 1971*Podophrya urostylae* (MAUPAS, 1881) JANKOWSKI, 1963**Genus Schizactinia** JANKOWSKI, 1967*Schizactinia multiramosa* (WENZEL, 1961) JANKOWSKI, 1967**Genus Sphaerophrya** CLAPARÈDE & LACHMANN, 1859*Sphaerophrya canelli* CLEMENT, 1967*Sphaerophrya epizoica* (HAMMANN, 1952) MATTHES, 1988*Sphaerophrya magna* MAUPAS, 1881*Sphaerophrya parurolepti* FOISSNER, 1980*Sphaerophrya stentoris* MAUPAS, 1881**Order Endogenida****Genus Acineta** EHRENBERG, 1833*Acineta compressa* CLAPARÈDE & LACHMANN, 1859*Acineta flava* KELLICOTT, 1885*Acineta fluviatilis* STOKES, 1885*Acineta tuberosa* EHRENBERG, 1834**Genus Dendrosoma** EHRENBERG, 1837*Dendrosoma radians* EHRENBERG, 1837**Genus Pseudogemma** COLLIN, 1909*Pseudogemma fraiponti* COLLIN, 1909**Genus Solenophrya** CLAPARÈDE & LACHMANN, 1859*Solenophrya crassa* CLAPARÈDE & LACHMANN, 1859**Genus Staurophrya** ZACHARIAS, 1893*Staurophrya elegans* ZACHARIAS, 1893**Genus Tokophrya** BÜTSCHLI, 1889*Tokophrya carchesii* (CLAPARÈDE & LACHMANN, 1859) BÜTSCHLI, 1889*Tokophrya cyclosum* (CLAPARÈDE & LACHMANN, 1859) BÜTSCHLI, 1889*Tokophrya infusionum* (STEIN, 1859) BÜTSCHLI, 1889*Tokophrya lemnarum* (STEIN, 1859) ENTZ, 1902*Tokophrya quadripartita* (CLAPARÈDE & LACHMANN, 1859) BÜTSCHLI, 1889

Genus Trichophrya CLAPARÈDE & LACHMANN, 1859*Trichophrya epistylidis* CLAPARÈDE & LACHMANN, 1859*Trichophrya melosirae* (GAJEWSKAJA, 1933) DOVGAL, 2002**Order Evaginogenida****Genus Dendrocometes** STEIN, 1852*Dendrocometes paradoxus* STEIN, 1852**Genus Discophrya** LACHMANN, 1859*Discophrya cothurnata* (WEISSE, 1847) LACHMANN, 1859*Discophrya cylindrica* (PERTY, 1852) COLLIN, 1912*Discophrya laccophili* MATTHES, 1954**Genus Enchelyomorpha** KAHL, 1930*Enchelyomorpha vermicularis* (SMITH, 1899) KAHL, 1930**Genus Heliophrya** SAEDELEER & TELLIER, 1930*Heliophrya minima* (RIEDER, 1936) FOISSNER, 1988*Heliophrya rotunda* (HENTSCHEL, 1916) MATTHES, 1954**Genus Periacineta** COLLIN, 1909*Periacineta buckei* (KENT, 1882) COLLIN, 1909**Genus Prodiscophrya** KORMOS, 1935*Prodiscophrya collini* (ROOT, 1914) KORMOS, 1935**Class Nassophorea****Order Synhymeniida****Genus Chilodontopsis** BLOCHMANN, 1895*Chilodontopsis depressa* (PERTY, 1852) BLOCHMANN, 1895*Chilodontopsis muscorum* KAHL, 1931*Chilodontopsis planicaudata* SONG & WILBERT, 1989*Chilodontopsis vorax* (STOKES, 1887) KAHL, 1931**Genus Nassulopsis** FOISSNER, BERGER & KOHMANN, 1994*Nassulopsis elegans* (EHRENBERG, 1833) FOISSNER, BERGER & KOHMANN, 1994*Nassulopsis paucivacuolata* (FOISSNER, 1979) FOISSNER, BERGER & KOHMANN, 1994**Genus Zosterodasys** DEROUX, 1978*Zosterodasys transversa* (KAHL, 1928) FOISSNER, BERGER & KOHMANN, 1994**Order Nassulida****Genus Furgasonia** JANKOWSKI, 1964*Furgasonia blochmanni* (FAURE-FREMIET, 1967) JANKOWSKI, 1964*Furgasonia rubens* (PERTY, 1849) FOISSNER, 1979*Furgasonia theresae* (FABRE-DOMERGUE, 1890) FOISSNER, AGATHA & BERGER, 2002**Genus Nassula** EHRENBERG, 1833*Nassula citrea* KAHL, 1931*Nassula lateritia* CLAPARÈDE & LACHMANN, 1859*Nassula longinassa* FOISSNER, 1980*Nassula ornata* EHRENBERG, 1833

Nassula rotunda GELEI, 1950

Genus Nassulides FOISSNER, AGATHA & BERGER, 2002

Nassulides vernalis (GELEI & SZABADOS, 1950) FOISSNER, AGATHA & BERGER, 2002

Genus Obertrumia FOISSNER & ADAM, 1981

Obertrumia aurea (EHRENBERG, 1833) FOISSNER, 1987

Obertrumia georgiana (DRAGESCO, 1972) FOISSNER & ADAM, 1981

Obertrumia gracilis FOISSNER, 1989

Order Microthoracida

Genus Drepanomonas FRESENIUS, 1858

Drepanomonas lunaris FOISSNER, 1979

Drepanomonas obtusa PENARD, 1922

Drepanomonas revoluta PENARD, 1922

Genus Leptopharynx MERMOD, 1914

Leptopharynx costatus MERMOD, 1914

Genus Microthorax ENGELMANN, 1862

Microthorax leptopharyngiformis FOISSNER, 1985

Microthorax pusillus ENGELMANN, 1862

Microthorax simplex FOISSNER, 1985

Microthorax simulans (KAHL, 1926) KAHL, 1931

Microthorax sulcatus ENGELMANN, 1862

Microthorax transversus FOISSNER, 1985

Microthorax tridentatus PENARD, 1922

Genus Pseudomicrothorax MERMOD, 1914

Pseudomicrothorax agilis MERMOD, 1914

Pseudomicrothorax dubius (MAUPAS, 1883) PENARD, 1922

Pseudomicrothorax foliformis FOISSNER, 1987

Genus Trochiliopsis PENARD, 1922

Trochiliopsis opaca PENARD, 1922

Class Colpodea

Order Bursariomorphida

Genus Bryometopus KAHL, 1932

Bryometopus chlorelligerus FOISSNER, 1980

Bryometopus edaphonus FOISSNER, 1980

Bryometopus sphagni (PENARD, 1922) KAHL, 1932

Genus Bursaria MÜLLER, 1773

Bursaria truncatella MÜLLER, 1773

Genus Bursaridium LAUTERBORN, 1894

Bursaridium pseudobursaria (FAURÉ-FREMIET, 1924) KAHL, 1927

Genus Paracondylostoma FOISSNER, 1980

Paracondylostoma setigerum FOISSNER, 1980

Genus Thylakidium SCHEWIAKOFF, 1892*Thylakidium pituitosum* FOISSNER, 1980*Thylakidium truncatum* SCHEWIAKOFF, 1892**Order Colpodida****Genus Bresslaua** KAHL, 1931*Bresslaua insidiatrix* CLAFF, DEWEY & KIDDER, 1941*Bresslaua vorax* KAHL, 1931**Genus Colpoda** MÜLLER, 1773*Colpoda cucullus* (MÜLLER, 1773) GMELIN, 1790*Colpoda distincta* (SMITH, 1899) FOISSNER, 1993*Colpoda ecaudata* (LIEBMANN, 1936) FOISSNER, BLATTERER, BERGER & KOHMANN, 1991*Colpoda flavicans* (STOKES, 1885) FOISSNER, 1993*Colpoda inflata* (STOKES, 1884) KAHL, 1931*Colpoda ovinucleata* FOISSNER, 1980*Colpoda magna* (GRUBER, 1880) LYNN, 1978*Colpoda steinii* MAUPAS, 1883*Colpoda variabilis* FOISSNER, 1980**Genus Hackenbergia** FOISSNER, 1997*Hackenbergia langae* FOISSNER, 1997**Genus Maryna** GRUBER, 1879*Maryna ovata* (GELEI, 1950) GELEI, 1954*Maryna socialis* GRUBER, 1879*Maryna umbrellata* (GELEI, 1950) FOISSNER, 1993**Genus Pseudochlamydonella** BUITKAMP, SONG & WILBERT, 1989*Pseudochlamydonella rheophila* BUITKAMP, SONG & WILBERT, 1989**Order Platyophryida****Genus Platyophrya** KAHL, 1926*Platyophrya citrina* FOISSNER, 1980*Platyophrya dubia* FOISSNER, 1980*Platyophrya hyalina* FOISSNER, 1980*Platyophrya sphagni* (PENARD, 1922) FOISSNER, 1993*Platyophrya vorax* KAHL, 1926**Genus Rostrophrya** FOISSNER, 1993*Rostrophrya camerounensis* (NJINE, 1979) FOISSNER, 1993**Order Cyrtolophosidida****Genus Cyrtolophosis** STOKES, 1885*Cyrtolophosis acuta* KAHL, 1926*Cyrtolophosis elongata* (SCHEWIAKOFF, 1892) KAHL, 1931*Cyrtolophosis mucicola* STOKES, 1885**Genus Kreyella** KAHL, 1931*Kreyella minuta* FOISSNER, 1979

Class Prostomatea**Order Prostomatida****Genus Apsiktrata** FOISSNER, BERGER & KOHMANN, 1994*Apsiktrata gracilis* (PENARD, 1922) FOISSNER, BERGER & KOHMANN, 1994**Genus Bursellopsis** CORLISS, 1960*Bursellopsis nigricans mobilis* (WANG & NIE, 1933) FOISSNER, BERGER & SCHAUMBURG, 1999*Bursellopsis spumosa* (SCHMIDT, 1920) CORLISS, 1960**Genus Vasicola** TATEM, 1869*Vasicola ciliata* TATEM, 1869*Vasicola lutea* KAHL, 1930**Order Prorodontida****Genus Balanion** WULFF, 1919*Balanion planctonicum* (FOISSNER, OLEKSIV & MÜLLER, 1990) FOISSNER, BERGER & KOHMANN, 1994**Genus Coleps** NITZSCH, 1827*Coleps elongatus* EHRENBERG, 1830*Coleps hirtus hirtus* (MÜLLER, 1786) NITZSCH, 1827*Coleps hirtus viridis* EHRENBERG, 1831*Coleps quadrispinus* FOISSNER, 1983*Coleps spetai* FOISSNER, 1984**Genus Holophrya** EHRENBERG, 1831*Holophrya coleps* EHRENBERG, 1831*Holophrya discolor* EHRENBERG, 1833*Holophrya nigricans* LAUTERBORN, 1894*Holophrya ovum* EHRENBERG, 1831*Holophrya saginata* PENARD, 1922*Holophrya teres* (EHRENBERG, 1833) FOISSNER, BERGER & KOHMANN, 1994**Genus Longifragma** FOISSNER, 1984*Longifragma obliqua* (KAHL, 1926) FOISSNER, 1984**Genus Malacophrys** KAHL, 1926*Malacophrys viridis* FOISSNER, 1980**Genus Nolandia** SMALL & LYNN, 1985*Nolandia nolandi* (KAHL, 1930) SMALL & LYNN, 1985**Genus Pantotrichum** EHRENBERG, 1830*Pantotrichum enchelys* EHRENBERG, 1831**Genus Parauotricha** FOISSNER, 1983*Parauotricha discolor* (KAHL, 1930) FOISSNER, 1983**Genus Pelagothrix** FOISSNER, BERGER & SCHAUMBURG, 1999*Pelagothrix plancticola* FOISSNER, BERGER & SCHAUMBURG, 1999**Genus Pinacocoleps** DIESING, 1866*Pinacocoleps incurvus* (EHRENBERG, 1833) DIESING, 1866

Genus Placus COHN, 1866*Placus luciae* (KAHL, 1926) KAHL, 1930*Placus ovum* (KAHL, 1926) KAHL, 1930**Genus Plagiocampa** SCHEWIAKOFF, 1892*Plagiocampa rouxi* KAHL, 1926**Genus Prorodon** EHRENBERG, 1833*Prorodon armatus* CLAPARÈDE & LACHMANN, 1859*Prorodon cinctus* FOISSNER, 1983*Prorodon ellipticus* (KAHL, 1930) FOISSNER, BERGER & KOHMANN, 1994*Prorodon niveus* EHRENBERG, 1833**Genus Urotricha** CLAPARÈDE & LACHMANN, 1859*Urotricha agilis* (STOKES, 1886) KAHL, 1930*Urotricha apsheronica* ALEKPEROV, 1984*Urotricha armata* KAHL, 1927*Urotricha castalia* MUNOZ, TELLEZ & FERNANDENZ-GALIANO, 1987*Urotricha corlissiana* SONG & WILBERT, 1989*Urotricha farcta* CLAPAREDE & LACHMANN, 1859*Urotricha furcata* SCHEWIAKOFF, 1892*Urotricha globosa* SCHEWIAKOFF, 1892*Urotricha lagenula* (EHRENBERG, 1831) KENT, 1881*Urotricha macrostoma* FOISSNER, 1983*Urotricha matthesi matthesi* KRAINER, 1995*Urotricha matthesi tristicha* FOISSNER & PFISTER, 1997*Urotricha ovata* KAHL, 1926*Urotricha pelagica* KAHL, 1935*Urotricha platystoma* STOKES, 1886*Urotricha psenneri* SONNTAG & FOISSNER, 2004*Urotricha pseudofurcata* KRAINER, 1995*Urotricha ristoii* KRAINER, 1995*Urotricha simonsbergeri* FOISSNER, BERGER & SCHAUMBURG, 1999*Urotricha spetai* FOISSNER, 2012*Urotricha venatrix* KAHL, 1935**Class Plagiopylea****Order Plagiopylida****Genus Plagiopyla** STEIN, 1860*Plagiopyla nasuta* STEIN, 1860**Order Trimyemida****Genus Discomorphella** CORLISS, 1960*Discomorphella pectinata* (LEVANDER, 1894) CORLISS, 1960**Genus Trimyema** LACKEY, 1925*Trimyema compressum* LACKEY, 1925

Order Odontostomatida**Genus Epalxella** CORLISS, 1960*Epalxella antiquorum* (PENARD, 1922) CORLISS, 1960*Epalxella bidens* (KAHL, 1932) CORLISS, 1960*Epalxella striata* (KAHL, 1926) CORLISS, 1960**Genus Mylestoma** KAHL, 1928*Mylestoma anatinum* (PENARD, 1922) KAHL, 1932**Genus Pelodinium** LAUTERBORN, 1908*Pelodinium reniforme* LAUTERBORN, 1908**Genus Saprodinium** LAUTERBORN, 1908*Saprodinium dentatum* (LAUTERBORN, 1901) LAUTERBORN, 1908*Saprodinium putrinium* LACKEY, 1925**Class Oligohymenophorea****Subclass Peniculia****Order Peniculida****Genus Clathrostoma** PENARD, 1922*Clathrostoma viminale* PENARD, 1922**Genus Disematostoma** LAUTERBORN, 1894*Disematostoma buetschlii* LAUTERBORN, 1894*Disematostoma colpidioides* GELEI, 1954**Genus Frontonia** EHRENBERG, 1833*Frontonia acuminata* (EHRENBERG, 1833) BÜTSCHLI, 1889*Frontonia angusta* KAHL, 1931*Frontonia atra* (EHRENBERG, 1833) BÜTSCHLI, 1889*Frontonia elliptica* BEARDSLEY, 1902*Frontonia leucas* (EHRENBERG, 1833) EHRENBERG, 1838*Frontonia rotunda* GELEI, 1954*Frontonia vernalis* (EHRENBERG, 1833) KAHL, 1931**Genus Lembadion** PERTY, 1849*Lembadion bullinum* (MÜLLER, 1786) PERTY, 1849*Lembadion lucens* (MASKELL, 1887) KAHL, 1931*Lembadion magnum* (STOKES, 1887) KAHL, 1931**Genus Marituja** GAJEWSKAJA, 1928*Marituja pelagica* GAJEWSKAJA, 1928**Genus Paramecium** MÜLLER, 1773*Paramecium aurelia*-Komplex*Paramecium bursaria* (EHRENBERG, 1831) FOCKE, 1836*Paramecium caudatum* EHRENBERG, 1833*Paramecium putrinum* CLAPARÈDE & LACHMANN, 1859**Genus Stokesia** WENRICH, 1929*Stokesia vernalis* WENRICH, 1929

Order Urocentrida**Genus Urocentrum** NITZSCH, 1827*Urocentrum turbo* (MÜLLER, 1786) NITZSCH, 1827**Subclass Scuticociliatia****Order Philasterida****Genus Balanonema** JANKOWSKI, 2007*Balanonema sapropelica* (FOISSNER, 1978) JANKOWSKI, 2007**Genus Cinetochilum** PERTY, 1849*Cinetochilum margaritaceum* (EHRENBERG, 1831) PERTY, 1849**Genus Dexiotricha** STOKES, 1885*Dexiotricha colpidiopsis* (KAHL, 1926) JANKOWSKI, 1964*Dexiotricha granulosa* (KENT, 1881) FOISSNER, BERGER & KOHMANN, 1994*Dexiotricha polystyla* FOISSNER, 1987*Dexiotricha tranquilla* (KAHL, 1926) AUGUSTIN & FOISSNER, 1992**Genus Dexiotrichides** KAHL, 1931*Dexiotrichides centralis* (STOKES, 1885) KAHL, 1931**Genus Kahlilembus** GROLIERE & COUTEAUX, 1984*Kahlilembus attenuatus* (SMITH, 1897) FOISSNER, BERGER & KOHMANN, 1994**Genus Loxocephalus** EBERHARD, 1862*Loxocephalus lucidus* SMITH, 1897*Loxocephalus luridus* EBERHARD, 1862**Genus Philasterides** KAHL, 1931*Philasterides armatus* (KAHL, 1926) KAHL, 1931**Genus Platynematum** FOISSNER, BERGER & KOHMANN, 1994*Platynematum sociale* (PENARD, 1922) FOISSNER, BERGER & KOHMANN, 1994**Genus Pseudocohnilembus** EVANS & THOMPSON, 1964*Pseudocohnilembus pusillus* (QUENNERSTEDT, 1869) FOISSNER & WILBERT, 1981*Pseudocohnilembus putrinus* (KAHL, 1928) FOISSNER & WILBERT, 1981**Genus Sathrophilus** CORLISS, 1960*Sathrophilus muscorum* (KAHL, 1931) CORLISS, 1960**Genus Uronema** DUJARDIN, 1841*Uronema biceps* PENARD, 1922*Uronema marinum* DUJARDIN, 1841*Uronema nigricans* (MÜLLER, 1786) FLORENTIN, 1901*Uronema parduczi* FOISSNER, 1971**Genus Uropedalium** KAHL, 1928*Uropedalium pyriforme* KAHL, 1928**Genus Urozona** SCHEWIAKOFF, 1889*Urozona buetschlii* SCHEWIAKOFF, 1889

Order Pleuronematida**Genus Calyptotricha** PHILLIPS, 1882*Calyptotricha chlorelligera* (LEPSI, 1957) FOISSNER, 1987*Calyptotricha lanuginosa* (PENARD, 1922) WILBERT & FOISSNER, 1980**Genus Conchophthirus** STEIN, 1861*Conchophthirus acuminatus* (CLAPARÈDE & LACHMANN, 1858) STEIN, 1861*Conchophthirus anodontae* (EHRENBERG, 1838) STEIN, 1861**Genus Cristigera** ROUX, 1899*Cristigera hammeri* WILBERT, 1986*Cristigera minor* PENARD, 1922*Cristigera phoenix* PENARD, 1922*Cristigera setosa* KAHL, 1928**Genus Ctedoctema** STOKES, 1884*Ctedoctema acanthocryptum* STOKES, 1884**Genus Cyclidium** MÜLLER, 1773*Cyclidium elongatum* CLAPARÈDE & LACHMANN, 1859*Cyclidium glaucoma* MÜLLER, 1773*Cyclidium heptatrichum* SCHEWIAKOFF, 1893*Cyclidium pellucidum* KAHL, 1931*Cyclidium versatile* PENARD, 1922**Genus Histiobalantium** STOKES, 1886*Histiobalantium bodamicum* KRÄINER & MÜLLER, 1995*Histiobalantium natans* (CLAPARÈDE & LACHMANN, 1859) KAHL, 1931**Genus Pleuronema** DUJARDIN, 1841*Pleuronema coronatum* KENT, 1881*Pleuronema crassum* DUJARDIN, 1841**Genus Protocyclidium** ALEKPEROV, 1993*Protocyclidium citrullus* (COHN, 1866) FOISSNER, AGATHA & BERGER, 2002**Subclass Hymenostomatia****Order Tetrahymenida****Genus Colpidium** STEIN, 1860*Colpidium colpoda* (LOSANA, 1829) STEIN, 1860*Colpidium kleini* FOISSNER, 1969**Genus Deltopylum** FAURÉ-FREMIET & MUGARD, 1946*Deltopylum rhabdoides* FAURE-FREMIET & MUGARD, 1946**Genus Dexiostoma** JANKOWSKI, 1967*Dexiostoma campylum* (STOKES, 1886) JANKOWSKI, 1967**Genus Dichilum** SCHEWIAKOFF, 1892*Dichilum platessoides* FAURÉ-FREMIET, 1924**Genus Epenardia** CORLISS, 1971*Epenardia myriophylli* (PENARD, 1922) CORLISS, 1971

Genus Espejoia BÜRGER, 1908*Espejoia culex* (SMITH, 1897) KAHL, 1931*Espejoia mucicola* (PENARD, 1922) KAHL, 1931**Genus Glaucoma** EHRENBERG, 1830*Glaucoma macrostoma* SCHEWIAKOFF, 1889*Glaucoma reniforme* SCHEWIAKOFF, 1892*Glaucoma scintillans* EHRENBERG, 1830*Glaucoma setosum* SCHEWIAKOFF, 1892**Genus Ichthyophthirius** FOUQUET, 1876*Ichthyophthirius multifiliis* FOUQUET, 1876**Genus Paracolpidium** GANNER & FOISSNER, 1989*Paracolpidium truncatum* (STOKES, 1885) GANNER & FOISSNER, 1989**Genus Spirozona** KAHL, 1926*Spirozona caudata* KAHL, 1926**Genus Stegophilum** SCHEWIAKOFF, 1892*Stegophilum schoenborni* FOISSNER, 1986**Genus Tetrahymena** FURGASON, 1940*Tetrahymena patula* (MÜLLER, 1786) CORLISS, 1951*Tetrahymena pyriformis*-Komplex**Genus Trichospira** ROUX, 1899*Trichospira inversa* (CLAPARÈDE & LACHMANN, 1859) KAHL, 1931**Genus Turaniella** CORLISS, 1960*Turaniella vitrea* (BRODSKY, 1925) CORLISS, 1960**Order Ophryoglenida****Genus Bursostoma** VÖRÖSVARY, 1950*Bursostoma bursaria* VÖRÖSVARY, 1950**Genus Ophryoglana** EHRENBERG, 1831*Ophryoglana flava* (EHRENBERG, 1833) CLAPARÈDE & LACHMANN, 1858*Ophryoglana flavicans* EHRENBERG, 1831*Ophryoglana hemophaga* MOLLOY, LYNN & GIAMBERINI, 2005*Ophryoglana inquieta* KAHL, 1931*Ophryoglana media* MUGARD, 1949**Subclass Apostomatia****Order Apostomatida****Genus Gymnodinioides** MINKIEWICZ, 1912*Gymnodinioides zonatum* (PENARD, 1922) CHATTON & LWOFF, 1935**Subclass Astomatia****Order Haptophryida****Genus Haptophrya** STEIN, 1867*Haptophrya planariorum* (SIEBOLD, 1839) STEIN, 1867

Genus Mesnilella CÉPÈDE, 1910*Mesnilella clavata* (LEIDY, 1855) CÉPÈDE, 1910**Subclass Peritrichia****Order Sessilida****Genus Apiosoma** BLANCHARD, 1885*Apiosoma piscicola* BLANCHARD, 1885*Apiosoma tintinnabulum* (KENT, 1881) ?STILLER, 1971**Genus Astylozoon** ENGELMANN, 1862*Astylozoon fallax* ENGELMANN, 1862*Astylozoon faurei* KAHL, 1935*Astylozoon vagans* (STILLER, 1939) DINGFELDER, 1962**Genus Campanella** GOLDFUSS, 1820*Campanella umbellaria* (LINNAEUS, 1758) GOLDFUSS, 1820**Genus Carchesium** EHRENBERG, 1831*Carchesium cyclopidarum* NENNINGER, 1948*Carchesium epistylis* CLAPARÈDE & LACHMANN, 1858*Carchesium polypinum* (LINNAEUS, 1758) EHRENBERG, 1830**Genus Cothurnia** EHRENBERG, 1831*Cothurnia annulata* STOKES, 1885*Cothurnia imberbis* EHRENBERG, 1831*Cothurnia patula* FROMENTEL, 1876*Cothurnia vaga* (SCHRANK, 1776) EHRENBERG, 1838**Genus Cyclodonta** MATTHES, 1958*Cyclodonta bipartita* (STOKES, 1885) MATTHES, 1958**Genus Daurotheca** JANKOWSKI, 1986*Daurotheca tespa* JANKOWSKI, 1986**Genus Epicarchesium** JANKOWSKI, 1985*Epicarchesium granulatum* (KELLCOTT, 1887) JANKOWSKI, 1985*Epicarchesium pectinatum* (ZACHARIAS, 1897) FOISSNER, BERGER & SCHAUMBURG, 1999**Genus Epistylis** EHRENBERG, 1830*Epistylis alpestris* FOISSNER, 1978*Epistylis anastatica* (LINNAEUS, 1767) EHRENBERG, 1831*Epistylis branchiophila* PERTY, 1852*Epistylis chrysemydis* BISHOP & JAHN, 1941*Epistylis coronata* NUSCH, 1970*Epistylis digitalis* (LINNAEUS, 1758) EHRENBERG, 1830*Epistylis entzii* STILLER, 1935*Epistylis galea* EHRENBERG, 1831*Epistylis hentscheli* KAHL, 1935*Epistylis kolbi* NENNINGER, 1948*Epistylis lacustris magna* NENNINGER, 1948*Epistylis niagarae* KELLCOTT, 1883*Epistylis nympharum* ENGELMANN, 1862

Epistylis plicatilis EHRENBERG, 1831

Epistylis procumbens ZACHARIAS, 1897

Epistylis pygmaeum (EHRENBERG, 1838) FOISSNER, BERGER & SCHAUMBURG, 1999

Epistylis sommerae SCHÖDEL, 1986

Epistylis variabilis STILLER, 1953

Genus Gerda CLAPARÈDE & LACHMANN, 1858

Gerda glans CLAPARÈDE & LACHMANN, 1858

Gerda picta (KENT, 1882) JANKOWSKI, 1976

Genus Hastatella ERLANGER, 1890

Hastatella radians ERLANGER, 1890

Hastatella aesculacantha JAROCKI & JAKUBOWSKA, 1927

Genus Heteropolaria FOISSNER & SCHUBERT, 1977

Heteropolaria lwoffii (FAURÉ-FREMIET, 1943) FOISSNER & SCHUBERT, 1977

Genus Intranstylum FAURE-FREMIET, 1904

Intranstylum eismondi KAHL, 1935

Intranstylum triformum SCHÖDEL, 1983

Genus Lagenophrys STEIN, 1852

Lagenophrys ampulla STEIN, 1852

Lagenophrys nassa STEIN, 1852

Lagenophrys vaginicola STEIN, 1852

Genus Opercularia GOLDFUSS, 1820

Opercularia archiorbopercularia FOISSNER, 1979

Opercularia articulata GOLDFUSS, 1820

Opercularia assellicola KAHL, 1935

Opercularia asymmetrica (BICZOK, 1956) AESCHT & FOISSNER, 1992

Opercularia coarctata (CLAPAREDE & LACHMANN, 1858) ROUX, 1901

Opercularia cylindrata WRZESNIEWSKI, 1866

Opercularia nutans (MÜLLER, 1773) STEIN, 1854

Opercularia venusta FOISSNER, 1979

Genus Ophrydium BORY, 1826

Ophrydium caudatum (PHILLIPS, 1883) ?FOISSNER, 1977

Ophrydium eutrophicum FOISSNER, 1979

Ophrydium hyalinum WRZESNIEWSKI, 1877

Ophrydium versatile (MÜLLER, 1786) BORY, 1824

Genus Opisthnecta FAURÉ-FREMIET, 1906

Opisthnecta bivacuolata FOISSNER, 1978

Opisthnecta dubia FOISSNER, 1975

Opisthnecta henneguyi FAURÉ-FREMIET, 1906

Opisthnecta minima FOISSNER, 1975

Genus Orbopercularia LUST, 1950

Orbopercularia nodosa FOISSNER, 1979

Genus Pelagovorticella JANKOWSKI, 1980*Pelagovorticella mayeri* (FAURÉ-FREMIET, 1923) JANKOWSKI, 1980*Pelagovorticella natans* (FAURÉ-FREMIET, 1924) JANKOWSKI, 1985**Genus Platycola** KENT, 1882*Platycola decumbens* (EHRENBERG, 1830) KENT, 1882*Platycola dilatata* (FROMENTEL, 1876) KENT, 1882**Genus Propyxidium** CORLISS, 1979*Propyxidium cothurnoides* (KENT, 1881) CORLISS, 1979**Genus Pseudocarchesium** SOMMER, 1951

Remarks: *Pseudocarchesium* was established by SOMMER (1951, p. 362, 396) with six species. However, since none of these species was fixed as type species, the genus *Pseudocarchesium* SOMMER, 1951 is invalid (ICZN 1999, Article 13.3). The subsequent designation of the originally included species *Carchesium aselli* ENGELMANN, 1862 as type species of *Pseudocarchesium* SOMMER, 1951 by JANKOWSKI (1994, p. 222) is likewise invalid, because for genera established after 1930 the type species has to be fixed in the original description (ICZN 1999, Articles 13.3, 68.2). AESCHT (2001, p. 135), who incorrectly assumed that SOMMER (1951) has established the genus *Pseudocarchesium* with only three species, has classified the genus as nomen nudum. Since the genus is not correctly published so far, the “*Pseudocarchesium*”-species recorded in Austria are listed according to their original combination. *Epistylis steinii* is often incorrectly assigned to PRECHT (1935, p. 442). However, he has only transferred the species from *Epistylis* to *Carchesium*. Perhaps *Carchesium steini* sensu PRECHT (1935, p. 442) is a misidentification.

Carchesium aselli ENGELMANN, 1862*Epistylis simulans* PLATE, 1888*Epistylis steinii* WRZESNIEWSKI, 1877*Pseudocarchesium ovatum* SOMMER, 1951**Genus Pseudohaplocaulus** WARREN, 1988*Pseudohaplocaulus infravacuolatus* FOISSNER & BROZEK, 1996**Genus Pseudovorticella** FOISSNER & SCHIFFMANN, 1974*Pseudovorticella chlamydophora* (PENARD, 1922) JANKOWSKI, 1976*Pseudovorticella difficilis magnistriata* FOISSNER & SCHIFFMANN, 1974*Pseudovorticella elongata* (FROMENTEL, 1876) LEITNER & FOISSNER, 1997*Pseudovorticella fasciculata* (MÜLLER, 1773) FOISSNER & BROZEK, 1996*Pseudovorticella margaritata* (FROMENTEL, 1876) JANKOWSKI, 1976*Pseudovorticella monilata* (TATEM, 1870) FOISSNER & SCHIFFMANN, 1974*Pseudovorticella mutans* (PENARD, 1922) JANKOWSKI, 1976*Pseudovorticella pseudocampanula* FOISSNER, 1979*Pseudovorticella quadrata* FOISSNER, 1979*Pseudovorticella sauwaldensis* FOISSNER & SCHIFFMANN, 1979*Pseudovorticella sphagni* FOISSNER & SCHIFFMANN, 1974**Genus Pyxicola** KENT, 1882*Pyxicola affinis* KENT, 1882*Pyxicola carteri* KENT, 1882*Pyxicola pusilla* KENT, 1882**Genus Rhabdostyla** KENT, 1881*Rhabdostyla inclinans* (MÜLLER, 1773) ROUX, 1901

Rhabdostyla dubia FOISSNER, 1979

Rhabdostyla longipes KENT, 1881

Genus Scyphidia DUJARDIN, 1841

Scyphidia limacina (MÜLLER, 1773) LACHMANN, 1856

Scyphidia physarum LACHMANN, 1856

Scyphidia rugosa DUJARDIN, 1841

Genus Telotrochidium KENT, 1881

Telotrochidium cylindricum FOISSNER, 1978

Telotrochidium elongatum FOISSNER, 1975

Telotrochidium johanninae FAURE-FREMIET, 1950

Genus Thuricola KENT, 1881

Thuricola folliculata (MÜLLER, 1786) KENT, 1881

Thuricola kellicottiana (STOKES, 1887) KAHL, 1935

Genus Usconophrys JANKOWSKI, 1985

Usconophrys aperta (PLATE, 1888) JANKOWSKI, 1985

Genus Vaginicola LAMARCK, 1816

Vaginicola crystallina EHRENBERG, 1830

Vaginicola ingenita (MÜLLER, 1786) LAMARCK, 1816

Vaginicola tinctoria EHRENBERG, 1830

Genus Vorticella LINNAEUS, 1767

Vorticella abbreviata KEISER, 1921

Vorticella alpestris FOISSNER, 1979

Vorticella campanula EHRENBERG, 1831

Vorticella chlorellata STILLER, 1940

Vorticella chlorostigma (EHRENBERG, 1831) EHRENBERG, 1838

Vorticella citrina MÜLLER, 1773

Vorticella constricta FROMENTEL, 1876

Vorticella convallaria-Komplex

Vorticella costata SOMMER, 1951

Vorticella cupifera KAHL, 1935

Vorticella extensa KAHL, 1935

Vorticella gracilis DUJARDIN, 1841

Vorticella kenti KAHL, 1935

Vorticella longifilum KENT, 1881

Vorticella microstoma-Komplex

Vorticella nutans MÜLLER, 1773

Vorticella octava-Komplex

Vorticella operculariformis FOISSNER, 1979

Vorticella picta (EHRENBERG, 1831) EHRENBERG, 1838

Vorticella putrina MÜLLER, 1776

Vorticella sepulcreti FOISSNER & SCHIFFMANN, 1975

Vorticella similis STOKES, 1887

Vorticella striata DUJARDIN, 1841

Vorticella utriculus STOKES, 1885

Vorticella vaga RÖMER, 1893

Vorticella vernalis STOKES, 1887

Genus Vorticellides FOISSNER, BLAKE, WOLF, BREINER & STOECK, 2009

Vorticellides aquadulcis (STOKES, 1887) FOISSNER, BLAKE, WOLF, BREINER & STOECK, 2009

Vorticellides astyliformis (FOISSNER, 1981) FOISSNER, BLAKE, WOLF, BREINER & STOECK, 2009

Vorticellides infusionum (DUJARDIN, 1841) FOISSNER, BLAKE, WOLF, BREINER & STOECK, 2009

Genus Zoothamnium BORY, 1826

Zoothamnium affine STEIN, 1854

Zoothamnium arbuscula (EHRENBERG, 1831) EHRENBERG, 1838

Zoothamnium aselli CLAPARÈDE & LACHMANN, 1858

Zoothamnium duplicatum KAHL, 1933

Zoothamnium elegans D'UDEKEM, 1864

Zoothamnium gammari KORFSMEIER, 1943

Zoothamnium kentii GRENFELL, 1884

Zoothamnium parasita STEIN, 1854

Zoothamnium procerius KAHL, 1935

Order Mobilida

Genus Pallitrichodina VAN AS & BASSON in AESCHT, 2001

Pallitrichodina stephani (VAN AS & BASSON, 1993) AESCHT, FOISSNER, BLATTERER, KOHMANN & BERGER, 2017

Remarks: The present species (original combination *Pallitrichodina stephani* VAN AS & BASSON, 1993) was established in the invalid genus *Pallitrichodina* VAN AS & BASSON, 1993 and was not combined with the valid genus *Pallitrichodina* VAN AS & BASSON in AESCHT, 2001 so far (for details on invalidity of *Pallitrichodina* VAN AS & BASSON, 1993 due to lack of type fixation, see AESCHT 2001, p. 113). This purely nomenclatural act was done in the German edition of the Fauna Aquatica Austriaca (AESCHT ET AL. 2017).

Genus Trichodina EHRENBERG, 1830

Trichodina domerguei megamicronucleata DOGIEL, 1940

Trichodina pediculus EHRENBERG, 1831

Genus Urceolariella CORLISS, 1977

Urceolariella mitra (SIEBOLD, 1850) CORLISS, 1977

Saprobic valencies

	S ¹	x	o	β	α	p	G	SI
Acinertia								
<i>A. incurvata</i> ²	p-i	-	-	-	3	7	4	3.7E
<i>A. uncinata</i>	a-p	-	-	2	4	4	2	3.2
Acineta								
<i>A. flava</i>	b	-	1	7	2	-	3	2.1
<i>A. grandis</i>	b-o	-	3	5	2	-	2	1.9
<i>A. tuberosa</i>	a-b	-	-	4	4	2	2	2.8
Actinobolina								
<i>A. radians</i>	b	-	1	7	2	-	3	2.1
<i>A. vorax</i>	o	-	7	3	-	-	4	1.3
Amphileptus								
<i>A. carchesii</i>	a	-	-	1	8	1	4	3.0
<i>A. claparedii</i>	a	-	-	2	8	-	4	2.8
<i>A. pleurosigma</i>	b-a	-	-	5	5	-	3	2.5
<i>A. procerus</i>	b-a	-	-	5	5	-	3	2.5
<i>A. punctatus</i>	a	-	-	1	9	-	5	2.9
Askenasia								
<i>A. volvox</i>	b	-	1	6	3	-	3	2.2
Aspidisca								
<i>A. cicada</i>	a-b	-	-	4	5	1	2	2.7
<i>A. lynceus</i>	b-a	-	1	4	4	1	1	2.5
<i>A. turrita</i>	a-b	-	-	4	6	-	3	2.6
Astylozoon								
<i>A. fallax</i>	b-a	-	-	5	5	-	3	2.5
<i>A. faurei</i>	b-a	-	-	5	5	-	3	2.5
Balanion								
<i>B. planctonicum</i>	o	-	7	3	-	-	4	1.3
Blepharisma								
<i>B. coeruleum</i>	b	-	2	8	-	-	4	1.8
<i>B. lateritium</i>	b	-	2	8	-	-	4	1.8
Bothrostoma								
<i>B. undulans</i> ²	p-m	-	-	-	-	10	5	4.0E
Brachonella								
<i>B. spiralis</i> ²	p-m	-	-	-	-	10	5	4.0E
Bursaria								
<i>B. truncatella</i>	b-a	-	2	4	3	1	1	2.3
Bursaridium								
<i>B. pseudobursaria</i>	o-b	-	6	4	-	-	3	1.4
Bursellopsis								
<i>B. spumosa</i>	o	-	7	3	-	-	4	1.3

	S ¹	x	o	β	α	p	G	SI
Caenomorpha								
<i>C. spp.</i> ²	p-m	-	-	-	-	10	5	4.0E
Calypotricha								
<i>C. lanuginosa</i>	a	-	-	3	7	-	4	2.7
Campanella								
<i>C. umbellaria</i>	a-b	-	-	3	6	1	3	2.8
Carchesium								
<i>C. pectinatum</i>	o-b	-	6	4	-	-	3	1.4
<i>C. polypinum</i>	a	-	-	2	7	1	3	2.9
Chaenea								
<i>C. stricta</i>	b-a	-	-	5	5	-	3	2.5
Chaetospira								
<i>C. muelleri</i>	b	-	1	8	1	-	4	2.0
<i>C. remex</i>	b-a	-	1	5	4	-	2	2.3
Chilodonella								
<i>C. uncinata</i>	a	-	-	2	6	2	3	3.0
Chilodontopsis								
<i>C. depressa</i>	b	-	1	7	2	-	3	2.1
Chlamydonella								
<i>C. alpestris</i>	b-a	-	2	4	4	-	2	2.2
Chlamydonellopsis								
<i>C. plurivacuolata</i>	b-a	-	-	5	5	-	3	2.5
Cinetochilum								
<i>C. margaritaceum</i>				eusaprobic				
Climacostomum								
<i>C. virens</i>	b	-	-	8	2	-	4	2.2
Codonella								
<i>C. cratera</i>	b-o	-	4	6	-	-	3	1.6
Coleps								
<i>C. hirtus</i>	a-b	-	1	3	4	2	1	2.7
<i>C. nolandi</i>	o-a	-	3	4	3	-	2	2.0
<i>C. spetai</i>	b	-	2	6	2	-	3	2.0
Colpidium								
<i>C. colpoda</i> ²	p-i	-	-	-	2	8	4	3.8E
<i>C. kleini</i>	p	-	-	-	3	7	4	3.7
Colpoda								
<i>C. cucullus</i>	p-a	-	-	-	4	6	3	3.6
<i>C. ecaudata</i> ²	p-i	-	-	-	1	9	5	3.9E
<i>C. inflata</i>	a-p	-	-	-	5	5	3	3.5
<i>C. magna</i>	a-p	-	-	2	5	3	2	3.1
<i>C. steinii</i> ³	a-p	-	-	-	5	5	3	3.5
<i>C. steinii</i> ⁴	b-a	-	2	4	3	1	1	2.3
Cothurnia								
<i>C. annulata</i>	o-b	-	6	4	-	-	3	1.4

	S ¹	x	o	β	α	p	G	SI
Ctedoctema								
<i>C. acanthocryptum</i>	b-a	-	1	4	4	1	1	2.5
Cyclidium								
<i>C. glaucoma</i>	a	-	-	1	7	2	3	3.1
<i>C. heptatrichum</i>	b	-	-	8	2	-	4	2.2
Cyrtolophosis								
<i>C. mucicola</i>	b-p	-	1	2	4	3	1	2.9
Dendrosoma								
<i>D. radians</i>	b-a	-	-	5	5	-	3	2.5
Dexiostoma								
<i>D. campylum</i> ²	p-i	-	-	-	1	9	5	3.9E
Dexiotricha								
<i>D. granulosa</i>	a-p	-	-	-	5	5	3	3.5
Dexiotrichides								
<i>D. centralis</i> ²	p-i	-	-	-	-	10	5	4.0E
Didinium								
<i>D. nasutum</i>	a-b	-	-	4	4	2	2	2.8
Dileptus								
<i>D. margaritifer</i>	b	-	2	5	3	-	2	2.1
Discomorphella								
<i>D. pectinata</i> ²	p-m	-	-	-	-	10	5	4.0E
Disematostoma								
<i>D. buetschlii</i>	b	-	1	7	2	-	3	2.1
<i>D. tetraedricum</i>	b	-	-	10	-	-	5	2.0
Drepanomonas								
<i>D. revoluta</i>	a-p	-	-	-	5	5	3	3.5
Dysteria								
<i>D. fluviatilis</i>	b	-	-	8	2	-	4	2.2
Enchelyodon								
<i>E. elegans</i>	a	-	-	-	10	-	5	3.0
Enchelyomorpha								
<i>E. vermicularis</i> ²	p-m	-	-	-	-	10	5	4.0E
Enchelys								
<i>E. gasterosteus</i>	b-a	-	-	5	5	-	3	2.5
Epalxella								
<i>E. spp.</i> ²	p-m	-	-	-	-	10	5	4.0E
Epenardia								
<i>E. myriophylli</i>	a-p	-	-	2	4	4	2	3.2
Epistylis								
<i>E. chrysemydis</i>	a	-	-	2	6	2	3	3.0
<i>E. coronata</i>	a	-	-	-	10	-	5	3.0
<i>E. digitalis</i>	o-b	-	5	5	-	-	3	1.5
<i>E. entzii</i>	a	-	-	2	7	1	3	2.9
<i>E. galea</i>	a	-	-	3	7	-	4	2.7
<i>E. hentscheli</i>	a-b	-	-	3	6	1	3	2.8

	S ¹	x	o	β	α	p	G	SI
<i>E. nympharum</i>	o-a	-	3	4	3	-	2	2.0
<i>E. plicatilis</i>	a-b	-	-	3	6	1	3	2.8
<i>E. procumbens</i>	o-b	-	5	5	-	-	3	1.5
Euplotes								
<i>E. aediculatus</i>	a	-	-	1	9	-	5	2.9
<i>E. affinis</i>	b-a	-	-	5	4	1	2	2.6
<i>E. eurystomus</i>	a	-	-	2	6	2	3	3.0
<i>E. moebiusi</i>	a	-	-	2	7	1	3	2.9
<i>E. patella</i>	b	-	-	7	3	-	4	2.3
Frontonia								
<i>F. acuminata</i>	b-a	-	2	4	4	-	2	2.2
<i>F. angusta</i>	b-a	-	-	5	5	-	3	2.5
<i>F. atra</i>	b-a	-	-	5	5	-	3	2.5
<i>F. leucas</i>	b-a	-	2	3	3	2	1	2.5
Gastronauta								
<i>G. clatratus</i>	b-a	-	2	4	4	-	2	2.2
<i>G. membranaceus</i>	b	-	2	6	2	-	3	2.0
Gastrostyla								
<i>G. mystacea</i>	p	-	-	-	3	7	4	3.7
<i>G. steinii</i>	a	-	-	2	7	1	3	2.9
Glaucoma								
<i>G. reniforme</i>	p	-	-	-	2	8	4	3.8
<i>G. scintillans</i> ⁵	p-a	-	-	-	4	6	3	3.6
Halteria								
<i>H. chlorelligera</i>	o	-	8	2	-	-	4	1.2
<i>H. grandinella</i>	b-a	-	1	6	3	-	3	2.2
Hastatella								
<i>H. radians</i>	b-a	-	1	6	3	-	3	2.2
Heliophrya								
<i>H. minima</i>	b-a	-	-	5	5	-	3	2.5
<i>H. rotunda</i>	b-a	-	-	5	5	-	3	2.5
Hexotricha								
<i>H. caudata</i> ²	p-m	-	-	-	-	10	5	4.0E
Histiculus								
<i>H. vorax</i>	a	-	-	-	10	-	5	3.0
Holophrya								
<i>H. discolor</i>	a-b	-	-	4	4	2	2	2.8
<i>H. ovum</i>	a-p	-	-	1	6	3	3	3.2
<i>H. teres</i>	b-p	-	-	3	4	3	2	3.0
Holosticha								
<i>H. kessleri</i>	a-b	-	-	4	5	1	2	2.7
<i>H. monilata</i>	a-b	-	-	3	6	1	3	2.8
<i>H. multistilata</i>	a-b	-	-	4	5	1	2	2.7
<i>H. pullaster</i>	b-a	-	1	4	4	1	1	2.5

	S ¹	x	o	β	α	p	G	SI
Homalozoon								
<i>H. vermiculare</i>	b-a	-	2	4	4	-	2	2.2
Hypotrichidium								
<i>H. conicum</i>	b-p	-	-	3	4	3	2	3.0
Kahlilembus								
<i>K. attenuatus</i>	b	-	-	10	-	-	5	2.0
Kerona								
<i>K. pediculus</i>	b-o	-	4	5	1	-	2	1.7
Lacrymaria								
<i>L. olor</i>	b	-	2	6	2	-	3	2.0
Lagenophrys								
<i>L. vaginicola</i>	o	-	9	1	-	-	5	1.1
Lagynophrya								
<i>L. acuminata</i>	o	-	8	2	-	-	4	1.2
Lagynus								
<i>L. elegans</i> ²	p-i	-	-	-	-	10	5	4.0E
Lembadion								
<i>L. bullinum</i>	b	-	-	9	1	-	5	2.1
<i>L. lucens</i>	b-a	-	-	6	4	-	3	2.4
<i>L. magnum</i>	b	-	2	8	-	-	4	1.8
Leptopharynx								
<i>L. costatus</i>	o-a	-	3	4	3	-	2	2.0
Linostoma								
<i>L. vorticella</i>	b-a	-	1	6	3	-	3	2.2
Litonotus								
<i>L. alpestris</i>	b-a	-	-	4	6	-	3	2.6
<i>L. crystallinus</i>	b-a	-	-	5	5	-	3	2.5
<i>L. cygnus</i>	b	-	-	10	-	-	5	2.0
<i>L. fusidens</i>	b-p	-	-	3	4	3	2	3.0
<i>L. lamella</i>	a	-	-	2	8	-	4	2.8
<i>L. varsaviensis</i> ^{2,6}	p-i	-	-	-	1	9	5	3.9E
<i>L. varsaviensis</i> ^{2,7}	b-a	-	-	5	5	-	3	2.5
Loxocephalus								
<i>L. luridus</i> ²	p-i	-	-	-	3	7	4	3.7E
Loxodes								
<i>L. spp.</i>	p	-	-	-	2	8	4	3.8
Loxophyllum								
<i>L. helus</i>	b	-	-	10	-	-	5	2.0
<i>L. meleagris</i>	b	-	-	8	2	-	4	2.2
<i>L. utriculariae</i>	b	-	1	8	1	-	4	2.0
Marituja								
<i>M. pelagica</i>	o	-	8	2	-	-	4	1.2
Mesodinium								
<i>M. acarus</i>	b	-	2	6	2	-	3	2.0
<i>M. pulex</i>	b	-	2	6	2	-	3	2.0

	S ¹	x	o	β	α	p	G	SI
Metacineta								
<i>M. cuspidata</i>	b-a	-	-	5	5	-	3	2.5
<i>M. mystacina</i>	b-a	-	-	5	5	-	3	2.5
Metopus								
<i>M. spp. sensu lato</i> ²	p-m	-	-	-	-	10	5	4.0E
Microthorax								
<i>M. pusillus</i>	a	-	-	2	8	-	4	2.8
Monilicaryon								
<i>M. monilatus</i>	b	-	-	7	3	-	4	2.3
Monodinium								
<i>M. balbianii</i>	o-a	-	3	3	4	-	2	2.1
Nassula								
<i>N. ornata</i>	b	-	2	6	2	-	3	2.0
<i>N. picta</i>	b	-	2	6	2	-	3	2.0
Nassulopsis								
<i>N. elegans</i>	b	-	1	8	1	-	4	2.0
Obertrumia								
<i>O. aurea</i>	b-a	-	-	6	4	-	3	2.4
Odontochlamys								
<i>O. alpestris</i>	b-a	-	-	5	5	-	3	2.5
Opercularia								
<i>O. articulata</i>	a-b	-	1	3	5	1	1	2.6
<i>O. coarctata</i>	a	-	-	2	7	1	3	2.9
<i>O. nutans</i>	b-a	-	-	5	5	-	3	2.5
Ophryidium								
<i>O. crassicaule</i>	b-a	-	-	5	5	-	3	2.5
<i>O. eutrophicum</i>	b-a	-	1	6	3	-	3	2.2
<i>O. sessile</i>	a-b	-	2	3	5	-	2	2.3
<i>O. versatile</i>	o	-	8	2	-	-	4	1.2
Ophryoglena								
<i>O. spp.</i>			not classified					
Opisthonecta								
<i>O. henneguyi</i>	b-p	-	-	3	4	3	2	3.0
Oxytricha								
<i>O. chlorelligera</i>	a	-	-	-	10	-	5	3.0
<i>O. fallax</i>	a	-	-	1	8	1	4	3.0
<i>O. ferruginea</i>	o	-	7	3	-	-	4	1.3
<i>O. haematoplasma</i>	b-a	-	-	6	4	-	3	2.4
<i>O. hymenostoma</i>	p	-	-	-	2	8	4	3.8
<i>O. saprobia</i>	a-p	-	-	-	6	4	3	3.4
<i>O. setigera</i>	a-b	-	-	4	6	-	3	2.6
<i>O. similis</i>	b-a	-	-	5	5	-	3	2.5
Paracolpidium								
<i>P. truncatum</i>	a	-	-	2	6	2	3	3.0

	S ¹	x	o	β	α	p	G	SI
Paradileptus								
<i>P. elephantinus</i>	b	-	3	6	1	-	3	1.8
Paramecium								
<i>P. aurelia</i> -Komplex	a-b	-	-	3	5	2	2	2.9
<i>P. bursaria</i>	b-a	-	-	6	3	1	3	2.5
<i>P. caudatum</i>	p-a	-	-	-	4	6	3	3.6
<i>P. putrinum</i>	p	-	-	1	2	7	3	3.6E
Parapodophrya								
<i>P. soliformis</i>	p	-	-	-	1	9	5	3.9
Paraurostyla								
<i>P. viridis</i>	b-a	-	-	5	5	-	3	2.5
<i>P. weissei</i>	a	-	-	2	7	1	3	2.9
Pelagohalteria								
<i>P. cirrifera</i>	o-b	-	6	4	-	-	3	1.4
Pelodinium								
<i>P. reniforme</i> ²	p-m	-	-	-	-	10	5	4.0E
Phascalodon								
<i>P. vorticella</i>	b-a	-	-	6	4	-	3	2.4
Phialina								
<i>P. spp.</i>				not classified				
Philasterides								
<i>P. armatus</i>	b-a	-	-	5	5	-	3	2.5
Placus								
<i>P. luciae</i>	b-o	-	4	4	2	-	2	1.8
Plagiocampa								
<i>P. rouxi</i>	a-b	-	-	4	6	-	3	2.6
Plagiopyla								
<i>P. nasuta</i> ²	p-i	-	-	-	-	10	5	4.0E
Platycola								
<i>P. decumbens</i>	b-a	-	2	4	4	-	2	2.2
Platynematum								
<i>P. sociale</i>	p	-	-	-	3	7	4	3.7
Platyophrya								
<i>P. vorax</i> ²	p-i	-	-	-	-	10	5	4.0E
Pleuronema								
<i>P. coronatum</i>	b	-	-	7	3	-	4	2.3
<i>P. crassum</i>	b-a	-	2	4	3	1	1	2.3
Pleurotricha								
<i>P. grandis</i>	b	-	-	10	-	-	5	2.0
Podophrya								
<i>P. fixa</i>	a	-	-	1	7	2	3	3.1
<i>P. maupasii</i>	a	-	-	1	9	-	5	2.9
Prodiscophrya								
<i>P. collini</i>	a-p	-	-	1	5	4	2	3.3

	S ¹	x	o	β	α	p	G	SI
Prorodon								
<i>P. ellipticus</i>	b-a	-	-	5	5	-	3	2.5
<i>P. niveus</i>	b-o	-	3	6	1	-	3	1.8
Pseudoblepharisma								
<i>P. tenue</i>	p	-	-	-	3	7	4	3.7
Pseudochilonopsis								
<i>P. algivora</i> ³	a	-	-	-	10	-	5	3.0
<i>P. algivora</i> ⁴	a-b	-	-	5	5	-	3	2.5
<i>P. fluviatilis</i>	b-a	-	-	5	3	2	2	2.7
<i>P. piscatoris</i>	b	-	-	7	3	-	4	2.3
Pseudocohnilembus								
<i>P. pusillus</i> ²	p-i	-	-	-	3	7	4	3.7E
Pseudomicrothorax								
<i>P. agilis</i>	b	-	1	8	1	-	4	2.0
Pseudovorticella								
<i>P. chlamydothora</i>	b-a	-	-	5	5	-	3	2.5
<i>P. monilata</i>	b-a	-	1	5	4	-	2	2.3
Pyxicola								
<i>P. carteri</i>	o-b	-	5	5	-	-	3	1.5
Rhabdostyla								
<i>R. inclinans</i>	a	-	-	-	10	-	5	3.0
Saprodinium								
<i>S. spp.</i> ²	p-m	-	-	-	-	10	5	4.0E
Sathrophilus								
<i>S. muscorum</i>	b-a	-	-	5	5	-	3	2.5
Scyphidia								
<i>S. rugosa</i>	a	-	-	-	8	2	4	3.2
Spathidium								
<i>S. sensu lato</i>				not classified				
Sphaerophrya								
<i>S. magna</i>	p	-	-	-	2	8	4	3.8
Spirostomum								
<i>S. ambiguum</i>	a	-	-	2	6	2	3	3.0
<i>S. caudatum</i>	o-b	-	6	4	-	-	3	1.4
<i>S. minus</i>	a-b	-	-	3	6	1	3	2.8
<i>S. teres</i>	p	-	-	1	2	7	3	3.6
Staurophrya								
<i>S. elegans</i>	o-a	-	3	4	3	-	2	2.0
Steinia								
<i>S. platystoma</i>	b-a	-	-	6	4	-	3	2.4
Stentor								
<i>S. amethystinus</i>	b	-	2	6	2	-	3	2.0
<i>S. coeruleus</i>	a-b	-	-	4	6	-	3	2.6
<i>S. igneus</i>	b	-	-	7	3	-	4	2.3
<i>S. muelleri</i>	b-a	-	-	5	5	-	3	2.5

	S ¹	x	o	β	α	p	G	SI
<i>S. multiformis</i>	b-a	-	-	5	5	-	3	2.5
<i>S. niger</i>	o-b	-	6	4	-	-	3	1.4
<i>S. polymorphus</i>	b-a	-	-	5	5	-	3	2.5
<i>S. roeselii</i>	a-b	-	1	4	5	-	2	2.4
Sterkiella								
<i>S. histriomuscorum</i>	a	-	-	2	6	2	3	3.0
Stichotricha								
<i>S. aculeata</i>	b-a	-	1	5	4	-	2	2.3
<i>S. secunda</i>	o	-	7	3	-	-	4	1.3
Stokesia								
<i>S. vernalis</i>	b	-	3	7	-	-	4	1.7
Strobilidium								
<i>S. caudatum</i>	o-b	-	5	5	-	-	3	1.5
<i>S. humile</i>	b	-	2	8	-	-	4	1.8
Strombidium								
<i>S. viride</i>	b	-	1	8	1	-	4	2.0
Stylonychia								
<i>S. mytilus</i> -Komplex	a	-	-	1	9	-	5	2.9
<i>S. pustulata</i>	b	-	1	7	2	-	3	2.1
<i>S. putrina</i>	a	-	-	2	7	1	3	2.9
<i>S. stylomuscorum</i>	b	-	-	10	-	-	5	2.0
<i>S. vorax</i>	b	-	-	10	-	-	5	2.0
Tachysoma								
<i>T. bicirratum</i>	a-p	-	-	2	4	4	2	3.2
<i>T. pellionellum</i>	b-a	-	1	4	4	1	1	2.5
Tetrahymena								
<i>T. pyriformis</i> -Komplex ²	p-i	-	-	-	3	7	4	3.7E
Thigmogaster								
<i>T. oppositevacuolatus</i>	a-b	-	-	3	5	2	2	2.9
<i>T. potamophilus</i>	b-a	-	-	5	5	-	3	2.5
Thuricola								
<i>T. folliculata</i>	b	-	2	6	2	-	3	2.0
<i>T. kellicottiana</i>	b	-	2	7	1	-	3	1.9
<i>T. vasiformis</i>	a	-	-	-	10	-	5	3.0
Tintinnidium								
<i>T. fluviatile</i>	o-b	-	5	5	-	-	3	1.5
<i>T. pusillum</i>	b	-	-	8	2	-	4	2.2
<i>T. semiciliatum</i>	b	-	2	6	2	-	3	2.0
Tintinnopsis								
<i>T. cylindrata</i>	b	-	-	7	3	-	4	2.3
Tokophrya								
<i>T. carchesii</i>	a	-	-	2	7	1	3	2.9
<i>T. infusionum</i>	b-a	-	2	5	3	-	2	2.1
<i>T. lemnarum</i>	a	-	-	1	7	2	3	3.1
<i>T. quadripartita</i>	a-b	-	-	3	5	2	2	2.9

	S ¹	x	o	β	α	p	G	SI
Trachelius								
<i>T. ovum</i>	a-b	-	1	4	4	1	1	2.5
Trachelophyllum								
<i>T. apiculatum</i>	b-a	-	-	5	5	-	3	2.5
Trichodina								
<i>T. pediculus</i>	b	-	2	6	2	-	3	2.0
Trimyema								
<i>T. compressum</i> ²	p-m	-	-	-	2	8	4	3.8E
Trithigmostoma								
<i>T. cucullulus</i>	a-p	-	-	2	5	3	2	3.1
<i>T. srameki</i>	b-a	-	1	6	3	-	3	2.2
<i>T. steini</i>	b-a	-	1	6	3	-	3	2.2
Trochilia								
<i>T. minuta</i>	b-a	-	-	5	5	-	3	2.5
Trochilioides								
<i>T. recta</i>	a	-	-	-	10	-	5	3.0
Tropidoattractus								
<i>T. acuminatus</i> ²	p-m	-	-	-	-	10	5	4.0E
Urocentrum								
<i>U. turbo</i>	a-b	-	-	4	4	2	2	2.8
Uroleptus								
<i>U. gallina</i>	b	-	-	10	-	-	5	2.0
<i>U. musculus</i>	a	-	-	1	8	1	4	3.0
<i>U. piscis</i>	a	-	-	3	7	-	4	2.7
<i>U. rattulus</i>	b	-	-	10	-	-	5	2.0
Uronema								
<i>U. nigricans</i>	a-p	-	-	1	6	3	3	3.2
Urostyla								
<i>U. grandis</i>	a	-	-	3	7	-	4	2.7
Urotricha								
<i>U. agilis</i>	b-a	-	-	5	5	-	3	2.5
<i>U. armata</i>	a	-	-	2	8	-	4	2.8
<i>U. farcta</i>	a-b	-	-	4	6	-	3	2.6
<i>U. furcata</i>	b	-	2	6	2	-	3	2.0
<i>U. globosa</i>	b	-	-	7	3	-	4	2.3
<i>U. ovata</i>	a-p	-	-	-	6	4	3	3.4
Urozoa								
<i>U. buetschlii</i> ²	p	-	-	-	2	8	4	3.8E
Vaginicola								
<i>V. ingenita</i>	b	-	2	6	2	-	3	2.0
<i>V. tincta</i>	o-b	-	5	5	-	-	3	1.5
Vorticella								
<i>V. aquadulcis</i> -Komplex	b-a	-	2	5	3	-	2	2.1
<i>V. campanula</i>	a-b	-	1	4	5	-	2	2.4
<i>V. convallaria</i> -Komplex	a	-	1	2	6	1	2	2.7

	S ¹	x	o	β	α	p	G	SI
<i>V. fromenteli</i>	a	-	-	2	8	-	4	2.8
<i>V. infusionum</i> -Komplex ²	p-a	-	-	1	4	5	2	3.4E
<i>V. marginata</i>	b	-	2	8	-	-	4	1.8
<i>V. mayeri</i>	b	-	-	10	-	-	5	2.0
<i>V. microstoma</i> -Komplex ²	p-a	-	-	-	5	5	3	3.5E
<i>V. natans</i>	b	-	3	7	-	-	4	1.7
<i>V. octava</i> -Komplex	b-a	-	2	4	4	-	2	2.2
<i>V. picta</i>	b	-	2	6	2	-	3	2.0
Zoothamnium								
<i>Z. arbuscula</i>	b-a	-	1	6	3	-	3	2.2
<i>Z. kentii</i>	b-a	-	-	5	5	-	3	2.5
<i>Z. procerius</i>	b-a	-	-	5	5	-	3	2.5
Zosterodasys								
<i>Z. transversa</i>	b	-	1	7	2	-	3	2.1

1: The verbal classification of the indicator organisms (S) is not uniform in SLADCEK et al. (1981) as well as other publications. The corrections herein are made without comment.

2: Species marked with an "E" are also classified within the table "eusaprobity".

3: If found as very abundant or in masses.

4: For single finds or occurring with low abundance.

5: Only useful as an indicator if it occurs with moderate abundance.

6: This animal is so categorized when it is found in low to mass abundances and the biological river water quality class corresponds to one with chemical loading.

7: This animal is so categorized when it is found in low to moderately high abundances and the biological river water quality class corresponds to one with chemical loading.

Definition of higher saprobic levels see introduction Edition 2002, page 41/42

	Eusaprobity							SI
	S ¹	α	p	i	m	h	G	
Acinertia								
<i>A. incurvata</i>	a-i	3	4	3	-	-	2	4.0
Bothrostoma								
<i>B. undulans</i>	m-i	-	1	4	5	-	2	5.4
Brachonella								
<i>B. spiralis</i>	m-i	-	1	4	5	-	2	5.4
Caenomorphia								
<i>C. spp.</i>	m-i	-	1	4	5	-	2	5.4
Colpidium								
<i>C. colpoda</i>	p-i	2	5	3	-	-	2	4.1
Colpoda								
<i>C. ecaudata</i>	p-i	1	6	3	-	-	3	4.2
Dexiostoma								
<i>D. campylum</i>	p-i	1	6	3	-	-	3	4.2
Dexiotrichides								
<i>D. centralis</i>	p-i	-	5	5	-	-	3	4.5
Discomorphella								
<i>D. pectinata</i>	m-i	-	1	4	5	-	2	5.4
Enchelyomorpha								
<i>E. vermicularis</i>	m-i	-	1	3	6	-	3	5.5
Epalxella								
<i>E. spp.</i>	m-i	-	1	4	5	-	2	5.4
Hexotricha								
<i>H. caudata</i>	p-m	-	3	4	3	-	2	5.0
Lagynus								
<i>L. elegans</i>	p-i	-	5	5	-	-	3	4.5
Litonotus								
<i>L. varsaviensis</i>	p-i	1	5	4	-	-	2	4.3
Loxocephalus								
<i>L. luridus</i>	a-i	3	4	3	-	-	2	4.0
Metopus								
<i>M. spp. sensu lato</i>	m-i	-	1	4	5	-	2	5.4
Paramecium								
<i>P. putrinum</i>	a-i	3	4	3	-	-	2	4.0
Pelodinium								
<i>P. reniforme</i>	m-i	-	1	4	5	-	2	5.4
Plagiopyla								
<i>P. nasuta</i>	p-i	-	5	5	-	-	3	4.5
Platyophrya								
<i>P. vorax</i>	p-i	-	5	5	-	-	3	4.5

	S ¹	α	p	i	m	h	G	SI
Pseudocohnilembus								
<i>P. pusillus</i>	a-i	3	4	3	-	-	2	4.0
Saprodinium								
<i>S. spp.</i>	m-i	-	1	4	5	-	2	5.4
Tetrahymena								
<i>T. pyriformis</i> -Komplex	a-i	3	4	3	-	-	2	4.0
Trimyema								
<i>T. compressum</i>	p-m	2	2	3	3	-	1	4.7
Tropidoattractus								
<i>T. acuminatus</i>	m-i	-	1	4	5	-	2	5.4
Urozona								
<i>U. buetschlii</i>	p-i	2	4	4	-	-	2	4.2
Vorticella								
<i>V. infusionum</i> -Komplex	a	5	2	2	1	-	1	3.9
<i>V. microstoma</i> -Komplex	a-p	5	3	2	-	-	2	3.7

¹: The verbal classification of the indicator organisms (S) is not uniform in SLADCEK et al. (1981) as well as other publications. The corrections herein are made without comment.

	Biomass		Occurrence				Community ³	Saprobity ⁴
	(mg of 10 ⁶ ind. ¹⁾)	Main food	Salinity tolerance ²	Preferred water type	Preferred habitat			
Acinertia								
<i>A. incurvata</i>	55	R	he	F,S,K	A,B	COL,HBE	p-i	
<i>A. uncinata</i>	10	R	os	F,S,K	A,B	COL,NBE	a-p	
Acineta								
<i>A. flava</i>	30	R	oe?	F,S	A,T		b	
<i>A. grandis</i>	150	R	oe?	F,S	A,T		b-o	
<i>A. tuberosa</i>	20	R	he	S,F,K	A,T		a-b	
Actinobolina								
<i>A. radians</i>	125	R	oe?	S,F	P,A		b	
<i>A. vorax</i>	250	R	oms?	S	P		o	
Amphileptus								
<i>A. carchesii</i>	200	R	os	S,F	A	CAR	a	
<i>A. claparedii</i>	60	R	he?	S,F	A	CAR	a	
<i>A. pleurosigma</i>	150	R	oms	S,F	A,B	STE	b-a	
<i>A. procerus</i>	160-1500	R	os	S,F	B		b-a	
<i>A. punctatus</i>	80	R	os	S,F	A,B		a	
Askenasia								
<i>A. volvox</i>	35	Al,Ki	oe?	S,F	P	OLI	b	
Aspidisca								
<i>A. cicada</i>	10	Ba	he?	F,S,K	B,A	TRI,CYR,NBE	a-b	
<i>A. lynceus</i>	17	Ba	ome?	F,S,K	B,A	TRI,CYR,NBE	b-a	
<i>A. turrita</i>	7	Ba	he	F,S,K	B,A	NBE	a-b	
Astylozoon								
<i>A. fallax</i>	30	Ba	os	S	P	MAR	b-a	
<i>A. faurei</i>	50	Ba	oms?	S,F	P	MAR	b-a	
Balanion								
<i>B. planctonicum</i>	0.3-3.6	Al	os	S	P	OLI	o	

	Biomass		Occurrence				Community ³	Saprobity ⁴
	(mg of 10 ⁶ ind. ¹⁾)	Main food	Salinity tolerance ²	Preferred water type	Preferred habitat			
Blepharisma								
<i>B. coeruleum</i>	250	Al (O)	os	S,F	B		b	
<i>B. lateritium</i>	250	Ba,Al	os	S	B,P		b	
Bursaria								
<i>B. truncatella</i>	50000	O	ome?	S,F	B,P		b-a	
Bursaridium								
<i>B. pseudobursaria</i>	342	Al	os	S,F	P		o-b	
Bursellopsis								
<i>B. spumosa</i>	18000	O	os	S,F	P		o	
Caenomorpha								
<i>C. spp.</i>	120 ⁵	Ba,Sb	os	S,F,K	Fs	MET	p-m	
Calyptricha								
<i>C. lanuginosa</i>	5	Ba,Al,Fl	ome	S,F	B,A	TRI	a	
Campanella								
<i>C. umbellaria</i>	850	Ba	oms	S,F	A,B,T	CAR	a-b	
Carchesium								
<i>C. pectinatum</i>	60	Ba?	he?	S,F	P		o-b	
<i>C. polypinum</i>	150	Ba	oe	F,S,K	B,A,T	TRI,CAR,NBE	a	
Chaenea								
<i>C. stricta</i>	10	Ba	os	F,S	B,A		b-a	
Chaetospira								
<i>C. muelleri</i>	80	Ba,Ki,Fl	he	S,F	B,A		b	
<i>C. remex</i>	250	Ba,Fl,Ki	oe	S,F	A,B		b-a	
Chilodonella								
<i>C. uncinata</i>	11	Ba	he?	F,S,K,Bo	A,B	TRI,CYR,NBE	a	
Chilodontopsis								
<i>C. depressa</i>	10	Ba,Al,Ki	he	F,S	A,B	PLE,CYR	b	

	Biomass		Occurrence				
	(mg of 10 ⁶ ind. ¹⁾)	Main food	Salinity tolerance ²	Preferred water type	Preferred habitat	Community ³	Saprobity ⁴
Chlamydonella							
<i>C. alpestris</i>	3	Ki,Ba	os	F,S,Bo	A,B	CYR	b-a
Chlamydonellopsis							
<i>C. plurivacuolata</i>	50	Ki	os	F	A,B	STE,CYR	b-a
Cinetochilum							
<i>C. margaritaceum</i>	5	Ba,Al	ome (he?)	S,F	A,B,P	NBE	10
Climacostomum							
<i>C. virens</i>	500	O	he?	S,F	B,P	MOO	b
Codonella							
<i>C. cratera</i>	20	Ki,Al?	oe	S,F	P	OLI	b-o
Coleps							
<i>C. hirtus</i>	21	O	oms (he?)	S,F	A,B,P	STE	a-b
<i>C. nolandi</i>	16	O	he	S,F	A,B	STE	o-a
<i>C. spetai</i>	60	Al,Cy	os	S	P		b
Colpidium							
<i>C. colpoda</i>	130	Ba,Fl,Al	ome	F,S,K	B	COL,TRI,HBE	p-i
<i>C. kleini</i>	65	Ba	os	F,S	B	TRI	p
Colpoda							
<i>C. cucullus</i>	70-140	Ba,Fl,Al	ome?	Bo,S,F	B,A	BOD	p-a
<i>C. ecaudata</i>	5-10	Ba	ome	Bo,K	B	HBE	p-i
<i>C. inflata</i>	40	Ba,Fl	ome?	Bo,S	B		a-p
<i>C. magna</i>	2400	Ba (O)	os	S	B	MAR	a-p
<i>C. steinii</i>	4	Ba	ome	Bo,S,F	P	BOD	a-p ⁹
Cothurnia							
<i>C. annulata</i>	14	Ba	oe?	S,F	A		o-b
Ctedoctema							
<i>C. acanthocryptum</i>	2	Ba	os	S,F	B	STE	b-a

	Biomass		Occurrence				
	(mg of 10 ⁶ ind. ¹⁾)	Main food	Salinity tolerance ²	Preferred water type	Preferred habitat	Community ³	Saprobity ⁴
Cyclidium							
<i>C. glaucoma</i>	1-3	Ba	he	F,S,K	B,A,P	TRI	a
<i>C. heptatrichum</i>	2	Ba	ome?	F,S	B,A,P		b
Cyrtolophosis							
<i>C. mucicola</i>	2	Ba	he?	Bo,S,F	B	BOD	b-p
Dendrosoma							
<i>D. radians</i>	.7	R	oms	S,F	A,B,T		b-a
Dexiostoma							
<i>D. campylum</i>	26	Ba,Fl,Al	oms	F,S,K	B	COL,HBE	p-i
Dexiotricha							
<i>D. granulosa</i>	20	Ba	oe (he?)	S,F	B,A	NBE	a-p
Dexiotrichides							
<i>D. centralis</i>	5	Ba	oe?	S,F,K	Fs,B		p-i
Didinium							
<i>D. nasutum</i>	500	R	oe?	S,F	B,P		a-b
Dileptus							
<i>D. margaritifer</i>	500	O	he	S,F	B,A	PLE	b
Discomorphella							
<i>D. pectinata</i>	50	Ba,Sb	he?	S,F	Fs	MET	p-m
Disematostoma							
<i>D. buetschlii</i>	400	Al,Ba	os	S	P	MAR	b
<i>D. tetraedricum</i>	150	Ki	os	S,F	P	MAR	b
Drepanomonas							
<i>D. revoluta</i>	1	Ba	oms?	Bo,S,F,K	B,A,P	BOD	a-p
Dysteria							
<i>D. fluviatilis</i>	5	?	os	F,S	A,B		b
Enchelyodon							
<i>E. elegans</i>	200	R?	oe?	S,F	B,A		a

	Biomass (mg of 10 ⁶ ind. ¹⁾	Main food	Salinity tolerance ²	Preferred water type	Occurrence Preferred habitat	Community ³	Saprobity ⁴
Enchelyomorpha							
<i>E. vermicularis</i>	3	-	oms?	K,F,S	Fs,B	MET,HBE	p-m
Enchelys							
<i>E. gasterosteus</i>	21	O	os	S,F	B,A	STE	b-a
Epalxella							
<i>E. spp.</i>	2-25	Sb	os	S,F	Fs	MET	p-m
Epenardia							
<i>E. myriophylli</i>	700	Ba,Sb	os	S,F	B		a-p
Epistylis							
<i>E. chrysemydis</i>	260-1300	Ba,Al	oe	F	A,T	CAR,NBE	a
<i>E. coronata</i>	90	Ba	os	S	A		a
<i>E. digitalis</i>	30	Ba?	os	S,F	T		o-b
<i>E. entzii</i>	300	Ba	os	S,F,K	A,T	CAR,NBE	a
<i>E. galea</i>	250	Ba	os	S,F	A		a
<i>E. hentscheli</i>	100	Ba	oms	F,S	A,B	CAR,NBE	a-b
<i>E. nympharum</i>	80	Ba	os	S,F	T		o-a
<i>E. plicatilis</i>	40	Ba	ome?	S,F,K	A,B,T	CAR,NBE	a-b
<i>E. procumbens</i>	40	Ba,Fl	oe?	S,F	P		o-b
Euplotes							
<i>E. aediculatus</i>	260	O	oe?	F,S,K	B	CYR,NBE	a
<i>E. affinis</i>	18	Ba,Ki,Al, Fl	he?	F,S,K	B,A	TRI,STE,CYR, NBE	b-a
<i>E. eurystomus</i>	400	O	he?	S,F	B		a
<i>E. moebiusi</i>	23	Ba,Ki,Fl	he	F,S,K	B,A	CYR,NBE	a
<i>E. patella</i>	93	O	he?	F,S,K	B,A	PLE,CYR,NBE	b
Frontonia							
<i>F. acuminata</i>	100	O	oms	S,F	B,A,P	STE	b-a
<i>F. angusta</i>	110	O	os	F,S	B,A,P	STE	b-a

	Biomass		Salinity tolerance ²	Preferred water type	Occurrence		Community ³	Saprobity ⁴
	(mg of 10 ⁶ ind. l ⁻¹)	Main food			Preferred habitat	Habitat		
<i>F. atra</i>	95	Ki	os?	S,F	B,P		b-a	
<i>F. leucas</i>	270	O	oe	S,F	B,A,P	STE	b-a	
Gastronauta								
<i>G. clatratus</i>	10	Ki	oms	F,S	A,B	PLECYR	b-a	
<i>G. membranaceus</i>	15	Ba	oe?	F,S	A,B	PLE,CYR	b	
Gastrostyla								
<i>G. mystacea</i>	120	O	os	S,Bo	B		p	
<i>G. steinii</i>	122	O	os	S,F,Bo	B		a	
Glaucoma								
<i>G. reniforme</i>	10	Ba,Al	os	S,F	B,A		p	
<i>G. scintillans</i>	25	Ba	oe	F,S,K	B,A	COL,TRI	p-a	
Halteria								
<i>H. chlorelligera</i>	45	Al	os	S,F	P,Fs?		o?	
<i>H. grandinella</i>	27	Ba,Al	he?	S,F,Bo	P,B	OLI,MAR	b-a	
Hastatella								
<i>H. radians</i>	30	Ba	oe?	S,F	P	MAR	b-a	
Heliophrya								
<i>H. minima</i>	10	R	os	S,F	A		b-a	
<i>H. rotunda</i>	40	R	oe?	S,F	A		b-a	
Hexotricha								
<i>H. caudata</i>	5	Ba	oms?	S,F,K	Fs		p-m	
Histriculus								
<i>H. vorax</i>	350	Ba	os	S,K	B		a	
Holophrya								
<i>H. discolor</i>	290	O	he	S,F,K	B,P		a-b	
<i>H. ovum</i>	400	Ba,Cy,Al	oms	S,F	B,P		a-p	
<i>H. teres</i>	1300	O	he	S,F	B,P		b-p	

	Biomass (mg of 10 ⁶ ind. ¹⁾	Main food	Salinity tolerance ²	Preferred water type	Occurrence Preferred habitat	Community ³	Saprobity ⁴
Holosticha							
<i>H. kessleri</i>	66	Ba,Ki	pe	S,F	B		a-b
<i>H. monilata</i>	52	Ba,Ki,Al	ome	F,S	B	STE,MOO	a-b
<i>H. multistilata</i>	109	O	ome	F,S,Bo	B		a-b
<i>H. pullaster</i>	12	Ba,Ki,Al	he	F,S	B	STE,CYR	b-a
Homalozoon							
<i>H. vermiculare</i>	300	O	oe	S,F	B,A		b-a
Hypotrichidium							
<i>H. conicum</i>	150	O	oms?	S	P	MAR	b-p
Kahlilembus							
<i>K. attenuatus</i>	3	Ba	he?	S,F,Bo	B,A	BOD	b
Kerona							
<i>K. pediculus</i>	230	Al,Ki ³	os	S,F	T,P		b-o
Lacrymaria							
<i>L. olor</i>	33	R	he	S,F	B,A	PLE	b
Lagenophrys							
<i>L. vaginicola</i>	40	Ba	os	S	T		o
Lagynophrya							
<i>L. acuminata</i>	25	Al	os	S	P		o
Lagynus							
<i>L. elegans</i>	200	O	he	S,F	Fs,B	MET	p-i
Lembadion							
<i>L. bullinum</i>	200	O	oe?	S,F	B	PLE	b
<i>L. lucens</i>	40	O	oms	S,F	B,P	STE	b-a
<i>L. magnum</i>	120	O	os	S,F	B,P	PLE	b
Leptopharynx							
<i>L. costatus</i>	5	Ba,Al	os	Bo,S,F	B,A,P	BOD,MOO	o-a

	Biomass			Occurrence			
	(mg of 10 ⁶ ind. ¹⁾)	Main food	Salinity tolerance ²	Preferred water type	Preferred habitat	Community ³	Saprobity ⁴
Linostoma							
<i>L. vorticella</i>	1000	O	oe?	S,F	P,B		b-a
Litonotus							
<i>L. alpestris</i>	2	Ba?,Fl?	os	F,S	B,A	STE,CYR	b-a
<i>L. crystallinus</i>	13-100	R?	os	S,F	B,A		b-a
<i>L. cygnus</i>	40	R	he	F,S	B,A	PLE,CYR	b
<i>L. fusidens</i>	20-80	R	he?	S,F	B,A		b-p
<i>L. lamella</i>	15	R	he?	F,S,K	B,A	TRI	a
<i>L. varsaviensis</i>	60	R	he?	F,S	B,A	COL	p-i
Loxocephalus							
<i>L. luridus</i>	300	Ba	oe	S,F	B,A,Fs		p-i
Loxodes							
<i>L. magnus</i>	960	O	os	S,F	B,P	MET	p
<i>L. rostrum</i>	250	O	oms	S,F	B,P	MET	p
<i>L. striatus</i>	200	Al,Ki,Cy	os	S,F	B,P	MET	p
Loxophyllum							
<i>L. helus</i>	160	R	he	S,F	A,B		b
<i>L. meleagris</i>	700	R	he?	S,F	A,B	PLE	b
<i>L. utriculariae</i>	90	R	oe?	F,S	A		b
Marituja							
<i>M. pelagica</i>	190	Ki,Cy,Al (O)	os	S	P		o
Mesodinium							
<i>M. acarus</i>	1.5	O	he	S,F	P,B		b
<i>M. pulex</i>	5	O	he	S,F	P,B		b
Metacineta							
<i>M. cuspidata</i>	16	R	os	S,F	A		b-a
<i>M. mystacina</i>	65	R	ome	S,F	A,T		b-a

	Biomass		Salinity tolerance ²	Preferred water type	Occurrence		Community ³	Saprobity ⁴
	(mg of 10 ⁶ ind. ¹)	Main food			Preferred habitat			
Metopus								
<i>M. spp. sensu lato</i>	15-500	Ba,Fl,Al	he	S,F,K	Fs	MET,HBE		p-m
Microthorax								
<i>M. pusillus</i>	1	Ba	he	S,F	B,A			a
Monilicaryon								
<i>M. monilatus</i>	900	O	os	S,F	B,A	PLE		b
Monodinium								
<i>M. balbianii</i>	55	R	he?	S,F	P,B	OLI		o-a
Nassula								
<i>N. ornata</i>	1600	Cy	oms	S,F	B,A,P	MAR		b
<i>N. picta</i>	224	Cy (O)	oe?	S,F,Bo	B,A,P	MAR		b
Nassulopsis								
<i>N. elegans</i>	400	Cy	he?	S,F	B,P	MAR		b
Obertrumia								
<i>O. aurea</i>	500	Cy	he?	S,F	B,P	MAR		b-a
Odontochlamys								
<i>O. alpestris</i>	10	Ba	os	F,Bo	A,B	CYR		b-a
Opercularia								
<i>O. articulata</i>	140	Ba	os	F,S,K	A,T	CAR,STE,NBE		a-b
<i>O. coarctata</i>	25	Ba	os	F,K	A,B	CAR,NBE		a
<i>O. nutans</i>	70	Ba	os	S,F	A,T	CAR,STE,NBE		b-a
Ophrydium								
<i>O. crassicaule</i>	180	Ba,Al	oms	S	A			b-a
<i>O. eutrophicum</i>	215	Ba	os	S	A,P			b-a
<i>O. sessile</i>	350	Ba	oe?	S	A			a-b
<i>O. versatile</i>	280	Ba,Al	he?	S	A,P			o
Ophryoglena								
<i>O. spp.</i>	-	histophag	-	S,F	B			-

	Biomass		Occurrence				Community ³	Saprobity ⁴
	(mg of 10 ⁶ ind. ¹)	Main food	Salinity tolerance ²	Preferred water type	Preferred habitat			
Opisthionecta								
<i>O. henneguyi</i>	1000	Ba,Fl	os	S,F,K	P,B	MAR	b-p	
Oxytricha								
<i>O. chlorelligera</i>	35	Ba,Fl,Ki	oms	S,F	B,A		a	
<i>O. fallax</i>	155	O	he?	S,F	B		a	
<i>O. ferruginea</i>	125	Ba,Cy,Al,Ki	oe?	F,S	B		o	
<i>O. haematoplasma</i>	80	O	os	F,S	B	STE	b-a	
<i>O. hymenostoma</i>	30	O	os	F,S,K	B,A		p	
<i>O. saprobia</i>	34	Ba,Fl	os	S,F	B		a-p	
<i>O. setigera</i>	8	Ba,Fl	os	F,S,Bo	B		a-b	
<i>O. similis</i>	14	Ba	he?	F,S	B		b-a	
Paracolpidium								
<i>P. truncatum</i>	30	Ba	os	F,S	B		a	
Paradileptus								
<i>P. elephantinus</i>	1000	O	os	S	P	OLI	b	
Paramecium								
<i>P. aurelia</i> -Komplex	150	Ba	ome	S,F,K	B,P	TRI,CAR	a-b	
<i>P. bursaria</i>	120	Ba,Al,Ki	ome	S,F	A,B,P	STE,MOO	b-a	
<i>P. caudatum</i>	500	Ba,Al	ome	S,F,K	B,P	COL,TRI,HBE	p-a	
<i>P. putrinum</i>	70	Ba,Sb,Cy,Fl	ome	F,S,K	B,A,P	COL,HBE	p	
Parapodophrya								
<i>P. soliformis</i>	65	R	oms?	S,K	Fs	HBE	p	
Paraurostyla								
<i>P. viridis</i>	87	Ba	os	S	B		b-a	
<i>P. weissei</i>	240	O	ome?	S,F	B		a	
Pelagohalteria								
<i>P. cirrifera</i>	35	Al	os	S,F	P		o-b	

	Biomass		Salinity tolerance ²	Preferred water type	Occurrence		Community ³	Saprobity ⁴
	(mg of 10 ⁶ ind. ¹)	Main food			Preferred habitat			
Pelodinium								
<i>P. reniforme</i>	20	Sb	he?	S,F	Fs	MET		p-m
Phascolodon								
<i>P. vorticella</i>	75	Al,Ki	oe	S,F	P	OLI,MAR		b-a
Phialina								
<i>P. spp.</i>	-	R	-	S,F,Bo	B,A	-		-
Philasterides								
<i>P. armatus</i>	25	histophag	os	S,F	B,A			b-a
Placus								
<i>P. luciae</i>	25	O	ome	S,F	B,A	PLE		b-o
Plagiocampa								
<i>P. rouxi</i>	7	Ba,Al	he	S,F,Bo	B,A,P			a-b
Plagiopyla								
<i>P. nasuta</i>	120	Ba,Sb,Al,Fl	oe?	S,F	Fs	MET		p-i
Platycola								
<i>P. decumbens</i>	35	Ba,Al,Fl	ome	S,F	A			b-a
Platynematum								
<i>P. sociale</i>	4	Ba	ome	S,F	B,A			p
Platyophrya								
<i>P. vorax</i>	5-12	O	os	Bo,S,F	B	BOD		p-i
Pleuronema								
<i>P. coronatum</i>	60	O	he?	S,F	B	PLE		b
<i>P. crassum</i>	60	Ba,Al,Ki	he	S,F	B			b-a
Pleurotricha								
<i>P. grandis</i>	1300	Ki,Al	oms?	S,F	B			b
Podophrya								
<i>P. fixa</i>	50	R	he?	S,F,K	A,B	NBE		a
<i>P. maupasii</i>	30-110	R	he	S,F	A,B	NBE		a

	Biomass		Occurrence				Community ³	Saprobity ⁴
	(mg of 10 ⁶ ind. ¹)	Main food	Salinity tolerance ²	Preferred water type	Preferred habitat			
Prodiscophrya								
<i>P. collini</i>	78	R	os	S,F,K	A,B	COL,NBE	a-p	
Prorodon								
<i>P. ellipticus</i>	190	R	he?	S,F	B,A		b-a	
<i>P. niveus</i>	2500	R	oms?	S,F	B		b-o	
Pseudoblepharisma								
<i>P. tenue</i>	30	Ba	os	S,Fs,F	B		p	
Pseudochilodonopsis								
<i>P. algivora</i>	9	Al,Ba	he?	S,F	B,P	CYR	a ⁸	
<i>P. fluviatilis</i>	15	Ki	os	F,K	A,B	STE,CYR	b-a	
<i>P. piscatoris</i>	19	Al,Ki	os	S,F	A	CYR	b	
Pseudocohnilembus								
<i>P. pusillus</i>	6	Ba	he	S,F,K,Bo	B,P	MET,COL	p-i	
Pseudomicrothorax								
<i>P. agilis</i>	14	Cy (Ba,Al)	oe?	S,F	A,B		b	
Pseudovorticella								
<i>P. chlamydophora</i>	50	Ba,Al	ome	S,F	A,B		b-a	
<i>P. monilata</i>	70	Ba	ome?(he?)	S,F	A,B	STE	b-a	
Pyxicola								
<i>P. carteri</i>	20	Ba	os	S	A		o-b	
Rhabdostyla								
<i>R. inclinans</i>	35	Ba	oms?	S,F	T		a	
Saprodinium								
<i>S. spp.</i>	17-50	Ba,Sb	os	S,F,K	Fs	MET	p-m	
Sathrophilus								
<i>S. muscorum</i>	12	Ba,Fl	os	Bo,S,F	A	BOD	b-a	
Scyphidia								
<i>S. rugosa</i>	90	Ba?	os	S,F	A,B		a	

	Biomass		Occurrence				
	(mg of 10 ⁶ ind. ¹)	Main food	Salinity tolerance ²	Preferred water type	Preferred habitat	Community ³	Saprobity ⁴
Spathidium							
<i>S. s. l.</i>	-	R	-	S,F	A,B,P	BOD	-
Sphaerophrya							
<i>S. magna</i>	65	R	he	S,F	A,B,P		p
Spirostomum							
<i>S. ambiguum</i>	14600	Ba,Fl,Al	oe	S,F	B,P	TRI	a
<i>S. caudatum</i>	130	Ba	he	S	B		o-b
<i>S. minus</i>	425	Ba	oe?	S,F	B,P	STE	a-b
<i>S. teres</i>	380	Sb,Ba,Al,Ki	oe (he?)	S,F,	B,P,Fs	COL,HBE	p
Staurophrya							
<i>S. elegans</i>	110	R	oe?	S,F	P		o-a
Steinia							
<i>S. platystoma</i>	75	O	os	S,F	A,B		b-a
Stentor							
<i>S. amethystinus</i>	4000	Ba,Al,Ki	os	S	P		b
<i>S. coeruleus</i>	12000	O	oe	S,F	B,A,P		a-b
<i>S. igneus</i>	450	Ba,Al,Ki	os	S,F	B,P	PLE	b
<i>S. muelleri</i>	4500	Ba,Al,Ki	ome	S,F	A	STE	b-a
<i>S. multiformis</i>	600	Al,Ba	he	S,F	B,A	STE	b-a
<i>S. niger</i>	1000	Al	oms	S,F	A,B		o-b
<i>S. polymorphus</i>	4500	O	oms	S,F	B,A	STE	b-a
<i>S. roeselii</i>	5000	O	oe	S,F	B,A	STE	a-b
Sterkiella							
<i>S. histriomuscorum</i>	72	O	os	F,S,K,Bo	B	NBE	a
Stichotricha							
<i>S. aculeata</i>	20	Ba,Al	he?	S,F	B	MAR	b-a
<i>S. secunda</i>	30	Ba,Al,Ki	ome	S,F	B,A	MAR	o

	Biomass (mg of 10 ⁶ ind. 1)	Main food	Salinity tolerance ²	Preferred water type	Occurrence Preferred habitat	Community ³	Saprobity ⁴
Stokesia							
<i>S. vernalis</i>	400	Ba,Al,Ki	os	S,F	P		b
Strobilidium							
<i>S. caudatum</i>	45	Ki,Al,Ba	oms?	S,F	B,P	PLE	o-b
<i>S. humile</i>	4	Ki	oms?	S	P,B	OLI	b
Strombidium							
<i>S. viride</i>	50	Ki,Al,Ba	oe	S,F	P	OLI	b
Stylonychia							
<i>S. mytilus</i> -Komplex	400	O	ome	S,F	B,A	TRI,CYR	a
<i>S. pustulata</i>	80	O	he?	S,F	B,A	CYR	b
<i>S. putrina</i>	68	O	ome	S,F	B		a
<i>S. stylomuscorum</i>	30	Ki,Fl	os	F	B		b
<i>S. vorax</i>	57	O	os	S	B		b
Tachysoma							
<i>T. bicirratum</i>	15	Ba,Al	os	S,F	B		a-p
<i>T. pellionellum</i>	15	Ba,Cy,Al,Ki	ome (he?)	F,S	B,A	STE,CYR	b-a
Tetrahymena							
<i>T. pyriformis</i> -Komplex	15	Ba ⁶	oms?	F,S,K	B	COL	p-i
Thigmogaster							
<i>T. oppositovacuolatus</i>	15	Ba	os	F,K	A,B	CYR	a-b
<i>T. potamophilus</i>	2.5	Ki,Al	os	F	A,B	CYR	b-a
Thuricola							
<i>T. folliculata</i>	120	Ba,Al	he	S,F	A		b
<i>T. kellicottiana</i>	200	Al	oms?	S,F	A		b
<i>T. vasiformis</i>	130	Ba	os	S	A,B		a
Tintinnidium							
<i>T. fluviatile</i>	50	Al,Ki	oe	S,F	P	OLI	o-b
<i>T. pusillum</i>	40	Al,Ki,Ba	oms?	S,F	P	OLI	b

	Biomass		Salinity tolerance ²	Preferred water type	Occurrence		Saprobity ⁴
	(mg of 10 ⁶ ind. ¹)	Main food			Preferred habitat	Community ³	
<i>T. semiciliatum</i>	40	Al,Ki	os	S,F	A,B	PLE	b
Tintinnopsis							
<i>T. cylindrata</i>	20	Al	os	S,F	P	OLI	b
Tokophrya							
<i>T. carchesii</i>	12	R	os	S,F	T	CAR	a
<i>T. infusionum</i>	30	R	os	S,F	A,B	CAR,NBE	b-a
<i>T. lemnarum</i>	16	R	oms?	S,F,K	A,B,T	CAR,NBE	a
<i>T. quadripartita</i>	75	R	oms?	S,F,K	A,B,T	CAR,NBE	a-b
Trachelius							
<i>T. ovum</i>	3000	R	oms	F,S	A,P	CAR	a-b
Trachelophyllum							
<i>T. apiculatum</i>	39	O	he?	S,F	A,B		b-a
Trichodina							
<i>T. pediculus</i>	80	Ba ⁴	he?	S,F	T,P		b
Trimyema							
<i>T. compressum</i>	10	Ba	he	S,F,K	Fs	MET,COL,HBE	p-m
Trithigmostoma							
<i>T. cucullulus</i>	50	Ki,Al,Cy,Ba	he?	F,S,K	A,B	COL,TRI,CYR	a-p
<i>T. srameki</i>	40	Ki	os	F,S	A,B	STE,CYR	b-a
<i>T. steini</i>	150	Ki	os	F,S	A,B	CYR	b-a
Trochilia							
<i>T. minuta</i>	1.5	Ba	os	F,K	A,B	STE,CYR	b-a
Trochilioides							
<i>T. recta</i>	25	Sb	he	F,S	A,B,Fs		a
Tropidoatractus							
<i>T. acuminatus</i>	20	Ba	os	S	Fs		p-m
Urocentrum							
<i>U. turbo</i>	70	Ba,Ki	he?	S,F	B,A,P		a-b

	Biomass		Occurrence				Saprobity ⁴
	(mg of 10 ⁶ ind. ¹)	Main food	Salinity tolerance ²	Preferred water type	Preferred habitat	Community ³	
Uroleptus							
<i>U. gallina</i>	72	Al	oms?	S,F	B		b
<i>U. musculus</i>	214	O	oms?	S,F	B,A		a
<i>U. piscis</i>	400	Ba,Cy,Ki	oe?	S,F	B,A		a
<i>U. rattulus</i>	400	Ba,Al	oe?	S,F	B,A		b
Uronema							
<i>U. nigricans</i>	5	Ba,Fl	he	F,S	B,A,P	TRI	a-p
Urostyla							
<i>U. grandis</i>	500	O	he?	S,F	B		a
Urotricha							
<i>U. agilis</i>	0.5	Ba,Fl	os	S	B,P	OLI,MAR	b-a
<i>U. armata</i>	15	R	oe (he?)	S,F	B,A	MAR	a
<i>U. farcta</i>	5	Ba,Al,Fl	oms?	S,F	B,P	OLI,MAR	a-b
<i>U. furcata</i>	3-4	Ba,Al	os	S,F	P	OLI,MAR	b
<i>U. globosa</i>	7	Ba,Al	he?	S	P	OLI,MAR	b
<i>U. ovata</i>	15	Al	oe?	S,F	B,P	OLI,MAR,MOO	a-p
Urozona							
<i>U. buetschlii</i>	3	Ba	os	S,K,F	B,P		p
Vaginicola							
<i>V. ingenita</i>	3-4	Ba	he	S,F	A,T		b
<i>V. tincta</i>	15	Ba	os	S,F	A		o-b
Vorticella							
<i>V. aquadulcis</i> -Komplex	15	Ba,Al	he?	S,F,K	A,B	STE	b-a
<i>V. campanula</i>	135	Ba,Al	oe (he?)	S,F,K	A,B,T	STE	a-b
<i>V. convallaria</i> -Komplex	50-75	Ba	he	S,F,K	A,B,T	TRI,CAR,NBE	a
<i>V. fromenteli</i>	35	Ba	oe	S	A		a
<i>V. infusionum</i> -Komplex	25	Ba	he?	S,F,K,Bo	A,B,T	COL,CAR,HBE	p-a
<i>V. marginata</i>	100	Ba	os	S,F	A,B		b

	Biomass		Salinity tolerance ²	Occurrence		Community ³	Saprobity ⁴
	(mg of 10 ⁶ ind. ¹)	Main food		Preferred water type	Preferred habitat		
<i>V. mayeri</i>	50	Ba	os	S,F	P		b
<i>V. microstoma</i> -Komplex	30	Ba,Al	oms?	S,F	A,B		p-a
<i>V. natans</i>	90	Ba,Al	oe?	S,F	P	OLI	b
<i>V. octava</i> -Komplex	20	Ba	oe	S,F	A		b-a
<i>V. picta</i>	40	Ba,Al	oe?	S,F	A	PLE	b
Zoothamnium							
<i>Z. arbuscula</i>	55	Ba	ome?	S,F	A		b-a
<i>Z. kentii</i>	40	Ba	ome	F,S	A	CAR,STE	b-a
<i>Z. procerius</i>	45	Ba	he	F,S	A,B,T	CAR,STE	b-a
Zosterodasys							
<i>Z. transversa</i>	300	Ki	he	F,S	A,B	CYR	b

a = alpha-mesosaprobic, **A** = Aufwuchs (periphyton), **Al** = algae (except of diatoms, but inclusively autotrophic flagellates), **b** = beta-meso-saprobic, **B** = benthos, **Ba** = bacteria, **Bo** = terrestrial soils, **BOD** = influence of soil and/or moss, **CAR** = *Carchesietosum polyipinae*, **COL** = *Colpidietum colpodae*, **Cy** = cyanobacteria, **CYR** = *Cyrtophoretea*, **F** = flowing waters, **FI** = heterotrophic flagellates, **Fs** = anaerobic mud (and anaerobic zones in the pelagial), **HBE** = high-load and/or oxygen deficient activated sludge, **he** = holo-euryhaline, **i** = isosaprobic, **K** = sewage treatment works (activated sludge plants), **Ki** = diatoms, **MAR** = *Marynetum*, **MET** = *Metopetum*, **MOO** = mire influence, **mpe** = meso- to poly-euryhaline, **mpe** = meso- to poly-stenohaline, **NBE** = normale activated sludge, **O** = omnivorous (feeds on autotrophic organisms and protozoans, sometimes even on small metazoans), **o** = oligosaprobic, **oe** = oligo-euryhaline, **OLI** = *Oligotrichetea* (lake influence), **ome** = oligo- to meso-euryhaline, **oms** = oligo- to meso-stenohaline, **os** = oligo-stenohaline, **p** = polysaprobic, **P** = planktonic, **pe** = poly-euryhaline, **PLE** = *Pleuronemetum coronatae*, **ps** = poly-stenohaline, **R** = predator (feeds on protozoans, mostly ciliates, some species even ingest small metazoans), **S** = stagnant waters, **Sb** = sulphur bacteria, **STE** = *Stentoretum*, **T** = epizoic, **TRI** = *Trithigmostometum cucullulae*, **x** = xenosaprobic.

- 1: Wet mass; $1 \mu\text{m}^3 = 1 \text{ pg}$, i.e. specific gravity of the protoplasm is 1.0 (FINLAY 1982)
- 2: For classification see Table 1. Data are often highly questionable and thus are then marked with a "?". Very few limnetic ciliates occur in truly marine environments although many species tolerate high salinities. Many freshwater species occur in saline estuaries together with some marine species, however, few marine ciliates occur in strongly saline inland waters.
- 3: See community plates. Many species cannot yet be classified into a certain community.
- 4: According to table 3 in FOISSNER et al. (1995)
- 5: Feeds also on epidermal cells, cnidocysts and food residues of *Hydra*.
- 6: Ingests also fish epidermal cells if the latter are very abundant.
- 7: For *Caenomorpha medusula*.
- 8: Also histophagous, i. e. feeding on cells of dying or dead metazoans.
- 9: Erroneously written "3.5 mg/l" in FOISSNER et al. (1994).
- 10: Not calculated because of complicated shape.
- 11: If very abundant, otherwise use a-b.
- 12: If very abundant, otherwise use b-a.
- 13: According to ALBRECHT (1984); erroneously classified as holo-euryhaline in FOISSNER et al. (1994).

Table 1: Salinity terminology (after ALBRECHT 1984). Cl = chloride (mg/l Cl^-), S = salinity (‰)

Cl	0 - 400	400 - 2000	2000 - 5000	5000 - 17 000	> 17 000
S	0 - 1	1 - 4	4 - 10	10 - 30	> 30
holo-euryhaline					
oligo-stenohaline		meso- to poly-euryhaline			
oligo- to meso-stenohaline			poly-euryhaline		
oligo-euryhaline				meso- to poly-stenohaline	
oligo- to meso-euryhaline					poly-stenohaline