

Review

An Updated Compilation of World Soil Ciliates (Protozoa, Ciliophora), with Ecological Notes, New Records, and Descriptions of new Species

Wilhelm Foissner

Universität Salzburg, Institut für Zoologie, Salzburg, Austria

Summary

This review compiles and analyses the taxonomic and biogeographical information available on soil ciliates. Key literature (257 references) is tabulated for each species. Furthermore, about 500 new records are reported from hundreds of samples collected world-wide, and four new species are described from Kenyan soils, including the new genus *Apybryophyllum*; *Metopus ovalis* Kahl, 1927 is redescribed. 643 ciliate species were originally described or reliably recorded from about 1000 soil samples world-wide, 49 (7.6%) of them were later recognized as junior synonyms, and 78 (13.2%) have been poorly described, leaving a total of 516 well-known species. However, the studied samples contained about 700 new species, most of which (about 500) have not yet been described. Thus, the total number of soil ciliates amounts to at least 1000 species. The overwhelming portion of soil ciliates belongs to three systematic assemblages, viz. the hypotrichs (39%), gymnostomatids (26%) and colpodids (13%) if the undescribed species are taken into account. Most soil ciliates feed on bacteria (39%) or are predatory (34%) or omnivorous (20%). Some are strictly mycophagous and highly characteristic for terrestrial habitats; and a few (mainly metopids) are anaerobic. Only about one fourth of the soil ciliate species has been reliably reported from freshwater habitats and from more than three out of five main biogeographical regions, indicating a high specificity of the soil ciliate biota and a limited distribution of at least some species. This is supported by the observation that some very conspicuous species, such as *Krassniggia auxiliaris* and *Bresslauides discoideus*, have so far been found only in Gondwanan, respectively, Laurasian soils.

Key words: Biodiversity; Community structure; Geographic distribution; New species; Soil ciliates; Taxonomy.

Introduction

The last comprehensive review on diversity and ecology of soil ciliates was published in 1987 [73]. It listed almost 300 species and showed that the soil ciliate

community is different from freshwater and other ciliate assemblages. Since then many new species and records of soil ciliates have been described. The present review summarises and analyses these and older taxonomic and biogeographical data. Furthermore, about 500 new records are listed from hundreds of samples collected world-wide. During these studies, more than 500 new species, a few of which are described here, were found, confirming a previous estimation that there are at least 1330–2000 species of soil ciliates [96].

Material and Methods

Areas and sampling

About 817 samples were collected and analysed from countries world-wide, including all main biogeographic regions (Tab. 1, Fig. 1). Generally, collections were made from a variety of biotopes covering most principal soil and vegetation types of the respective region. Detailed site descriptions for the new records will be provided in later publications.

The material collected usually included mineral top soil (0–5 cm depth) with fine plant roots, the humic layer, and the deciduous and/or grass litter from the soil surface. Furthermore, many samples contained some terrestrial or arboricolous mosses with adhering soil and/or bark from trees. All these habitats are referred to as “terrestrial”, as opposed to freshwater, because they contain, although in varying amounts, true humic and mineral soil [73]. Usually, about 10 small subsamples were taken from an area of about 100 m² and mixed to a composite sample. All samples were air-dried for at least one month and then sealed in plastic bags. Such samples can be stored for years without any significant loss of species [73, and unpubl. results].

Sample processing and community analysis

All collections were analysed with the non-flooded Petri dish method as described in [73, 83]. Briefly, this simple method involves placing 10–50 g terrestrial material in a Petri

Table 1. Origin and number of samples investigated.

Biogeographic region	Number of samples investigated
HOLARCTIS	
Austria	about 300 ^{1,3)}
Germany	about 10 ^{1,3)}
Denmark	12 ²⁾
Finland	10
Iceland	11 ²⁾
Portugal	2
Italy	2
Greece	10 ²⁾
Japan	23 ²⁾
	----- subtotal 380
PALAEOTROPIS	
Kenya	56 ²⁾
Tanzania	6
Cameroon	8
Namibia (Etosha Pan)	12
Republic of South Africa	10
	----- subtotal 92
AUSTRALIS	
Australia and Tasmania	158 ²⁾
NEOTROPIS	
Costa Rica	33 ²⁾
Venezuela	16
Brazil	5 ²⁾
Peru	4
Chile	29
	----- subtotal 87
ARCHINOTIS	
Continental and maritime Antarctica	about 73 ¹⁾
Gough & Marion Islands	27 ¹⁾
	----- subtotal 100
Total number of samples investigated	817

¹⁾ Taxonomic and faunistic data almost completely published, e.g. [2, 9, 12–14, 16, 62–65, 69, 70, 73–75, 77, 81, 94, 95, 108, 111, 113, 161, 162, 183, 184, 189].

²⁾ Taxonomic and faunistic data partially published, e.g. [8, 11, 20, 21, 49, 71, 72, 73, 75, 78, 87, 93, 97]. Note, however, that almost all colpodids from all samples investigated so far have been published [84–86, 89, 90, 93].

³⁾ Many sites were investigated several times during a period of 1–3 years; all samples from the other countries are from different sites. Details of these sites will be provided in later papers.

dish (10–15 cm in diameter) and saturating but not flooding it with distilled water. Such cultures were analysed for ciliates by inspecting about 2 ml of the run-off on days 2, 7, 14, 21, and 28. The non-flooded Petri dish method is selective, that is probably only a small proportion of the resting cysts present in a sample is reactivated, and undescribed species or species with specialized demands are very likely undersampled [96]. Thus, the real number of species, described and undescribed, in the samples investigated is probably much higher. Unfortunately, a better method for broad analysis of soil ciliates is not known.

Identification of species and species concept

Identification of species was according to the literature cited in this paper. Most of the species found were either new or described or redescribed by my students and myself. Thus, identification was mainly of live specimens using a high-power oil immersion objective and differential interference contrast. However, all “difficult”, new or supposedly new, species were treated with the silver staining techniques described in [82]. Usually, these methods yield permanent slides which have been or will be deposited in the Oberösterreichische Landesmuseum in Linz (LI).

The species concept, of course, influences the number of species found and/or recognised as “undescribed”. I usually apply the morphospecies concept which is, according to Finlay et al. [55], as valid as any, and probably more pragmatic than any other. I do not consider myself a splitter, that is I classify new species as such only when populations can be separated from their nearest relatives by at least one distinct (non-morphometric) morphological character. For examples, see papers cited and species described in this review.

Species descriptions

The species described were found in soils from Kenya, equatorial Africa. The soils of this region contain a very rich ciliate community. 507 species were found in 92 samples, 240 of them were undescribed [96].

Protospadidium terricola: Grassland soil (0–3 cm) from Mt. Kenya near the lodge “The Ark”, about 2300 m above sea level. Soil reddish, humic, pH 6.4. Collected on 1. 7. 1985, investigated on 25. 7. 1986.

Apobryophyllum terricola and *Tachysoma humicola longisetum*: Litter, humus, and mineral soil from a small plant island within young (some hundred years) lava masses of the Shetani volcano, a black and almost bare, fascinating area in the Tsavo National Park. Soil brownish, pH 7.6. Collected on 8. 7. 1985, investigated on 1. 4. 1986.

Metopus ovalis and *Keronopsis dieckmanni*: Grassland soil from shore of Lake Baringo, one of the highly saline, outstanding Flamingo lakes in Kenya. Soil brownish, humic and saline (salt crystals were formed when a drop of the soil water evaporated on the slide), pH 7.3. Collected on 2. 7. 1985, investigated in October 1985.

Methods follow those used in my previous papers [e.g. 82, 84, 95]. Two or three type slides (1 holotype and 1 to 2 paratypes) each of the new species described and two voucher slides of the species redescribed have been deposited in the Oberösterreichische Landesmuseum in Linz (LI), Austria. The slides usually contain many protargol-impregnated cells, with relevant specimens marked by a black ink circle on the cover glass.




Fig. 1. Biogeographic regions according to [174]. Hatched areas are transition zones.

Results and Discussion

The data are summarized in Tables 2–4, as in my previous review [73]. The present compilation is more complete, considers extensively the older literature, and includes many new records from soils world-wide (Tab. 2). Furthermore, the quality of species descriptions was analysed and detailed taxonomic literature compiled for each species (Tab. 3).

The new compilation largely confirms and strengthens the conclusions from my previous review [73], for instance, that soil ciliates are on average smaller and more slender than freshwater ciliates (Tab. 4). I believe that these are adaptations to the film-like distribution of the water and the narrowness of the habitat, that is the soil pores. Thus, the following sections concentrate on aspects which could not be successfully treated in 1987 [73] due to the limited data available, for instance, concerning geographic distribution.

A special note is necessary on my unpublished taxonomic material which is occasionally referred to. It amounts to about 500 new species, including many new genera and some new families. These species, the full description of which will require years of work, were found in the samples listed in Table 1 and in about 150 other samples not included in this review because the data have not been sufficiently checked.

Uniqueness of the soil ciliate biota

Only 25% (16% if the undescribed species are included) of the described species have been reliably recorded from both soil and freshwater habitats (Tab. 4). Thus, there would appear to be a high proportion of species which occur only or mainly in terrestrial environments and very likely evolved in such habitats [73]. The most characteristic soil ciliates are colpodids, but many autochthonous genera and species occur also among hypotrichs (e.g., *Circinella* spp., *Hemisincirra* spp., *Erniella filiformis*, *Terricirra* spp.) and gymnostomatids (e.g., *Coriplites terricola*, *Pleuroplitoides smithi*, *Spathidium* spp., *Epispathidium* spp.).

Community structure

Most soil ciliates belong to three systematic assemblages (Tab. 4), that is hypotrichs (36%), colpodids (22%) and gymnostomatids (17%). However, these proportions change considerably if the as yet undescribed species mentioned above are included (Tab. 4): 39% hypotrichs, 26% gymnostomatids, 13% colpodids. The dominance of colpodids in the published material is caused by the fact that nearly all new species found have been described [84–86, 89, 90, 93; Tab. 4]. However, even if the lower, more realistic figure (13%) is taken, the diversity of colpodids is much higher in soil than in freshwater (13% vs. 5% [73]). The reason for the high diversi-

ty of colpodids in terrestrial environments is not well understood. Possibly, their ability to evolve diverse feeding strategies (bacterivory, fungivory, omnivory, predation) and body sizes (10–1000 µm) favoured colonization of and adaptive radiation in such habitats, as indicated by the high proportion of autochthonous species [84, 92]. The overwhelming diversity of hypotrichs is more easily explained. All are ciliated only on the ventral (oral) surface and most are very slender and/or strongly flattened. Such organisation is pre-adaptive in terrestrial habitats, where water is frequently available only as a thin film covering soil particles and pores. The same structural organisation is found in cyrtophorid ciliates, a conspicuous and diverse group in the Aufwuchs (periphyton) of freshwaters [114]. Their low diversity (2.2%) in soils is possibly related to food: most cyrtophorids prefer diatoms which are, compared to freshwaters, rare in soil. The considerable number of gymnostomatids, most of which are rapacious carnivores, indicates that the soil ciliate community is well-structured.

The compilation contains also some planktonic (e.g., *Linostoma vorticella*, *Hastatella radians*) and anaerobic (*Metopus* spp., *Brachonella* spp.) species (Tab. 2). These were mainly found in soils from swamps and/or floodplains. Most were recorded in few samples only, except for some *Metopus* species, which are rather common and excellent indicators of anaerobic conditions [115].

The number of ciliate species living in soil

Table 2 lists 643 species found in 817 samples analysed by my group (Tab. 1) and in an unknown number of collections investigated by others, especially Greeff [135], Penard [177–179], Kahl [152–157], Gellér [126–130], Buitkamp [26–28], and Hemberger [138]. Supposing that these authors probably studied about 200 samples, the compilation would be based on roughly 1000 collections.

Of the 643 ciliate species originally described or reliably recorded from terrestrial habitats, 49 (7.6%) were later recognized as junior, subjective synonyms, and 78 (13.2%) have been poorly described (Tab. 2–4), leaving a total of 516 well-known species. If we add 500 new, not yet described species mentioned above, we obtain roughly 1000 species of soil ciliates, that is about half the number recently estimated to be present by a probability calculation [96]. The synonymy rate is low (7.6%) as compared, for instance, to insects, where rates between 20–50% have been reported [121, 122, 168], and to ciliates in general, for which Finlay et al. [55] estimated a rate of 20–30%. The low synonymy rate in soil ciliates is very likely caused by the limited number of specialists working in this field, the distinctiveness of the biota, and the fortunate situation that more than half of the species were rather recently and

thoroughly described using appropriate techniques like silver impregnation.

Further progress and a distinct rise in species number may be expected when specialists will begin to investigate the epizoic ciliate biota of euterrestrial invertebrates, an almost unexplored field, with a few exceptions, viz. *Pyxidium longicollum* attached to shells of testate amoebae [19] and *P. tardigradum* living on tardigrades [233]. Likewise, ciliates attached to soil particles and moss leaves via a stalk are poorly known because they are usually not found with the non-flooded Petri dish method. Few of the sessile, muscicolous peritrichs and suctoria described by Penard [177, 178] have been rediscovered.

Another rise in the number of species can be expected when the large floodplains of the world are explored for soil ciliates. Twenty-one new species were found in two samples from the Amazon floodplain near Manaus [97].

Geographic distribution

The high ciliate diversity in terrestrial environments remained unrecognised for a long time because of the lack of an appropriate isolation method (still a great problem [96]) and the view of many ecologists that soil ciliates are common invaders from freshwaters and activated sludge (for reviews, see [73, 91, 92]). Thus, most of the older faunistic and ecological literature is burdened with misidentifications and is almost useless. Accordingly, it has been excluded from this review.

The reliable information on geographic distribution of soil ciliates is compiled in Table 2, and includes about 500 new records mainly from tropical soils in Africa, Australia, Costa Rica, and South America. A rough framework of only five widely recognised biogeographic regions (Holarctic, Palaeotropis, Australis, Neotropis, Archinotis; Fig. 1) has been used because of the very limited data available. Substantial biogeographic information has been extracted mainly from the following studies, most of which concern the Holarctic region: [2, 16, 17, 20, 26, 27, 30, 35, 62, 71–74, 78, 84–86, 93–95, 97, 108, 113, 128, 129, 132, 143, 152, 154, 155, 157, 161, 162, 182, 185, 189, 190, 195, 198, 199, 205, 206, 229, 230, 240].

While ecologists tend to consider protozoa as cosmopolitan [55], taxonomists have argued for a restricted distribution of at least some genera and species [73, 91, 92, 96]. In my experience, many morphological species of soil ciliates are cosmopolitan, depending mainly on appropriate habitats. This is also indicated by the rather limited genetic differentiation found among 15 isolates from three continents of a common soil ciliate, *Colpoda inflata* [24].

The present review cannot provide a definite conclusion on the extent of endemism in soil ciliates because

Table 2. Taxonomic and ecological summary of world soil ciliate fauna¹⁾. For statistics (number of species, percentage of synonyms, etc.), see Table 4.

Species ²⁾	Taxonomic group ³⁾	Biomass of 10 ⁶ indiv. (mg) ⁴⁾	Food ⁵⁾	Geographic distribution ⁶⁾					Degree of autochthonism ⁷⁾
				H	P	A	N	Ar	
<i>Acaryophrya collaris</i> (Kahl, 1926) Dingfelder, 1962	GY	113	B,C	+	+	-	-	-	*
<i>Acineria uncinata</i> Tucolesco, 1962	GY	10	C,F	+	+	-	+	+	*
<i>Acineta flava</i> Kellicott, 1885	SU	30	C?	+	-	-	-	-	*
<i>Acropisthium mutabile</i> Perty, 1852	GY	35	C	+	-	+	+	-	*
<i>Actinobolina vorax</i> (Wenrich, 1929) Kahl, 1930	GY	250	C,R	-	+	-	-	-	*
<i>Alinostoma multivacuolatum</i> Alekperov, 1993	CY	700	?	+	-	-	-	-	**
<i>Amphibotrella enigmatica</i> Grandori & Grandori, 1934	?	1750	?	+	-	-	-	-	**
<i>Amphisiella binucleata</i> (Hemberger, 1985) Foissner, 1988	HY	96	F?	-	+	-	+	-	**
<i>Amphisiella magnigranulosa</i> Foissner, 1988	HY	90	C,G,H	+	+	+	+	-	**
<i>Amphisiella polycirrata</i> Berger & Foissner, 1989	HY	120	C,F	-	+	-	+	-	**
<i>Amphisiella quadrinucleata</i> Berger & Foissner, 1989	HY	70	C	+	-	-	-	-	**
<i>Amphisiella raptans</i> Buitkamp & Wilbert, 1974	HY	1280	C,N	+	-	-	-	-	***
<i>Amphisiella terricola</i> Gellért, 1955	HY	100	C	+	+	+	+	-	***
<i>Amphisiella vitiphila</i> (Foissner, 1987) Foissner, 1988	HY	92	C,F	-	+	-	+	-	**
<i>Amphisiellides atypicus</i> (Hemberger, 1985) Foissner, 1988	HY	184	?	-	-	-	+	-	**
<i>Amphisiellides illuvialis</i> Eigner & Foissner, 1994	HY	38	B,C	+	-	-	-	-	*
<i>Apoamphisiella tihanyiensis</i> (Gellért & Tamás, 1958) Foissner, 1997	HY	300	C,E,F,H,N, R,T	-	-	-	+	-	*
<i>Apobryophyllum terricola</i> Foissner (this paper)	GY	30	C?	-	+	-	-	-	***
<i>Apocolpoda africana</i> Foissner, 1993	CO	16	B	-	+	-	-	-	***
<i>Archinassula muscicola</i> Kahl, 1935	NA?	77	B	+	-	-	-	-	**
<i>Arcuospathidium atypicum</i> (Wenzel, 1953) nov. comb.	GY	20	C,H	+	+	+	+	-	**
<i>Arcuospathidium australe</i> Foissner, 1988, synonym with <i>A. atypicum</i>	GY	38	C?	-	-	-	-	+	**
<i>Arcuospathidium cooperi</i> Foissner, 1996	GY	180	C	+	+	+	+	-	***
<i>Arcuospathidium cultriforme</i> (Penard, 1922) Foissner, 1984	GY	23	C	+	-	-	+	-	**
<i>Arcuospathidium japonicum</i> Foissner, 1988	GY	400	C	+	+	+	+	-	***
<i>Arcuospathidium lionotiforme</i> (Kahl, 1930) Foissner, 1984	GY	87	C,H	+	+	+	+	-	**
<i>Arcuospathidium muscorum</i> (Dragesco & Dragesco-Kernéis, 1979) Foissner, 1984	GY	28	C?	+	+	-	-	-	***
<i>Aspidisca cicada</i> (Müller, 1786) Claparède & Lachmann, 1858	HY	5	B	+	-	-	-	-	*
<i>Aspidisca lynceus</i> (Müller, 1773) Ehrenberg, 1830-	HY	6	B	+	-	+	-	-	*
<i>Australocirrus octonucleatus</i> Foissner, 1988	HY	720	B,C,F,G,H	+	+	+	+	-	**
<i>Australocirrus oscitans</i> Blatterer & Foissner, 1988	HY	990	C,F,G,H,T	+	+	+	+	-	***
<i>Australothrix alwinae</i> Blatterer & Foissner, 1988	HY	1200	B,C,F,H,R,T	-	-	+	-	-	**
<i>Australothrix australis</i> Blatterer & Foissner, 1988	HY	1120	C,F,H,R,S,T	-	-	+	-	-	**
<i>Australothrix simplex</i> Shen, Liu, Song & Gu, 1992	HY	270	?	+	-	-	-	-	***
<i>Australothrix steineri</i> Foissner, 1995	HY	81	B,F,H	-	-	+	+	-	**
<i>Avestina acuta</i> (Buitkamp, 1977) Jankowski, 1980	CO	4	B	+	+	-	-	-	**
<i>Avestina ludwigi</i> Aesch & Foissner, 1990	CO	5	H	+	-	+	+	-	***
<i>Bakuella edaphoni</i> Song, Wilbert & Berger, 1992	HY	600	B,C,D,F,T	+	-	-	-	-	**
<i>Bakuella pampinaria</i> Eigner & Foissner, 1992	HY	80	C,F,H	+	-	-	-	-	**
<i>Balantidiooides bivacuolata</i> Kahl, 1932	HE	280	C	+	-	-	-	-	***
<i>Balantidiooides corbifera</i> (Fryd-Versavel & Tuffrau, 1978) Foissner, Adam & Foissner, 1982	HE	360	C,G	+	-	-	-	-	**
<i>Balantidiooides dragescoi</i> Foissner, Adam & Foissner, 1982	HE	135	C	+	+	+	-	-	***
<i>Balantidiooides muscicola</i> (Penard, 1922) Penard, 1930 in Kahl (1930)	HE	120	R	+	-	-	-	-	***
<i>Bardeliella pulchra</i> Foissner, 1984	CO	8	B	+	+	-	-	-	***
<i>Bicoronella costaricana</i> Foissner, 1995	HY	260	C,E,H,T	-	-	-	+	-	**
<i>Birojimia muscorum</i> (Kahl, 1932) Berger & Foissner, 1989	HY	100	C,F,H,S	+	+	+	+	+	**
<i>Birojimia terricola</i> Berger & Foissner, 1989	HY	140	C,F,H	+	-	-	-	-	**
<i>Blepharisma americanum</i> Suzuki, 1954	HE	640	B,C,F	-	-	-	+	-	*
<i>Blepharisma biancae</i> Lepsi, 1948	HE	10	?	+	-	-	-	-	**

Table 2. continued.

Species ²⁾	Taxonomic group ³⁾	Biomass of 10 ⁶ indiv. (mg) ⁴⁾	Food ⁵⁾	Geographic distribution ⁶⁾					Degree of autochthonism ⁷⁾
				H	P	A	N	Ar	
<i>Blepharisma bimicronucleatum</i> Villeneuve-Brachon, 1940	HE	42	B	+	+	+	+	+	*
<i>Blepharisma hyalinum</i> Perty, 1849	HE	11	B,H	+	+	+	+	+	*
<i>Blepharisma ichthyoides</i> Gelei, 1939, synonym with <i>B. lateritium</i>									
<i>Blepharisma lateritium</i> (Ehrenberg, 1831) Kahl, 1932	HE	525	B,G	+	-	-	-	-	*
<i>Blepharisma ovatum</i> (Stokes, 1884) Penard, 1922	HE	750	C,D,G	-	+	-	-	-	*
<i>Blepharisma steini</i> Kahl, 1932	HE	69	B	+	+	+	+	+	*
<i>Blepharisma undulans</i> Stein, 1867	HE	225	B	+	+	+	-	-	*
<i>Brachonella cydonia</i> (Kahl, 1927) Jankowski, 1964	HE	113	B	-	-	-	+	-	*
<i>Bresslaua insidiatrix</i> Claff, Dewey & Kidder, 1941	CO	192	C,F	+	+	-	-	-	**
<i>Bresslaua sicaria</i> Claff, Dewey & Kidder, 1941	CO	200	C,H	+	-	-	+	-	**
<i>Bresslaua vorax</i> Kahl, 1931	CO	113	B,C,F,G,H	+	+	+	+	-	**
<i>Bresslauides australis</i> Blatterer & Foissner, 1988	CO	320	B,C,F,G,H,N	-	-	+	-	-	***
<i>Bresslauides discoideus</i> (Kahl, 1931) Foissner, 1993	CO	9000	C,E,R	+	-	-	-	-	***
<i>Bresslauides terricola</i> (Foissner, 1987) Foissner, 1993	CO	2200	C,H	+	+	+	+	-	***
<i>Bryometopus alekperovi</i> nov. comb., nov. nom. ¹⁰⁾	CO	3200	?	+	-	-	-	-	**
<i>Bryometopus atypicus</i> Foissner, 1980	CO	45	B,F,G	+	+	+	+	-	*
<i>Bryometopus balantidiooides</i> Foissner, 1993	CO	30	B	-	-	+	-	-	**
<i>Bryometopus edaphonus</i> Foissner, 1980	CO	24	B?	+	-	-	-	-	**
<i>Bryometopus hawaiiensis</i> Foissner, 1994	CO	30	B,F,G,S	+	-	+	-	-	*
<i>Bryometopus pseudochilodon</i> Kahl, 1932	CO	22	B,G	+	+	+	+	+	**
<i>Bryometopus sphagni</i> (Penard, 1922) Kahl, 1932	CO	198	B,C,D,G	+	+	+	-	+	*
<i>Bryometopus triquetrus</i> Foissner, 1993	CO	10	B,E,F	+	+	+	+	-	**
<i>Bryophrya bavaricensis</i> (Kahl, 1931) Wenzel, 1953	CO	85	E	+	-	-	-	-	***
<i>Bryophrya rubescens</i> (Penard, 1922) Foissner, 1993	CO	150	E	+	-	-	-	-	**
<i>Bryophyllum loxophylliforme</i> Kahl, 1931	GY	252	C,E,H,R	+	+	+	+	+	**
<i>Bryophyllum tegularum</i> Kahl, 1931	GY	100	C,R	+	-	+	-	-	**
<i>Buitkampia angusta</i> Foissner, 1985, synonym with <i>Platyophrya vorax</i>									
<i>Bursaria truncatella</i> Müller, 1773	CO	50000	C,D,F,G,R,T	+	+	+	+	-	*
<i>Chaenea clavata</i> Grandori & Grandori, 1934	GY	50	?	+	-	-	-	-	**
<i>Chaenea humicola</i> Gellért, 1957	GY	4	?	+	-	-	-	-	**
<i>Chilodon geographicus</i> Penard, 1922, synonym with <i>Odontochlamys gourandi</i>									
<i>Chilodonella aplanata</i> Kahl, 1931	CY	10	?	+	-	-	-	-	*
<i>Chilodonella nigra</i> Lepsi, 1951	CY	?	?	+	-	-	-	-	**
<i>Chilodonella uncinata</i> (Ehrenberg, 1838) Strand, 1928	CY	13	B	+	+	+	+	-	*
<i>Chilodonella uncinata</i> (Ehrenberg) in Gellért [126], synonym with <i>Pseudochilodonopsis mutabilis</i>									
<i>Chilodontopsis muscorum</i> Kahl, 1931	NA	15	?	+	+	-	-	-	***
<i>Chilophrya terricola</i> Foissner, 1984	PR	9	C?	+	-	-	-	-	**
<i>Chlamydonella alpestris</i> Foissner, 1979	CY	4	B,D	+	-	-	-	-	*
<i>Cinetochilum margaritaceum</i> (Ehrenberg, 1830) Perty, 1852	HM	10	B,D,G	+	+	+	+	-	*
<i>Cinetochilum marinum</i> Pomp & Wilbert, 1988	HM	5	B	-	-	+	-	-	**
<i>Circinella arenicola</i> Foissner, 1994	HY	120	B	+	-	-	-	-	***
<i>Circinella filiformis</i> (Foissner, 1982) Foissner, 1994	HY	5	?	+	+	+	-	-	***
<i>Circinella vettorsi</i> (Berger & Foissner, 1989) Foissner, 1994	HY	23	?	+	+	-	+	-	***
<i>Cirrophrya australis</i> Foissner, 1993	CO	20	?	-	-	+	-	-	***
<i>Cirrophrya haptica</i> Gellért, 1950	CO	25	B,G	+	-	-	-	-	***
<i>Cirrophrya terricola</i> Foissner, 1987	CO	20	C	+	-	-	-	-	***
<i>Cladotricha australis</i> Blatterer & Foissner, 1988	HY	20	B,H	-	-	+	-	-	**
<i>Colpoda aspera</i> Kahl, 1926	CO	6	B	+	+	+	+	+	*
<i>Colpoda atra</i> Alekperov, 1993	CO	1600	B,F	+	-	-	-	-	**
<i>Colpoda augustini</i> Foissner, 1987	CO	20	B	+	+	+	+	-	***
<i>Colpoda bifurcata</i> Alekperov, 1993, synonym with <i>C. lucida</i>									
<i>Colpoda californica</i> Kahl, 1931	CO	5	?	+	-	-	+	-	**
<i>Colpoda cavicola</i> Kahl, 1935	CO	240	B,G,H	+	-	-	+	-	**
<i>Colpoda colpodiopsis</i> Kahl, 1931	CO	12	B,F	+	+	-	-	-	*
<i>Colpoda cucullus</i> (Müller, 1773) Gmelin, 1790	CO	74	B,F,G	+	+	+	+	+	*

Table 2. continued.

Species ²⁾	Taxonomic group ³⁾	Biomass of 10 ⁶ indiv. (mg) ⁴⁾	Food ⁵⁾	Geographic distribution ⁶⁾					Degree of autochthonism ⁷⁾	
				H	P	A	N	Ar		
<i>Colpoda duodenaria</i> Taylor & Furgason, 1938, synonym with <i>C. steinii</i>										
<i>Colpoda ecaudata</i> (Liebmann, 1936) Foissner, Blatterer, Berger & Kohmann, 1991	CO	9	B	+	+	+	+	+	*	
<i>Colpoda edaphoni</i> Foissner, 1980	CO	10	B	+	+	+	+	-	***	
<i>Colpoda elliotti</i> Bradbury & Outka, 1967	CO	5	B	+	+	+	+	-	**	
<i>Colpoda eurystoma</i> Gellért, 1950, synonym with <i>Kalometopia duplicita</i>										
<i>Colpoda fastigata</i> Kahl, 1931, synonym with <i>C. maupasi</i>										
<i>Colpoda flavicans</i> (Stokes, 1885) Foissner, 1993	CO	80	B,E	+	+	+	+	-	*	
<i>Colpoda henneguyi</i> Fabre-Domergue, 1889	CO	122	B,H	+	+	+	+	+	**	
<i>Colpoda inflata</i> (Stokes, 1884) Kahl, 1931	CO	40	B,F	+	+	+	+	+	*	
<i>Colpoda irregularis</i> Kahl, 1931	CO	72	?	+	+	-	-	-	*	
<i>Colpoda lucida</i> Greeff, 1888	CO	230	B	+	+	+	+	+	**	
<i>Colpoda magna</i> (Gruber, 1879) Lynn, 1978	CO	1400	B,C,F,G	+	+	+	+	-	*	
<i>Colpoda maupasi</i> Enriques, 1908	CO	39	B	+	+	+	+	+	**	
<i>Colpoda minima</i> (Alekperov, 1985) Foissner, 1993	CO	420	B	+	+	+	+	-	*	
<i>Colpoda orientalis</i> Foissner, 1993	CO	300	B	+	+	-	+	-	**	
<i>Colpoda praestans</i> Penard, 1922	CO	750	B,E,H	+	-	-	+	-	**	
<i>Colpoda reniformis</i> Kahl, 1931	CO	180	E?	+	-	-	-	-	**	
<i>Colpoda simulans</i> Kahl, 1931	CO	70	?	+	-	-	-	-	**	
<i>Colpoda spiralis</i> Novotny, Lynn & Evans, 1977, synonym with <i>C. cavicola</i>										
<i>Colpoda steinii</i> Maupas, 1883	CO	5	B,E,G,H	+	+	+	+	+	*	
<i>Colpoda tripartita</i> Kahl, 1931	CO	240	B	+	+	+	+	-	***	
<i>Colpoda variabilis</i> Foissner, 1980	CO	300	B,G	+	+	-	+	+	*	
<i>Colpodidium caudatum</i> Wilbert, 1982	NA	30	B	+	+	+	+	-	***	
<i>Colpodidium viridis</i> (Mirabdullaev, 1986) Jankowski, 1992	NA	80	?	-	+	-	+	-	*	
<i>Condylostoma terricola</i> Foissner, 1995	HE	70	C,F,H,S	-	-	-	+	-	**	
<i>Corallocolpoda grelli</i> (Foissner, 1993) Foissner, 1993	CO	20	C,H	-	-	+	-	-	***	
<i>Corallocolpoda pacifica</i> Alekperov, 1991	CO	280	C	-	+	+	+	-	***	
<i>Coriplites terricola</i> Foissner, 1988	GY	7	C,F	+	+	+	+	-	**	
<i>Corticocolpoda kaneshiroae</i> Foissner, 1993	CO	1050	C	-	-	+	-	-	***	
<i>Cosmocolpoda naschbergeri</i> Foissner, 1993	CO	75	B	-	-	-	+	-	***	
<i>Cothurnia minutissima</i> (Penard, 1914) Kahl, 1935	PE	15	B	+	-	-	-	-	**	
<i>Cothurnia regalis</i> Penard, 1914, synonym with <i>Thuricola innixa</i>										
<i>Cothurnia richtersi</i> (Penard, 1914) Kahl, 1935	PE	12	B	+	-	-	-	+	**	
<i>Cothurniopsis elastica</i> Penard, 1914, synonym with <i>C. valvata</i>										
<i>Cothurniopsis valvata</i> Stokes, 1893	PE	8	B	+	-	-	-	-	*	
<i>Cyclidium glaucoma</i> Müller, 1773	HM	2	B,G	+	+	-	+	+	*	
<i>Cyclidium muscicola</i> Kahl, 1931	HM	1	B	+	+	+	+	+	**	
<i>Cyclidium opisthostoma</i> Grandori & Grandori, 1934, synonym with <i>Homalogastra setosa</i>										
<i>Cyclidium terricola</i> Kahl, 1931	HM	3	B	+	+	+	-	-	**	
<i>Cyrtobymena australis</i> Foissner, 1995	HY	1330	C,H,T	-	-	+	+	-	**	
<i>Cyrtobymena balladynula</i> (Kahl, 1932) Foissner, 1989	HY	4	?	+	-	-	-	-	**	
<i>Cyrtobymena candens</i> (Kahl, 1932) Foissner, 1989	HY	150	B,C,F,N	+	+	+	+	+	**	
<i>Cyrtobymena candens depressa</i> (Gellért, 1942) Foissner, 1989	HY	120	B,D,F	+	+	+	+	-	**	
<i>Cyrtobymena citrina</i> (Berger & Foissner, 1987) Foissner, 1989	HY	54	C,D,H,T	+	+	+	+	-	**	
<i>Cyrtobymena dubia</i> (Gellért, 1956) nov. comb., synonym with <i>C. muscorum</i>										
<i>Cyrtobymena gracilis</i> (Kahl, 1932) Foissner, 1989	HY	60	?	+	-	-	-	-	**	
<i>Cyrtobymena granulata</i> (Kahl, 1932) Foissner, 1989	HY	40	?	+	-	-	-	-	**	
<i>Cyrtobymena muscorum</i> (Kahl, 1932) Foissner, 1989	HY	225	C,D,F,H,T	+	+	-	-	-	***	
<i>Cyrtobymena primicirrata</i> (Berger & Foissner, 1987) nov. comb.	HY	77	C,D,F,H	+	+	+	+	-	**	
<i>Cyrtobymena quadrinucleata</i> (Dragesco & Njiné, 1971) Foissner, 1989	HY	81	C	+	+	+	+	+	*	
<i>Cyrtobymena tetricirrata</i> (Gellért, 1942) Foissner, 1989	HY	150	B,C,F	+	+	+	-	-	***	
<i>Cyrtolophosis acuta</i> Kahl, 1926	CO	1	B	+	+	+	-	+	*	

Table 2. continued.

Species ²⁾	Taxonomic group ³⁾	Biomass of 10 ⁶ indiv. (mg) ⁴⁾	Food ⁵⁾	Geographic distribution ⁶⁾					Degree of autochthonism ⁷⁾
				H	P	A	N	Ar	
<i>Cyrtolophosis colpidiformis</i> Foissner, 1993	CO	7	B	+	-	-	-	-	**
<i>Cyrtolophosis elongata</i> (Schewiakoff, 1892) Kahl, 1931	CO	1	B	+	+	+	+	+	*
<i>Cyrtolophosis minor</i> Vuxanovici, 1963	CO	1	B	-	-	+	+	-	*
<i>Cyrtolophosis mucicola</i> Stokes, 1885	CO	2	B	+	+	+	+	+	*
<i>Dapedophrya flexilis</i> (Penard, 1922) Foissner, 1995	CO	36	B,H	+	-	-	+	-	***
<i>Deviata bacilliformis</i> (Gelei, 1954) Eigner, 1995	HY	47	B,G	+	+	+	-	-	*
<i>Didinium nasutum</i> (Müller, 1773) Stein, 1859	GY	500	C,G	+	+	-	-	-	*
<i>Dileptus alpinus</i> Kahl, 1931 ¹¹⁾	GY	23	C	+	+	+	+	+	**
<i>Dileptus americanus</i> Kahl, 1931 ¹¹⁾	GY	63	C,F	+	+	+	-	-	**
<i>Dileptus anguillula</i> Kahl, 1931 ¹¹⁾	GY	15	C	+	+	+	+	-	**
<i>Dileptus binucleatus</i> Kahl, 1931 ¹¹⁾	GY	158	C	+	-	-	-	-	**
<i>Dileptus breviproboscis</i> Foissner, 1981, synonym with <i>D. anguillula</i>									
<i>Dileptus conspicuus</i> Kahl, 1931 ¹¹⁾	GY	168	C,D,F	+	-	+	-	-	***
<i>Dileptus conspicuus telobivacuolatus</i> Gellért, 1955 ¹¹⁾	GY	81	C	+	-	-	-	-	***
<i>Dileptus costaricanus</i> Foissner, 1995 ¹¹⁾	GY	198	C	-	-	-	+	-	**
<i>Dileptus edaphoni</i> Song, 1994 ¹¹⁾	GY	82	C	+	-	-	-	-	**
<i>Dileptus falciformis</i> Kahl, 1931 ¹¹⁾	GY	247	C	+	-	-	-	-	**
<i>Dileptus gracilis</i> Kahl, 1931 ¹¹⁾	GY	43	C	+	+	+	-	-	**
<i>Dileptus kahli</i> Srámek-Hušek, 1957 ⁸⁾ ¹¹⁾	GY	180	C	+	-	-	-	-	*
<i>Dileptus margarifer</i> (Ehrenberg, 1838) Wirnsberger, Foissner & Adam, 1984 ¹¹⁾	GY	500	C,E,G,N,R	+	+	-	-	-	*
<i>Dileptus mucronatus</i> Penard, 1922 ¹¹⁾	GY	456	C	+	+	+	-	-	*
<i>Dileptus orientalis</i> Song, Packroff & Wilbert, 1988 ¹¹⁾	GY	31	B	+	-	-	-	-	**
<i>Dileptus polyvacuolatus</i> Foissner, 1989 ¹¹⁾	GY	212	C	+	+	-	+	-	*
<i>Dileptus similis</i> Foissner, 1995 ¹¹⁾	GY	314	C	-	-	-	+	-	**
<i>Dileptus tenuis</i> Penard, 1922 ¹¹⁾	GY	7	C	+	-	-	-	-	**
<i>Dileptus terrenus</i> Foissner, 1981 ¹¹⁾	GY	226	C	+	+	-	+	-	**
<i>Dileptus visscheri</i> Dragesco, 1963 ¹¹⁾	GY	64	C	+	-	+	-	-	*
<i>Dimacrocyaron amphileptoides</i> (Kahl, 1931) Jankowski, 1967	GY	127	G,H,T	+	+	+	+	+	**
<i>Drepanomonas borzai</i> Lepsi, 1948, synonym with <i>D. revoluta</i>									
<i>Drepanomonas dentata</i> Fresenius, 1858	NA	5	B	-	-	-	+	-	*
<i>Drepanomonas exigua</i> Penard, 1922	NA	1	B	+	-	-	+	-	***
<i>Drepanomonas muscicola</i> Foissner, 1987	NA	1	B	+	+	+	+	+	**
<i>Drepanomonas pauciliata</i> Foissner, 1987	NA	1	B	+	+	+	+	+	**
<i>Drepanomonas revoluta</i> Penard, 1922	NA	1	B,F	+	+	+	+	+	*
<i>Drepanomonas sphagni</i> Kahl, 1931	NA	1	B	+	+	+	+	+	**
<i>Enchelydium piliforme</i> (Kahl, 1930) Foissner, 1984	GY	100	C	+	-	-	-	-	*
<i>Enchelydium polynucleatum</i> Foissner, 1984	GY	255	C	+	+	+	+	+	**
<i>Enchelydium terrenum</i> Foissner, 1984	GY	57	C	+	-	+	+	-	**
<i>Enchelyodon californicus</i> Kahl, 1935	GY	110	C?	+	-	-	-	-	**
<i>Enchelyodon lagenula</i> (Kahl, 1930) Blatterer & Foissner, 1988	GY	10	C?	+	+	+	+	-	**
<i>Enchelyodon longinucleatus</i> Foissner, 1984	GY	63	C?	+	+	+	+	-	**
<i>Enchelyodon nodosus</i> Berger, Foissner & Adam, 1984	GY	367	C?	+	+	-	-	-	**
<i>Enchelyodon terrenus</i> Foissner, 1984	GY	132	C?	+	+	-	-	-	**
<i>Enchelyodon tratzi</i> Foissner, 1987	GY	23	C?	+	+	-	+	-	**
<i>Enchelyomorpha vermicularis</i> (Smith, 1899) Kahl, 1930	SU	2	C?	+	-	-	-	-	*
<i>Enchelyotricha binucleata</i> Foissner, 1987	GY	16	C	+	-	-	-	-	**
<i>Enchelys agricola</i> Horváth, 1956, synonym with <i>Platyophrya spumacola</i>									
<i>Enchelys multmicronucleata</i> Alekperov, 1993	GY	3	?	+	-	-	-	-	**
<i>Enchelys multinucleata</i> (Dragesco & Dragesco-Kernéis, 1979) Berger, Foissner & Adam, 1984	GY	141	C	+	+	+	+	-	**
<i>Enchelys terricola</i> Foissner, 1987	GY	99	C	+	-	-	-	-	**
<i>Enchelys tokkuri</i> Shibuya, 1930	GY	157	?	+	-	-	-	-	**
<i>Enchelys vermiformis</i> Foissner, 1987	GY	50	C?	+	-	-	-	-	***
<i>Engelmanniella mobilis</i> (Engelmann, 1862) Foissner, 1982	HY	34	B	+	+	+	-	-	*

Table 2. continued.

Species ²⁾	Taxonomic group ³⁾	Biomass of 10 ⁶ indiv. (mg) ⁴⁾	Food ⁵⁾	Geographic distribution ⁶⁾					Degree of autochthonism ⁷⁾
				H	P	A	N	Ar	
<i>Epispathidium amphoriforme</i> (Greeff, 1888) Foissner, 1984	GY	180	C	+	+	+	+	-	**
<i>Epispathidium ascendens</i> (Wenzel, 1955) Foissner, 1987	GY	64	C	+	+	+	+	+	**
<i>Epispathidium papilliferum</i> (Kahl, 1930) Foissner, 1984	GY	135	C	+	-	+	+	-	**
<i>Epispathidium regium</i> Foissner, 1984	GY	360	C	+	+	-	-	-	**
<i>Epispathidium terricola</i> Foissner, 1987	GY	101	C	+	+	+	+	+	**
<i>Epistylis alpestris</i> Foissner, 1978	PE	79	B	+	+	+	+	-	*
<i>Erniella filiformis</i> Foissner, 1987	HY	100	B?,C	-	+	-	-	-	***
<i>Eschaneustyla brachytoma</i> Stokes, 1886	HY	146	C,D,G,H	+	+	-	+	-	**
<i>Eschaneustyla terricola</i> Foissner, 1982, synonym with <i>E. brachytoma</i>									
<i>Euploites corsica</i> Berger & Foissner, 1989 ¹¹⁾	HY	8	B?	+	-	-	-	-	**
<i>Euploites finki</i> Foissner, 1982 ¹¹⁾	HY	17	B	+	-	-	+	-	**
<i>Euploites labiatus</i> Ruinen, 1938 ¹¹⁾	HY	15	B?	+	-	+	-	-	*
<i>Euploites muscicola</i> Kahl, 1932 ¹¹⁾	HY	49	B,E	+	+	+	+	-	**
<i>Euploites muscorum</i> Dragesco, 1970, synonym with <i>E. muscicola</i>									
<i>Euploites terricola</i> Penard, 1922 ¹¹⁾	HY	36	B?	+	-	-	-	-	**
<i>Frontonia depressa</i> (Stokes, 1886) Kahl, 1931	HM	95	C,G,H,N	+	+	+	+	+	*
<i>Frontonia parameciiformis</i> Wenzel, 1953	HM	20	?	+	-	-	-	-	**
<i>Frontonia parvula</i> Penard, 1922, synonym with <i>F. depressa</i>									
<i>Frontonia solea</i> Foissner, 1987	HM	120	E	-	+	-	-	-	*
<i>Frontonia terricola</i> Foissner, 1987 ⁸⁾	HM	192	C	+	+	-	-	-	**
<i>Furgasonia trichocystis</i> (Stokes, 1894) Jankowski, 1964	NA	72	B	-	+	-	-	-	*
<i>Fuscheria lacustris</i> Song & Wilbert, 1989	GY	23	C?	+	-	+	-	+	*
<i>Fuscheria nodosa</i> Foissner, 1983	GY	42	C	-	+	+	-	-	*
<i>Fuscheria terricola</i> Berger, Foissner & Adam, 1983	GY	20	C	+	+	-	+	+	***
<i>Gastronauta derouxi</i> Blatterer & Foissner, 1992	CY	48	B?,H?	+	+	-	+	+	***
<i>Gastronauta membranaceus</i> Bütschli, 1889	CY	15	B	+	-	-	-	-	*
<i>Gastrostyla dorsicirrata</i> Foissner, 1982	HY	96	C,H	+	-	-	-	-	**
<i>Gastrostyla minima</i> Hemberger, 1985	HY	195	B,D,G	-	-	-	+	-	*
<i>Gastrostyla muscorum</i> Kahl, 1932	HY	380	C,G	+	-	-	-	-	**
<i>Gastrostyla philippinensis</i> Shibuya, 1931, synonym with <i>G. steinii</i>									
<i>Gastrostyla steinii</i> Engelmann, 1862	HY	122	C	+	+	+	+	-	*
<i>Glaucoma gigantea</i> Grandori & Grandori, 1934	HM	420	E	+	-	-	-	-	**
<i>Glaucoma terricola</i> Gellért, 1957, synonym with <i>Tetrahymena rostrata</i>									
<i>Gonostomum affine</i> (Stein, 1859) Sterki, 1878	HY	39	B,F,S	+	+	+	+	+	*
<i>Gonostomum algicola</i> Gellért, 1942, synonym with <i>G. affine</i> ⁹⁾									
<i>Gonostomum andoi</i> Shibuya, 1929, synonym with <i>G. affine</i>									
<i>Gonostomum bryonicolum</i> Gellért, 1956, synonym with <i>G. affine</i> ⁹⁾									
<i>Gonostomum ciliophorum</i> Gellért, 1956, synonym with <i>G. affine</i> ⁹⁾									
<i>Gonostomum geleii</i> Gellért, 1957, synonym with <i>G. affine</i> ⁹⁾									
<i>Gonostomum kuehnelti</i> Foissner, 1987	HY	30	B,F	+	+	+	+	+	***
<i>Gonostomum spirotrichoides</i> Gellért, 1956, synonym with <i>G. affine</i> ⁹⁾									
<i>Gonostomum strenua</i> (Engelmann, 1862) Sterki, 1878	HY	60	B,G	+	-	-	-	-	**
<i>Grandoria aculeata</i> (Grandori & Grandori, 1934) Corliss, 1960	CO	5	?	+	-	-	-	-	**
<i>Grossglockneria acuta</i> Foissner, 1980	CO	5	H	+	+	+	+	+	***
<i>Grossglockneria hyalina</i> Foissner, 1985	CO	4	H	+	+	+	+	-	***
<i>Halteria decem sulcata</i> Szabó, 1934, synonym with <i>H. grandinella</i>									
<i>Halteria grandinella</i> (Müller, 1773) Dujardin, 1841	OL	12	B,F,G	+	+	+	+	+	*
<i>Halterioforma caudata</i> Horváth, 1956	PR?	2	B	+	-	-	-	-	**
<i>Haplocaulus terrenus</i> Foissner, 1981	PE	5	B	+	+	+	-	+	**
<i>Hastatella radians</i> Erlanger, 1890	PE	30	B	-	+	-	-	-	*
<i>Hausmanniella discoidea</i> (Gellért, 1956) Foissner, 1984	CO	67	B,C,G,H	+	+	+	+	-	***
<i>Hausmanniella patella</i> (Kahl, 1931) Foissner, 1984	CO	192	C,G	+	+	+	+	-	***
<i>Hausmanniella quinquecircularis</i> (Gellért, 1955) Foissner, 1993	CO	440	G,H	+	-	-	-	-	***
<i>Hemiamphisiella granulifera</i> (Foissner, 1987) Foissner, 1988	HY	52	F,H,N	+	+	+	-	-	**

Table 2. continued.

Species ²⁾	Taxonomic group ³⁾	Biomass of 10 ⁶ indiv. (mg) ⁴⁾	Food ⁵⁾	Geographic distribution ⁶⁾					Degree of autochthonism ⁷⁾
				H	P	A	N	Ar	
<i>Hemiamphisiella quadrinucleata</i> (Foissner, 1984) Foissner, 1988	HY	42	F,G,H	+	-	-	-	-	**
<i>Hemiamphisiella terricola</i> Foissner, 1988	HY	192	C,D,F,H	+	+	+	+	-	**
<i>Hemiamphisiella wilberti</i> (Foissner, 1982) Foissner, 1988	HY	152	T	+	+	+	-	-	**
<i>Hemisincirra gellerti</i> (Foissner, 1982) Foissner, 1984	HY	10	B	+	+	+	+	+	***
<i>Hemisincirra gracilis</i> (Foissner, 1982) Foissner, 1984	HY	11	B?H	+	+	+	+	-	***
<i>Hemisincirra heterocirrata</i> Hemberger, 1985	HY	26	?	-	-	-	+	-	***
<i>Hemisincirra inquieta</i> Hemberger, 1985	HY	7	B	+	+	+	+	-	***
<i>Hemisincirra interrupta</i> (Foissner, 1982) Foissner, 1984	HY	8	?	+	+	+	-	-	***
<i>Hemisincirra kahli</i> (Buitkamp, 1977) Hemberger, 1985	HY	8	B	+	+	-	-	-	***
<i>Hemisincirra muelleri</i> Foissner, 1986	HY	11	B	+	-	-	-	-	***
<i>Hemisincirra octonucleata</i> Hemberger, 1985	HY	24	?	-	-	-	+	-	***
<i>Hemisincirra polynucleata</i> Foissner, 1984	HY	30	?	+	+	-	-	+	***
<i>Hemisincirra pori</i> (Wilbert & Kahan, 1986) Foissner, 1987	HY	19	B,G	+	-	-	-	-	***
<i>Hemisincirra quadrinucleata</i> Hemberger, 1985	HY	26	?	-	-	-	+	-	***
<i>Hemisincirra similis</i> (Foissner, 1982) Foissner, 1984	HY	32	B?	+	+	+	+	-	***
<i>Hemisincirra vermiculare</i> Hemberger, 1985	HY	24	?	-	-	-	+	-	***
<i>Hemisincirra wenzeli</i> Foissner, 1987	HY	14	B,F	+	-	+	+	-	***
<i>Histiculus similis</i> (Quennerstedt) in Buitkamp [27], synonym with <i>Sterkiella cavigola</i>									
<i>Holophrya bimacronucleata</i> Grandori & Grandori, 1934	PR	69	?	+	-	-	-	-	**
<i>Holosticha adami</i> Foissner, 1982	HY	66	C,D,R,S,T	+	-	+	-	-	**
<i>Holosticha australis</i> Blatterer & Foissner, 1988	HY	65	C,T	+	+	+	+	-	**
<i>Holosticha bergeri</i> Foissner, 1987	HY	10	E,F	+	+	+	-	+	***
<i>Holosticha distyla</i> Buitkamp, 1977	HY	136	F,T	-	+	-	-	-	**
<i>Holosticha islandica</i> Berger & Foissner, 1989	HY	30	B,H	+	-	-	-	-	**
<i>Holosticha longiseta</i> Lepsi, 1951	HY	30	?	+	-	-	-	-	**
<i>Holosticha manca mononucleata</i> Gellért, 1955	HY	10	F,H	+	-	-	-	-	**
<i>Holosticha manca plurinucleata</i> Gellért, 1955	HY	26	B,H,S	+	-	-	-	-	**
<i>Holosticha monilata</i> Kahl, 1928	HY	52	B,D,E	+	+	-	-	-	*
<i>Holosticha multistilata</i> Kahl, 1928	HY	109	C,D,E,F,T	+	+	+	+	+	*
<i>Holosticha muscicola</i> Gellért, 1956	HY	60	B,S	+	-	-	-	-	**
<i>Holosticha muscorum</i> (Kahl, 1932) Foissner, 1982	HY	360	C,D,T	+	+	+	+	-	*
<i>Holosticha sigmoidea</i> Foissner, 1982	HY	38	B,H	+	-	+	+	+	**
<i>Holosticha stueberi</i> Foissner, 1987	HY	525	C,F,H,N	+	+	+	-	-	**
<i>Holosticha sylvatica</i> Foissner, 1982	HY	181	C,E,H,S	+	+	+	+	-	***
<i>Holosticha tetricirrata</i> Buitkamp & Wilbert, 1974	HY	39	B,C,D,G,H	+	+	+	+	+	**
<i>Holostichides chardezi</i> Foissner, 1987	HY	63	E,H,S	+	+	+	-	-	**
<i>Holostichides terricola</i> Foissner, 1988	HY	33	B,H,S	+	+	+	+	-	**
<i>Holostichides typicus</i> (Song & Wilbert, 1988) Eigner, 1994	HY	170	?	+	-	-	-	-	**
<i>Holostichides wilberti</i> (Song, 1990) Eigner, 1994	HY	27	B?	+	-	-	-	-	**
<i>Homalogastra setosa</i> Kahl, 1926	HM	1	B	+	+	+	+	+	*
<i>Idiocolpoda pelobia</i> Foissner, 1993	CO	3	B	-	-	+	-	-	**
<i>Ilsiella palustris</i> Foissner, 1993	CO	65	B	+	+	+	-	-	**
<i>Ilsiella venusta</i> Foissner, 1987	CO	20	B	-	+	-	-	-	**
<i>Jaroschia sumptuosa</i> Foissner, 1993	CO	100	C?F?	-	-	+	-	-	***
<i>Kahlia acrobates</i> (Horváth, 1932) Corliss, 1960	HY	?	B,C,G	+	-	-	-	-	**
<i>Kahlia simplex</i> (Horváth, 1934) Corliss, 1960	HY	157	B,D,F,H	+	+	-	-	-	**
<i>Kahlilembus attenuatus</i> (Smith, 1897) Foissner, Berger & Kohmann, 1994	HM	1	B	+	+	+	+	-	*
<i>Kahlilembus fusiformis</i> (Kahl, 1926), synonym with <i>K. attenuatus</i>									
<i>Kalometopia duplicata</i> (Penard, 1922) Foissner, 1993	CO	400	C,G,S	+	-	-	-	-	***
<i>Keronella gracilis</i> Wiackowski, 1985	HY	300	C,E,G,H	+	-	-	-	-	**
<i>Keronopsis algivora</i> (Gellért, 1942) nov. comb.	HY	64	G	+	-	-	-	-	**
<i>Keronopsis dieckmanni</i> Foissner (this paper)	HY	300	C,F	-	+	-	-	-	**
<i>Keronopsis helluo</i> Penard, 1922	HY	800	C,R	+	-	-	-	-	**

Table 2. continued.

Species ²⁾	Taxonomic group ³⁾	Biomass of 10 ⁶ indiv. (mg) ⁴⁾	Food ⁵⁾	Geographic distribution ⁶⁾					Degree of autochthonism ⁷⁾
				H	P	A	N	Ar	
<i>Keronopsis tasmaniensis</i> Blatterer & Foissner, 1988	HY	260	C,H	-	+	+	-	-	**
<i>Keronopsis wetzeli</i> Wenzel, 1953	HY	90	C,G	+	-	-	-	-	**
<i>Keronopsis alpestris</i> Kahl, 1932	HY	96	C,G	+	-	-	-	-	**
<i>Krassnigga auxiliaris</i> Foissner, 1987	CO	880	C	-	+	+	-	-	***
<i>Kreyella muscicola</i> Kahl, 1931	CO	5	?	+	-	-	-	-	**
<i>Kuehneltiella muscicola</i> Foissner, 1993	CO	115	C	+	+	-	+	-	***
<i>Kuehneltiella terricola</i> Foissner, 1990	CO	384	C,F	-	-	+	-	-	***
<i>Kuklikophrya ougandae</i> (Dragesco, 1972) Foissner, 1993	CO	195	E	+	+	-	-	-	*
<i>Lacrymaria cohni</i> Kent in Buitkamp & Wilbert (1974), synonym with <i>Phialina terricola</i> Foissner, 1984									
<i>Lacrymaria pulchra</i> Wenzel, 1953	GY	5	?	+	-	-	-	-	**
<i>Lagynophrya armata</i> Kahl, 1935	GY	20	?	+	-	-	-	-	**
<i>Lagynophrya geleii</i> Foissner, 1981	GY	25	C?	+	-	+	-	-	**
<i>Lagynophrya trichocystis</i> Foissner, 1981	GY	3	C?	+	-	-	-	-	**
<i>Lamostylo abdita</i> Foissner, 1997	HY	27	B,F,H,N	-	-	+	-	-	**
<i>Lamostylo australis</i> (Blatterer & Foissner, 1988) Petz & Foissner, 1996	HY	50	B,C,F,G,H	+	+	+	+	-	***
<i>Lamostylo edaphoni</i> Berger & Foissner, 1987	HY	11	B	+	+	+	+	+	***
<i>Lamostylo granulifera</i> Foissner, 1997	HY	90	C,F,N	+	+	+	+	-	***
<i>Lamostylo hyalina</i> (Berger, Foissner & Adam, 1984) Berger & Foissner, 1987	HY	4	?	+	-	-	+	+	***
<i>Lamostylo islandica</i> Berger & Foissner, 1988	HY	20	B,F	+	+	+	+	+	***
<i>Lamostylo kirkeniensis</i> Berger & Foissner, 1988	HY	20	B,C,F,H	+	-	+	-	-	***
<i>Lamostylo lamottei</i> Buitkamp, 1977	HY	26	C,D	-	+	-	-	-	***
<i>Lamostylo longa</i> (Hemberger, 1985) Berger & Foissner, 1988	HY	135	?	-	-	-	+	-	***
<i>Lamostylo perisincirra</i> (Hemberger, 1985) Berger & Foissner, 1987	HY	7	?	+	+	-	-	+	***
<i>Lamostylo raptans</i> (Hemberger, 1985) Foissner, 1997	HY	150	?	-	-	-	+	-	***
<i>Laurentiella strenua</i> (Dingfelder, 1962) Berger & Foissner, 1989	HY	3000	B,C,D	+	-	+	-	-	*
<i>Leptopharynx costatus</i> Mermod, 1914	NA	5	B,F	+	+	+	+	+	*
<i>Leptopharynx eurystoma</i> (Kahl, 1931) nov. comb.	NA	10	E?	+	-	+	+	-	*
<i>Leptopharynx macrostoma</i> Njine, 1979, synonym with <i>L. eurystoma</i>									
<i>Leptopharynx minimus</i> Alekperov, 1993, synonym with <i>L. costatus</i>									
<i>Leptopharynx stenostoma</i> (Gellért, 1942) nov. comb., synonym with <i>L. costatus</i>									
<i>Linostoma vorticella</i> (Ehrenberg, 1833) Jankowski, 1978	HE	1000	C,D,G	-	+	-	-	-	*
<i>Litonotus crinitus</i> Grandori & Grandori, 1934	GY	40	C?	+	-	-	-	-	**
<i>Litonotus digitatus</i> Grandori & Grandori, 1934	GY	39	C?	+	-	-	-	-	**
<i>Litonotus muscorum</i> (Kahl, 1931) Blatterer & Foissner, 1988	GY	12	C	+	+	+	+	-	**
<i>Maryna acuminata</i> (Gellért, 1955) Foissner, 1993	CO	1	B	+	-	-	-	-	**
<i>Maryna antarctica</i> Foissner, 1993	CO	7	B	+	-	-	-	+	**
<i>Maryna atra</i> (Gelei, 1950) Foissner, 1993	CO	180	B	+	+	-	-	-	*
<i>Maryna lichenicola</i> (Gelei, 1950) Foissner, 1993	CO	9	B	+	+	-	-	-	**
<i>Maryna minima</i> (Gelei, 1950) Foissner, 1993	CO	7	?	+	-	-	-	-	*
<i>Maryna ovata</i> (Gelei, 1950) Foissner, 1993	CO	40	B	+	+	-	+	-	**
<i>Maryna pinguis</i> Dingfelder, 1962	CO	270	?	+	-	-	-	-	**
<i>Maryna rotunda</i> Dingfelder, 1962	CO	113	?	-	+	-	-	-	**
<i>Maryna socialis</i> Gruber, 1879	CO	830	?	+	-	-	-	-	*
<i>Maryna umbrellata</i> (Gelei, 1950) Foissner, 1993	CO	523	B,S?	-	+	-	-	-	**
<i>Metacineta micraster</i> (Penard, 1914) Batisse, 1967	SU	33	C	+	-	-	-	-	*
<i>Metacineta mystacina</i> (Ehrenberg, 1831) Bütschli, 1889	SU	65	C	+	-	-	-	-	*
<i>Metopus contractus</i> Penard, 1922	HE	100	B	-	-	-	+	-	*
<i>Metopus es</i> (Müller, 1776) Lauterborn, 1916	HE	95	B,E,F,S	-	+	-	+	-	*
<i>Metopus basei</i> Sondheim, 1929	HE	14	B	+	+	+	+	-	*
<i>Metopus ovalis</i> Kahl, 1927	HE	240	B	-	+	-	+	-	*
<i>Metopus palaeformis</i> Kahl, 1927	HE	10	B	-	-	-	+	-	*

Table 2. continued.

Species ²⁾	Taxonomic group ³⁾	Biomass of 10 ⁶ indiv. (mg) ⁴⁾	Food ⁵⁾	Geographic distribution ⁶⁾					Degree of autochthonism ⁷⁾
				H	P	A	N	Ar	
<i>Metopus rectus</i> (Kahl, 1932) Foissner, 1980	HE	72	B	+	-	-	-	-	*
<i>Metopus setosus</i> Kahl, 1927	HE	32	B	-	-	-	+	-	*
<i>Metopus setosus minor</i> Kahl, 1927	HE	18	B	-	-	-	+	-	*
<i>Microdiaphanosoma arcuatum</i> (Grandori & Grandori, 1934) Wenzel, 1953	CO	0.5	B	+	+	+	+	+	***
<i>Microdiaphanosoma terricola</i> Foissner, 1993	CO	0.5	?	+	-	+	-	-	***
<i>Microthorax simulans</i> (Kahl, 1926) Kahl, 1931	NA	0.7	B	+	+	+	+	+	**
<i>Monodinium balbiani</i> Fabre-Domergue, 1888	GY	55	C,G	+	+	-	+	-	*
<i>Mycterothrix tuamotuensis</i> (Balbiani, 1887) Kahl, 1931	CO	20	B	+	-	+	-	-	***
<i>Mykophagophrys terricola</i> (Foissner, 1985) Foissner, 1995	CO	3	H	+	+	+	+	+	***
<i>Nassula citrea</i> Kahl, 1930	NA	150	E	+	+	-	-	-	*
<i>Nassula picta</i> Greeff, 1888	NA	224	E	+	+	-	-	+	*
<i>Nassula protectissima</i> Penard, 1922	NA	40	E	+	-	-	-	-	**
<i>Nassula terricola</i> Foissner, 1989	NA	180	E	+	-	-	-	-	**
<i>Nassula tumida obscura</i> Lepsi, 1951	NA	50	?	+	-	-	-	-	**
<i>Neogeneia hortualis</i> Eigner, 1995	HY	38	B	+	-	-	-	-	**
<i>Nivaliella plana</i> Foissner, 1980	CO	0.5	H	+	+	+	+	+	***
<i>Notohymena antarctica</i> Foissner, 1996	HY	45	B?C,F	-	-	-	-	+	**
<i>Notohymena australis</i> (Foissner & O'Donoghue) Blatterer & Foissner, 1988	HY	45	B	-	-	-	+	-	*
<i>Notohymena rubescens</i> Blatterer & Foissner, 1988	HY	40	F,H,N	+	-	+	+	-	**
<i>Notohymena selvatica</i> (Hemberger, 1985) Blatterer & Foissner, 1988	HY	560	?	-	+	-	+	-	**
<i>Notoxoma parabryophryides</i> Foissner, 1993	CO	10	B	+	+	+	+	-	***
<i>Notoxoma sigmoides</i> Foissner, 1993	CO	3	B	-	+	-	-	-	***
<i>Obertrumia kahli</i> Foissner, 1989	NA	180	E	+	-	-	-	-	**
<i>Odontochlamys alpestris</i> Foissner, 1981	CY	10	B,F	+	+	+	+	-	*
<i>Odontochlamys convexa</i> (Kahl, 1931) Blatterer & Foissner, 1992	CY	9	B?	+	+	+	+	-	***
<i>Odontochlamys gouraudi</i> Certes, 1891	CY	13	B	+	+	+	+	-	*
<i>Odontochlamys wisconsinensis</i> (Kahl, 1931) Petz & Foissner, 1997	CY	6	B,G?	+	-	-	-	+	**
<i>Onychodromopsis flexilis</i> Stokes, 1887	HY	80	B,C,F	+	-	-	-	+	*
<i>Opercularia arboricola</i> (Biegel, 1954) Foissner, 1981, synonym with <i>O. curvicaule</i>									
<i>Opercularia arenicola</i> Greeff, 1873	PE	?	?	+	-	-	-	-	**
<i>Opercularia asymmetrica</i> (Biczok, 1956) Aesch & Foissner, 1992	PE	13	B	+	-	-	-	-	*
<i>Opercularia curvicaule</i> (Penard, 1922) nov. comb.	PE	48	B	+	+	+	+	+	**
<i>Ophryoglena marginata</i> Greeff, 1888	HM	?	?	+	-	-	-	-	**
<i>Opisthonecta minima</i> Foissner, 1975	PE	143	B	+	+	-	+	-	*
<i>Opisthotricha elongata</i> Grandori & Grandori, 1934	HY	20	?	+	-	-	-	-	**
<i>Opisthotricha procera</i> Kahl, 1932	HY	20	?	+	-	-	-	-	**
<i>Opisthotricha terrestris</i> Horváth, 1956, synonym with <i>Sterkiella histriomuscorum</i>									
<i>Opisthotricha terricola</i> Gellért, 1957	HY	24	B	+	-	-	-	-	**
<i>Orthoamphisiella franzi</i> (Foissner, 1982) Eigner, 1995	HY	30	B,F,S	+	-	-	-	-	*
<i>Orthoamphisiella grelli</i> Eigner & Foissner, 1993	HY	20	B,F	-	-	-	-	+	**
<i>Orthoamphisiella stramenticola</i> Eigner & Foissner, 1991	HY	60	B?C,E,H	+	-	-	+	-	**
<i>Orthokreyella schiffmanni</i> Foissner, 1984	CO	0.2	B?	+	-	-	-	-	***
<i>Oxytricha auripunctata</i> Blatterer & Foissner, 1988	HY	40	C,F,H	-	+	+	+	-	**
<i>Oxytricha bimembranata</i> Shibuya, 1929	HY	300	?	+	-	-	-	-	**
<i>Oxytricha crassistilata</i> Kahl in Alekperov [4], synonym with <i>Sterkiella histriomuscorum</i>									
<i>Oxytricha gigantea</i> Horváth, 1933	HY	270	B,C,F,H	+	+	-	-	-	*
<i>Oxytricha granulifera</i> Foissner & Adam, 1983	HY	72	B,F,H	+	+	+	+	+	**
<i>Oxytricha granulifera quadricirrata</i> Blatterer & Foissner, 1988	HY	30	B,F,H	-	+	+	+	-	**
<i>Oxytricha hengshanensis</i> Shen, Liu, Song & Gu, 1992	HY	252	G,N	+	-	-	-	-	**

Table 2. continued.

Table 2. continued.

Species ²⁾	Taxonomic group ³⁾	Biomass of 10 ⁶ indiv. (mg) ⁴⁾	Food ⁵⁾	Geographic distribution ⁶⁾					Degree of autochthonism ⁷⁾
				H	P	A	N	Ar	
<i>Platyophrya macrostoma</i> Foissner, 1980	CO	1.5	B	+	+	+	+	+	**
<i>Platyophrya paoletti</i> Foissner, 1997	CO	3	B	-	-	-	+	-	***
<i>Platyophrya similis</i> (Foissner, 1980) Foissner, 1987	CO	180	C,D,F,G	+	-	+	+	-	**
<i>Platyophrya spumacola</i> Kahl, 1927	CO	40	B,C,E,F,G, H,R,T	+	+	+	+	+	**
<i>Platyophrya vorax</i> Kahl, 1926	CO	12	B,F,G,H	+	+	+	+	+	*
<i>Platyophryides dragescoi</i> Foissner, 1987	CO	64	C	+	-	+	+	+	***
<i>Platyophryides latus</i> (Kahl, 1930) Foissner, 1987	CO	108	C,G,H	+	-	+	+	-	***
<i>Platyophryides magnus</i> Foissner, 1993	CO	700	C	-	+	-	-	-	***
<i>Pleuroplites australis</i> Foissner, 1988	GY	8	C,F	+	+	+	+	+	***
<i>Pleuroplitooides smithi</i> Foissner, 1996	GY	56	C	+	+	+	+	+	***
<i>Podophrya halophila</i> Kahl, 1934	SU	17	C	-	-	+	+	-	*
<i>Protocyclidium terrenum</i> Alekperov, 1993	HM	2	B	+	-	-	-	-	**
<i>Protospathidium bonnetti</i> (Buitkamp, 1977) Foissner, 1981	GY	5	C	+	+	+	+	-	***
<i>Protospathidium muscicola</i> Dragesco & Dragesco-Kernéis, 1979	GY	7	F	+	+	-	+	-	**
<i>Protospathidium serpens</i> (Kahl, 1930) Foissner, 1981	GY	5	C?	+	+	+	+	+	**
<i>Protospathidium terricola</i> Foissner (this paper)	GY	28	C?	-	+	-	-	+	**
<i>Pseudocarchesium claudicans</i> (Penard, 1922) Foissner, 1989	PE	3	B	+	-	+	+	-	**
<i>Pseudochilodonopsis mutabilis</i> Foissner, 1981	CY	8	B	+	+	+	+	+	**
<i>Pseudochilodonopsis polyvacuolata</i> Foissner & Didier, 1981	CY	30	B	+	+	-	-	-	*
<i>Pseudocohnilembus marinus</i> Thompson, 1966	HM	3	B?	-	-	+	-	-	*
<i>Pseudocohnilembus persalinus</i> Evans & Thompson, 1964, synonym with <i>P. pusillus</i>									
<i>Pseudocohnilembus pusillus</i> (Quennerstedt, 1869)									
Foissner & Wilbert, 1981	HM	3	B	+	-	-	-	-	*
<i>Pseudocohnilembus putrinus</i> (Kahl, 1928)	HM	2	B	+	+	+	+	-	*
Foissner & Wilbert, 1981									
<i>Pseudocohnilembus veisovi</i> Alekperov & Musaev, 1988, synonym with <i>P. pusillus</i>									
<i>Pseudocristigera hymenofera</i> Horváth, 1956, synonym with <i>Drepanomonas revoluta</i>									
<i>Pseudocyrtolophosis alpestris</i> Foissner, 1980	CO	2	B	+	+	+	+	+	**
<i>Pseudocyrtolophosis terricola</i> Foissner, 1993	CO	3	B	-	+	+	+	-	**
<i>Pseudoglaucoma muscorum</i> (Kahl, 1931) Wenzel, 1953	CO	1	?	+	-	-	-	-	**
<i>Pseudoholophrya terricola</i> Berger, Foissner & Adam, 1984	GY	32	C?	+	+	+	+	+	**
<i>Pseudokreyella terricola</i> Foissner, 1985	CO	3	B?	+	+	+	-	+	***
<i>Pseudokreyella australis</i> Foissner, 1993	CO	3	B	+	-	+	-	-	***
<i>Pseudomicrothorax agilis</i> Mermod, 1914	NA	14	B,E,G	-	+	-	+	-	*
<i>Pseudomicrothorax dubius</i> (Maupas, 1883) Penard, 1922	NA	50	B,E	-	+	-	-	-	*
<i>Pseudoplatyophrya nana</i> (Kahl, 1926) Foissner, 1980	CO	3	H	+	+	+	+	+	***
<i>Pseudoplatyophrya saltans</i> Foissner, 1988	CO	2	H	+	+	+	+	+	***
<i>Pseudothuricola dionysii</i> (Penard, 1922) Kahl, 1935	PE	9	B	+	-	-	-	-	**
<i>Pseudouroleptus buitkampi</i> (Foissner, 1982)	HY	72	B,C,S	+	-	-	-	-	***
Berger & Foissner, 1987									
<i>Pseudouroleptus procerus</i> Berger, Foissner, 1987	HY	120	C,F,G,N	+	+	-	-	-	***
<i>Pseudouroleptus terrestris</i> Hemberger, 1985	HY	330	?	-	-	-	+	-	**
<i>Pseudourostyla franzi</i> Foissner, 1987	HY	375	C,H,N,T	-	+	+	+	-	**
<i>Pseudovorticella mutans</i> (Penard, 1922) Foissner, 1979	HY	14	B	+	-	-	-	-	*
<i>Pseudovorticella sphagni</i> Foissner & Schiffmann, 1974	PE	14	B	+	+	+	+	-	*
<i>Psilotricha succisa</i> (Müller, 1786) Foissner, 1983	HY	66	G	-	-	+	-	-	*
<i>Pyxidium invaginatum muscorum</i> Kahl, 1935, synonym with <i>Opercularia curvicaule</i>									
<i>Pyxidium longicollum</i> Biegel, 1954	PE	7	?	+	-	-	-	-	**
<i>Pyxidium tardigradum</i> Van der Land, 1964	PE	7	B?	+	-	-	-	-	**
<i>Reticulowoodruffia terricola</i> Foissner, 1993	CO	54	?	+	-	-	-	-	***
<i>Rhabdostyla (?) arborea</i> Greeff, 1888	PE	?	?	+	-	-	-	-	**
<i>Rhabdostyla muscorum</i> Kahl, 1935	PE	6	?	+	-	-	-	-	**
<i>Rhabdotricha terricola</i> Greeff, 1888	HY	?	?	+	-	-	-	-	**

Table 2. continued.

Species ²⁾	Taxonomic group ³⁾	Biomass of 10 ⁶ indiv. (mg) ⁴⁾	Food ⁵⁾	Geographic distribution ⁶⁾					Degree of autochthonism ⁷⁾
				H	P	A	N	Ar	
<i>Rhopalophrya elegans</i> Horváth, 1956	GY	2	B,F	+	-	-	-	-	**
<i>Rhopalophrya pentacerca</i> Grandori & Grandori, 1934	GY	4	?	+	-	-	-	-	**
<i>Rostrophrya camerounensis</i> (Njine, 1979) Foissner, 1993	CO	640	D,E	-	+	-	-	-	*
<i>Rostrophrya terricola</i> Foissner, 1993	CO	360	?	+	-	+	-	-	***
<i>Rostrophryides africana</i> Foissner, 1987	CO	14	?	+	+	-	+	-	***
<i>Rostrophryides australis</i> Blatterer & Foissner, 1988	CO	30	B,G,H	+	-	+	+	-	***
<i>Sagittaria australis</i> Pomp & Wilbert, 1988	CO	9	?	-	-	+	-	-	**
<i>Sagittaria hyalina</i> Foissner, Czapik & Wiackowski, 1981	CO	4	B	+	+	+	+	-	*
<i>Sagittaria polygonalis</i> Grandori & Grandori, 1934	CO	5	?	+	-	-	-	-	**
<i>Sathrophilus muscorum</i> (Kahl, 1931) Corliss, 1960	HM	12	B	+	+	+	+	+	**
<i>Sathrophilus simonis</i> (Lepsi, 1948) Corliss, 1960	HM	6	?	+	-	-	-	-	**
<i>Semiplatyophrya foissneri</i> Wilbert & Kahan, 1986	CO	13	B	+	+	+	-	-	**
<i>Solenophrya flavescens</i> Penard, 1914	SU	33	C	+	-	-	-	-	**
<i>Solenophrya massula</i> Penard, 1914	SU	3	C	+	-	-	-	-	**
<i>Solenophrya sacculus</i> Penard, 1914	SU	5	C	+	-	-	-	-	**
<i>Sorogena stoianovitchae</i> Bradbury & Olive, 1980	CO	54	C	+	+	+	+	-	**
<i>Spathidium alpinum</i> Gellért, 1956	GY	30	C	+	-	-	-	-	**
<i>Spathidium anguilla</i> Vuxanovici, 1962	GY	25	C?	+	+	+	+	-	*
<i>Spathidium bavaricense</i> Kahl, 1930	GY	130	C	+	+	+	+	-	**
<i>Spathidium claviforme</i> Kahl, 1930	GY	31	C	+	+	+	+	-	**
<i>Spathidium falciforme</i> (Penard, 1922) Kahl, 1930	GY	4	?	+	-	-	-	-	**
<i>Spathidium furcatum</i> Shibuya, 1930	GY	60	C?	+	-	-	-	-	**
<i>Spathidium geobium</i> Lepsi, 1951	GY	10	C?	+	-	-	-	-	**
<i>Spathidium holsatiae</i> Kahl, 1930	GY	45	C?	+	-	-	-	-	**
<i>Spathidium lagyniforme</i> Kahl, 1930	GY	34	C?	+	-	-	-	-	*
<i>Spathidium longicaudatum</i> (Buitkamp & Wilbert, 1974) Buitkamp, 1977	GY	18	C,N	+	+	+	+	-	***
<i>Spathidium metabolicum</i> Pomp & Wilbert, 1988, synonym with <i>S. anguilla</i>									
<i>Spathidium multinucleatum</i> Gellért, 1955	GY	23	C	+	-	-	-	-	**
<i>Spathidium muscicola</i> Kahl, 1930	GY	84	C	+	+	+	+	-	**
<i>Spathidium procerum</i> Kahl, 1930	GY	15	C?	+	+	+	+	-	**
<i>Spathidium rusticum</i> Foissner, 1981	GY	25	C?	+	+	-	-	-	**
<i>Spathidium scalpriforme</i> (Kahl, 1930) Kahl, 1930	GY	220	C?F	+	-	-	-	-	**
<i>Spathidium seppelti</i> Petz & Foissner, 1997	GY	65	C	-	-	-	-	+	**
<i>Spathidium spathula</i> (Müller, 1773) Moody, 1912	GY	45	C	+	+	+	+	-	*
<i>Spathidium spathuloides</i> Gellért, 1955	GY	6	C	+	-	-	-	-	**
<i>Spetazoon austriense</i> Foissner, 1994	GY	200	C	-	+	+	-	-	**
<i>Sphaerophrya parva</i> Greeff, 1888	SU	?	?	+	-	-	-	-	**
<i>Sphaerophrya terricola</i> Foissner, 1986	SU	8	C	+	+	+	+	+	**
<i>Spirofilopsis tubicola</i> (Gruber, 1879) Dingfelder, 1962	HY	30	?	+	-	-	-	-	**
<i>Stammeridium kabli</i> (Wenzel, 1953) Wenzel, 1969	NA	0.6	B	+	+	+	+	-	***
<i>Stegochilum smalli</i> Alekperov, 1993, synonym with <i>Tetrahymena rostrata</i>									
<i>Steinia platystoma</i> (Ehrenberg, 1831) Diesing, 1866	HY	175	C,F,G,D	+	+	-	-	-	*
<i>Sterkiella caricola</i> (Kahl, 1935) Foissner et al., 1991 (formerly <i>Histiculus</i>)	HY	540	C,N	+	+	+	+	-	**
<i>Sterkiella histriomuscorum</i> (Foissner et al., 1991) Foissner et al., 1991 (formerly <i>Histiculus muscorum</i>)	HY	72	B,C,F,N	+	+	+	+	+	*
<i>Sterkiella similis</i> f. <i>tricirrata</i> (Buitkamp, 1977) nov. comb.	HY	234	B,E	-	+	-	-	-	**
<i>Sterkiella thompsoni</i> Foissner, 1996	HY	90	B,D,F,G	-	-	-	-	+	*
<i>Stichotricha aculeata</i> Wrześniowski, 1866	HY	20	B,G	+	+	+	-	-	*
<i>Stichotricha socialis</i> Gruber, 1879	HY	63	?	+	-	-	-	-	*
<i>Strongylidium californicum</i> Kahl, 1932	HY	100	?	+	-	-	-	-	**
<i>Strongylidium muscorum</i> Kahl, 1932	HY	30	?	+	-	-	-	-	**
<i>Styloynchia bifaria</i> (Stokes, 1887) nov. comb. ⁸⁾	HY	100	B,C	+	-	-	+	-	*
<i>Styloynchia mytilus</i> (Müller, 1773) Ehrenberg, 1830	HY	70	B,C,G,Z	+	-	+	+	-	*

Table 2. continued.

Species ²⁾	Taxonomic group ³⁾	Biomass of 10 ⁶ indiv. (mg) ⁴⁾	Food ⁵⁾	Geographic distribution ⁶⁾					Degree of autochthonism ⁷⁾
				H	P	A	N	Ar	
<i>Styloynchia pustulata</i> (Müller, 1786) Ehrenberg, 1835	HY	80	B,C,F,G	-	+	+	-	-	*
<i>Styloynchia quadrinucleata</i> Alekperov & Musaev, 1988	HY	3500	C	+	-	-	-	-	**
<i>Styloynchia stylomuscorum</i> (Foissner, Blatterer, Berger & Kohmann, 1991) Foissner, Blatterer, Berger & Kohmann, 1991	HY	30	D,F,G	+	-	-	-	-	*
<i>Tachysoma granulifera</i> Berger & Foissner, 1987	HY	26	F,G,H,N	+	+	+	-	-	***
<i>Tachysoma humicola</i> Gellért, 1957	HY	9	B,T	+	+	+	+	-	***
<i>Tachysoma humicola longisetum</i> Foissner (this paper)	HY	12	B	-	+	-	+	-	***
<i>Tachysoma terricola</i> Hemberger, 1985	HY	45	?	-	-	-	+	-	***
<i>Tectohymena terricola</i> Foissner, 1993	CO	10	B	+	+	-	-	-	***
<i>Telostomatella ferroii</i> (Grandori, 1935) Foissner, 1985	CO	11	E,F	+	-	-	-	-	**
<i>Telotrochidium cylindricum</i> Foissner, 1978	PE	66	B	+	+	-	-	-	*
<i>Terricirra livida</i> (Berger & Foissner, 1987) Berger & Foissner, 1989	HY	24	B	+	+	+	-	-	***
<i>Terricirra matsusakai</i> Berger & Foissner, 1989	HY	45	B	+	+	+	+	-	***
<i>Terricirra viridis</i> (Foissner, 1982) Berger & Foissner, 1989	HY	33	B	+	+	+	-	-	***
<i>Territricha stramenticola</i> Berger & Foissner, 1988	HY	100	C,H	+	-	-	-	-	**
<i>Tetrahymena edaphoni</i> Foissner, 1987	HM	8	B	+	-	-	-	-	**
<i>Tetrahymena rostrata</i> (Kahl, 1926) Corliss, 1952	HM	12	B,G	+	+	+	+	-	**
<i>Thuricola innixa</i> Stokes, 1882	PE	70	B	+	-	-	-	-	*
<i>Thuricola kellicottiana</i> (Stokes, 1887) Kahl, 1935	PE	200	E	+	-	-	-	-	*
<i>Thylakidium macrostomum</i> Alekperov, 1991, synonym with <i>Bryometopus sphagni</i>									
<i>Thylakidium typicum</i> Gellért, 1955	CO	175	B	+	-	-	-	-	**
<i>Tokophrya muscicola</i> Penard, 1914	SU	7	C	+	-	-	+	-	**
<i>Trachelochaeta gonostomoida</i> Hemberger, 1985	HY	204	?	-	-	-	+	-	**
<i>Trachelophyllum apiculatum</i> (Perty, 1852) Claparède & Lachmann, 1859	GY	39	C	+	+	+	+	-	*
<i>Trachelostyla canadensis</i> Buitkamp & Wilbert, 1974, synonym with <i>Gonostomum algicola</i>									
<i>Transitella lichenicola</i> Gellért, 1950, synonym with <i>Balantidioides bivacuolata</i>									
<i>Tricoronella pulchra</i> Blatterer & Foissner, 1988	HY	540	C,F,H,N,S,T	-	+	-	-	-	***
<i>Trihydromena terricola</i> Foissner, 1988	CO	2	B?	+	+	+	+	-	***
<i>Trithigmostoma bavariensis</i> (Kahl, 1931) Foissner, 1987	CY	96	B	+	+	-	+	-	**
<i>Urliella terricola</i> Foissner, 1989	NA	6	B	+	-	-	-	-	**
<i>Uroleptoides kibni</i> Wenzel, 1953	HY	15	?	+	-	-	-	-	*
<i>Uroleptoides qingdaensis</i> Song & Wilbert, 1989, synonym with <i>Hemiamphisiella terricola</i>									
<i>Uroleptus humicola</i> Gellért, 1956	HY	37	S	+	-	-	-	-	**
<i>Uroleptus lepisma</i> (Wenzel, 1953) nov. comb.	HY	120	B,C,F,H,N,T	+	+	+	+	-	**
<i>Uroleptus matthesi</i> Wenzel, 1953	HY	15	?	+	-	-	-	-	**
<i>Uroleptus musculus</i> (Kahl, 1932) Foissner et al., 1991	HY	214	B,C,G	-	+	-	-	-	*
<i>Uroleptus notabilis</i> (Foissner, 1982) nov. comb.	HY	32	F,H,T	+	+	+	+	+	**
<i>Uronema nigricans</i> (Müller, 1786) Florentin, 1901	HM	5	B,F	+	+	-	-	+	*
<i>Urosoma acuminata</i> (Stokes, 1887) Kahl, 1932	HY	73	D,H,G	+	+	+	+	-	*
<i>Urosoma cienkowskii</i> Kowalewski, 1882	HY	34	B,C,F,H	+	+	-	-	-	*
<i>Urosoma karinae</i> Foissner, 1987 nom. corr.	HY	60	B?F	+	+	+	-	-	**
<i>Urosoma macrostoma</i> Gellért, 1957	HY	14	S	+	-	-	-	-	**
<i>Urosoma macrostylia</i> (Wrześniowski, 1866) Kahl, 1932	HY	41	B	+	+	+	+	-	*
<i>Urosoma octonucleata</i> Berger & Foissner, 1989	HY	67	B	+	-	-	-	-	**
<i>Urosomoida agiliformis</i> Foissner, 1982	HY	30	B,C	+	+	+	+	-	**
<i>Urosomoida agilis</i> (Engelmann, 1862) Hemberger, 1985	HY	30	B,D,N,S,T	+	+	+	+	-	*
<i>Urosomoida antarctica</i> Foissner, 1996	HY	25	B,F?N?	-	-	-	-	+	**
<i>Urosomoida dorsiincisura</i> Foissner, 1982	HY	38	B,C,N	+	+	-	+	-	**
<i>Urosomoida granulifera</i> Foissner, 1996	HY	28	B?	-	-	-	-	+	**
<i>Urosomoida minima</i> Hemberger, 1985	HY	12	?	-	-	-	+	-	**
<i>Urostyla grandis</i> Ehrenberg, 1830	HY	500	C,G,R,T	+	-	-	-	-	*
<i>Urostyla muscorum</i> Kahl, 1932	HY	630	C,R	+	-	-	-	-	**
<i>Urotricha atypica</i> Alekperov, 1993	PR	44	?	+	-	-	-	-	**

Table 2. continued.

Species ²⁾	Taxonomic group ³⁾	Biomass of 10 ⁶ indiv. (mg) ⁴⁾	Food ⁵⁾	Geographic distribution ⁶⁾					Degree of autochthonism ⁷⁾
				H	P	A	N	Ar	
<i>Urotricha mamilla</i> Lepsi, 1951	PR	2	?	+	-	-	-	-	**
<i>Urotricha terricola</i> Alekperov & Musayev, 1988	PR	20	?	+	-	-	-	-	**
<i>Vaginicola chaperoni</i> (Penard, 1914) Kahl, 1935	PE	30	B	+	-	-	-	-	**
<i>Vaginicola doliolum</i> (Penard, 1914) Kahl, 1935	PE	9	B	+	-	-	-	-	**
<i>Vaginicola terricola</i> Greeff, 1888	PE	22	B	+	-	-	-	+	**
<i>Vaginicola virgula</i> (Penard, 1914) Kahl, 1935	PE	8	B	+	-	-	-	-	**
<i>Vorticella astyliformis</i> Foissner, 1981	PE	13	B,S	+	+	+	+	+	**
<i>Vorticella coeni</i> Lepsi, 1948	PE	160	?	+	-	-	-	-	**
<i>Vorticella infusionum</i> Dujardin, 1841	PE	18	B	+	+	+	-	+	*
<i>Vorticella lichenicola</i> Greeff, 1888	PE	23	B	+	-	-	-	-	**
<i>Vorticella microstoma</i> Ehrenberg, 1830	PE	30	B,G	+	-	-	+	-	*
<i>Vorticella muralis</i> Penard, 1922	PE	63	B	+	-	-	-	-	**
<i>Vorticella operculariformis</i> Foissner, 1979	PE	40	B	+	+	-	-	-	*
<i>Vorticella pileolata</i> Lepsi, 1948	PE	20	?	+	-	-	-	-	**
<i>Vorticella similis</i> Stokes, 1887	PE	75	B	+	+	+	+	-	*
<i>Wallacia bujoreana</i> (Lepsi, 1951) Berger & Foissner, 1989	HY	15	B	+	-	-	-	-	**
<i>Woodruffia australis</i> Foissner, 1993	CO	25	B?	-	+	+	+	-	***
<i>Woodruffia lichenicola</i> Gellért, 1955, synonym with <i>Platyophryides latus</i>									
<i>Woodruffia rostrata</i> Kahl, 1931	CO	126	C	+	-	+	-	-	*
<i>Woodruffia sinistromembranellata</i> Gellért, 1955, synonym with <i>Platyophrya spumacola</i>									
<i>Woodruffides metabolicus</i> (Johnson & Larson, 1938) Foissner, 1987	CO	1000	C	+	+	+	+	-	**
<i>Woodruffides terricola</i> Foissner, 1987	CO	110	?	+	+	-	-	-	***

¹⁾ This list contains (i) all species originally described from terrestrial habitats (soil, litter, terrestrial mosses etc.) and (ii) species originally described from freshwater habitats but later reliably recorded (by silver impregnation) from terrestrial environments. True *Sphagnum* species and species from ephemeral puddles were excluded, if not later reliably recorded from terrestrial habitats. Colpodids were selected from Foissner's [84] monograph. As far as it concerns my papers, this summary supersedes my earlier species lists.

²⁾ For literature of original species descriptions, recent combinations, important redescriptions, and synonymies (most subjective), see Table 3.

³⁾ Classification mainly after [34], that of colpodids after [84]. CO = colpodid, CY = cyrtophorid, GY = gymnostomatid (haptorid), HE = heterotrich, HM = hymenostome, HY = hypotrich, NA = nassulid, OL = oligotrich, PE = peritrich, PR = prostomatid, SU = suctorian.

⁴⁾ Rough estimation obtained by reducing the shape of the cells to simple geometric figures and assuming a specific gravity of 1.

⁵⁾ Food items were determined mainly by examination of the food vacuoles. B = bacteria, C = ciliates, D = diatoms, E = blue green algae (cyanobacteria), F = colourless flagellates, G = green algae, including autotrophic flagellates, H = hyphae and/or spores of fungi and yeasts, N = naked amoebae, R = rotifers and/or nematodes, S = inorganic and organic soil particles ("detritus"), T = testate amoebae.

⁶⁾ Classification of biogeographic regions, see Fig. 1. H = Holarctis (North America, Greenland, Eurasia with Iceland, Canary Islands, Korea, Japan, and north Africa), P = Palaeotropis (Africa south of Sahara desert, Madagascar, India), A = Australis (mainly Australia), N = Neotropis (Central and South America), Ar = Archinotis (Antarctica and islands in the southern oceans). Most are original records published here for the first time (see text). Note that freshwater and/or marine records are not included.

⁷⁾ * low, reliably recorded also from freshwater habitats; ** probably strong; includes most of the new species and many moss inhabitants; *** probably found exclusively in true terrestrial habitats (litter, soil, humus under and in moss etc.). Only species with specific food requirements or very characteristic morphological adaptations have been classified to this level; furthermore, species belonging to genera known from soil only, were usually also classified to this level.

⁸⁾ Misidentified as *Dileptus monilatus* Stokes, 1886 by Song [205] according to [98]; misidentified as *Styloynchia vorax* by [245]; misidentified by Song [206] as *Frontonia acuminata*.

⁹⁾ Synonymy questionable.

¹⁰⁾ For *Thylakidium magnum* Alekperov, 1991 [3], which becomes, after transfer to *Bryometopus*, a secondary homonym of *B. magnus* Foissner, 1980 [61], which is, however, very likely a junior synonym of *B. sphagni* Penard, 1922 [84].

¹¹⁾ Note that the genera *Dileptus* and *Euplates* were split in several genera by some authors [98, 256].

Table 3. Quality of descriptions (+ poor; ++ incomplete, i.e. either life aspect or infraciliature insufficiently described; +++ excellent, i.e. original description and/or redescriptions include morphometry and appropriate figures from live and silver prepared specimens) and key literature. Usually, the original reference is provided, except for those species treated in our recent monographs [84, 114–117].

Species	Species
<i>Acaryophryxa collaris</i> +++ [38, 68, 148, 149, 152, 191]	<i>Bresslaua sicaria</i> + [84]
<i>Acineria uncinata</i> +++ [107, 117, 207, 232]	<i>Bresslaua vorax</i> +++ [84, 154]
<i>Acineta flava</i> +++ [117]	<i>Bresslauides australis</i> +++ [20, 84]
<i>Acropisthium mutabile</i> +++ [69, 152, 181]	<i>Bresslauides discoideus</i> +++ [84, 154]
<i>Actinobolina vorax</i> ++ [117, 152]	<i>Bresslauides terricola</i> +++ [75, 84]
<i>Alinostoma multivacuolatum</i> ++ [4]	<i>Bryometopus alekperovi</i> ++ [3]
<i>Amphibotrella enigmatica</i> + [132, 133]	<i>Bryometopus atypicus</i> +++ [61, 84]
<i>Amphisella binucleata</i> +++ [11, 78, 138]	<i>Bryometopus balantidioides</i> +++ [84]
<i>Amphisella magnigranulosa</i> +++ [78]	<i>Bryometopus edaphonus</i> + [28, 61, 84]
<i>Amphisella polycirrata</i> +++ [11]	<i>Bryometopus hawaiiensis</i> +++ [89]
<i>Amphisella quadrinucleata</i> +++ [11]	<i>Bryometopus pseudochilodon</i> +++ [84, 155]
<i>Amphisella raptans</i> ++ [28]	<i>Bryometopus sphagni</i> +++ [3, 84, 155, 179]
<i>Amphisella terricola</i> +++ [69, 128]	<i>Bryometopus triquetrus</i> +++ [84]
<i>Amphisella vitiphila</i> +++ [75, 78]	<i>Bryophryxa bavaricensis</i> +++ [84, 154, 240]
<i>Amphisellides atypicus</i> ++ [78, 138]	<i>Bryophryxa rubescens</i> ++ [84, 179]
<i>Amphisellides illuvialis</i> +++ [49]	<i>Bryophyllum loxophylliforme</i> ++ [154]
<i>Apoamphisella tihanyiensis</i> +++ [97, 131]	<i>Bryophyllum tegularum</i> +++ [69, 154]
<i>Apobryophyllum terricola</i> +++ (this paper)	<i>Bursaria truncatella</i> +++ [84]
<i>Apocolpoda africana</i> +++ [84]	<i>Chaena clavata</i> + [132]
<i>Archinassula muscicola</i> + [157]	<i>Chaenea humicola</i> + [130]
<i>Arcuospathidium atypicum</i> +++ [78, 240]	<i>Chilodonella aplanata</i> + [154]
<i>Arcuospathidium cooperi</i> +++ [94]	<i>Chilodonella nigra</i> + [164]
<i>Arcuospathidium cultriforme</i> +++ [69, 152, 179]	<i>Chilodonella uncinata</i> +++ [114, 154]
<i>Arcuospathidium japonicum</i> +++ [78]	<i>Chilodontopsis muscorum</i> +++ [69, 154]
<i>Arcuospathidium litorntiforme</i> +++ [69, 152, 179]	<i>Chilophryxa terricola</i> +++ [69]
<i>Arcuospathidium muscorum</i> +++ [13, 41, 69]	<i>Chlamydonella alpestris</i> +++ [114]
<i>Arcuospathidium vermiforme</i> +++ [69]	<i>Cinetochilum margaritaceum</i> +++ [116]
<i>Aspidisca cicada</i> +++ [114]	<i>Cinetochilum marinum</i> ++ [185]
<i>Aspidisca lynceus</i> +++ [114]	<i>Circinella arenicola</i> +++ [88]
<i>Australocirrus octonucleatus</i> +++ [78]	<i>Circinella filiformis</i> +++ [65, 88]
<i>Australocirrus oscitans</i> +++ [20]	<i>Circinella vettversi</i> +++ [11, 88]
<i>Australothrix alwinae</i> +++ [20]	<i>Cirrophryxa australis</i> ++ [84]
<i>Australothrix australis</i> +++ [20]	<i>Cirrophryxa haptica</i> ++ [84]
<i>Australothrix simplex</i> ++ [252]	<i>Cirrophryxa terricola</i> +++ [77, 84]
<i>Australothrix steineri</i> +++ [93]	<i>Cladotricha australis</i> +++ [20]
<i>Avestina acuta</i> ++ [26, 84]	<i>Colpoda aspera</i> +++ [4, 84, 148, 154]
<i>Avestina ludwigi</i> +++ [84]	<i>Colpoda atra</i> + [4]
<i>Bakuella edaphoni</i> +++ [23, 212]	<i>Colpoda augustini</i> +++ [75, 84]
<i>Bakuella pampinaria</i> +++ [23, 47]	<i>Colpoda californica</i> + [84, 154]
<i>Balantidioides bivacuolata</i> +++ [111, 119, 127, 155, 230, 240]	<i>Colpoda cavicola</i> +++ [84, 157]
<i>Balantidioides corbifera</i> +++ [111, 119]	<i>Colpoda colpidiopsis</i> ++ [4, 84, 154]
<i>Balantidioides dragescoi</i> +++ [111]	<i>Colpoda cucullus</i> +++ [4, 84, 154]
<i>Balantidioides muscicola</i> + [152, 179]	<i>Colpoda ecaudata</i> +++ [84, 114]
<i>Bardeliella pulchra</i> +++ [69, 84]	<i>Colpoda edaphoni</i> +++ [84]
<i>Bicoronella costaricana</i> +++ [93]	<i>Colpoda ellioti</i> +++ [4, 84]
<i>Birojimia muscorum</i> +++ [11, 65, 155]	<i>Colpoda flavicans</i> +++ [84]
<i>Birojimia terricola</i> +++ [11]	<i>Colpoda henneguyi</i> +++ [84, 154]
<i>Blepharisma americanum</i> +++ [107, 139, 226]	<i>Colpoda inflata</i> +++ [4, 84, 154, 206]
<i>Blepharisma biancae</i> + [163]	<i>Colpoda irregularis</i> ++ [84, 154]
<i>Blepharisma bimicronucleatum</i> +++ [81, 139, 234]	<i>Colpoda lucida</i> +++ [4, 84, 135]
<i>Blepharisma hyalinum</i> +++ [27, 81, 139, 180]	<i>Colpoda magna</i> +++ [84, 137]
<i>Blepharisma lateritium</i> +++ [81, 115, 124, 139, 155, 160]	<i>Colpoda maupasi</i> +++ [84, 154]
<i>Blepharisma ovatum</i> + [139, 155, 179, 219]	<i>Colpoda minima</i> +++ [84]
<i>Blepharisma steini</i> +++ [4, 20, 81, 139, 155]	<i>Colpoda orientalis</i> +++ [84]
<i>Blepharisma undulans</i> +++ [81, 139, 155, 216]	<i>Colpoda praestans</i> ++ [84, 179]
<i>Brachonella cydonia</i> ++ [144, 150, 155]	<i>Colpoda reniformis</i> + [84, 154]
<i>Bresslaua insidiatrix</i> +++ [84]	<i>Colpoda simulans</i> + [84, 154]

Table 3. continued.

Species	Species
<i>Colpoda steinii</i> +++ [84, 154]	<i>Drepanomonas revoluta</i> +++ [27, 77, 129, 143, 154, 163, 240]
<i>Colpoda tripartita</i> +++ [20, 84, 154, 230]	<i>Drepanomonas sphagni</i> +++ [77, 154]
<i>Colpoda variabilis</i> +++ [84]	<i>Enchelydium piliforme</i> +++ [69, 102, 152, 153]
<i>Colpodidium caudatum</i> +++ [93, 243]	<i>Enchelydium polynucleatum</i> +++ [69, 103]
<i>Colpodidium viridis</i> +++ [147, 170]	<i>Enchelydium terrenum</i> +++ [69]
<i>Condylostoma terricola</i> +++ [93]	<i>Enchelyodon californicus</i> + [157]
<i>Corallocolpoda grelli</i> +++ [84]	<i>Enchelyodon lagenula</i> +++ [20, 152, 153]
<i>Corallocolpoda pacifica</i> +++ [84]	<i>Enchelyodon longinucleatus</i> +++ [69]
<i>Coriplites terricola</i> +++ [78]	<i>Enchelyodon nodosus</i> +++ [14]
<i>Corticocolpoda kaneshiroae</i> +++ [86]	<i>Enchelyodon terrenus</i> +++ [69]
<i>Cosmocolpoda naschbergeri</i> +++ [84, 105]	<i>Enchelyodon tratzi</i> +++ [74]
<i>Cothurnia minutissima</i> ++ [157, 177, 239]	<i>Enchelyomorpha vermicularis</i> +++ [6, 106, 132, 152, 153, 200]
<i>Cothurnia richtersi</i> ++ [157, 177, 179, 239]	<i>Enchelyotricha binucleata</i> +++ [77]
<i>Cothurniopsis valvata</i> ++ [157, 224, 239]	<i>Enchelys multimicronucleata</i> ++ [4]
<i>Cyclidium glaucoma</i> +++ [95, 116, 154]	<i>Enchelys multinucleata</i> +++ [14, 41]
<i>Cyclidium muscicola</i> +++ [93, 154]	<i>Enchelys terricola</i> +++ [77]
<i>Cyclidium terricola</i> + [154]	<i>Enchelys tokkuri</i> + [193]
<i>Cyrtobymena australis</i> +++ [93]	<i>Enchelys vermiformis</i> +++ [77]
<i>Cyrtobymena balladynula</i> ++ [81, 155]	<i>Engelmanniella mobilis</i> +++ [51, 65, 246, 247]
<i>Cyrtobymena candens</i> +++ [65, 81, 155]	<i>Epispadidium amphoriforme</i> +++ [69, 128, 135, 152, 179]
<i>Cyrtobymena candens depressa</i> ++ [81, 126]	<i>Epispadidium ascendens</i> +++ [77, 241]
<i>Cyrtobymena citrina</i> +++ [8, 81]	<i>Epispadidium papilliferum</i> +++ [69, 152, 153]
<i>Cyrtobymena gracilis</i> + [81, 155]	<i>Epispadidium regium</i> +++ [69]
<i>Cyrtobymena granulata</i> + [81, 155]	<i>Epispadidium terricola</i> +++ [77]
<i>Cyrtobymena muscorum</i> +++ [11, 26, 65, 81, 129, 155, 254]	<i>Epistylis alpestris</i> +++ [57, 59]
<i>Cyrtobymena primicirrata</i> +++ [8, 69]	<i>Erniella filiformis</i> +++ [75]
<i>Cyrtobymena quadrinucleata</i> +++ [43, 69, 81]	<i>Eschanestyla brachytoma</i> +++ [44, 65, 220]
<i>Cyrtobymena tetracirrata</i> +++ [8, 81, 126]	<i>Euplotes corsica</i> +++ [11]
<i>Cyrtolophosis acuta</i> +++ [84, 148]	<i>Euplotes finki</i> +++ [65, 257]
<i>Cyrtolophosis colpidiformis</i> +++ [84]	<i>Euplotes labiatus</i> +++ [20, 188]
<i>Cyrtolophosis elongata</i> +++ [27, 84, 154, 230]	<i>Euplotes muscicola</i> +++ [40, 65, 155, 204]
<i>Cyrtolophosis minor</i> +++ [84]	<i>Euplotes terricola</i> + [179]
<i>Cyrtolophosis mucicola</i> +++ [84, 154]	<i>Frontonia depressa</i> +++ [77, 126, 154, 179, 221]
<i>Dapedophrya flexilis</i> +++ [84, 93, 179]	<i>Frontonia parameciiformis</i> + [240]
<i>Deviata bacilliformis</i> +++ [8, 45, 125]	<i>Frontonia solea</i> ++ [77]
<i>Didinium nasutum</i> +++ [69, 117, 152]	<i>Frontonia terricola</i> +++ [77]
<i>Dileptus alpinus</i> +++ [39, 81, 154, 240]	<i>Furgasonia trichocystis</i> +++ [81, 144, 154, 225]
<i>Dileptus americanus</i> ++ [39, 154]	<i>Fuscheria lacustris</i> +++ [210]
<i>Dileptus anguillula</i> +++ [39, 64, 69, 154]	<i>Fuscheria nodosa</i> +++ [68, 107]
<i>Dileptus binucleatus</i> ++ [39, 154]	<i>Fuscheria terricola</i> +++ [4, 13, 104]
<i>Dileptus conspicuus</i> +++ [39, 81, 154]	<i>Gastronauta derouxi</i> +++ [21]
<i>Dileptus conspicuus telobivacuolatus</i> ++ [39, 128]	<i>Gastronauta membranaceus</i> +++ [4, 21, 76, 114]
<i>Dileptus costaricanus</i> +++ [93]	<i>Gastrostyla dorsicirrata</i> +++ [65]
<i>Dileptus edaphoni</i> +++ [205]	<i>Gastrostyla minima</i> +++ [98, 138]
<i>Dileptus falciformis</i> ++ [39, 154]	<i>Gastrostyla muscorum</i> ++ [155]
<i>Dileptus gracilis</i> +++ [39, 81, 154]	<i>Gastrostyla steinii</i> +++ [51, 65, 155, 194]
<i>Dileptus kahli</i> +++ [98, 205, 214]	<i>Glaucoma gigantea</i> + [132, 133]
<i>Dileptus margaritifer</i> +++ [39, 117]	<i>Gonostomum affine</i> +++ [26, 28, 65, 126, 129, 130, 165, 192, 213, 215, 217, 240]
<i>Dileptus mucronatus</i> +++ [39, 69, 179, 206]	<i>Gonostomum kuehnelti</i> +++ [77]
<i>Dileptus orientalis</i> +++ [211]	<i>Gonostomum strenua</i> +++ [51, 165, 202, 217]
<i>Dileptus polyvacuolatus</i> +++ [81]	<i>Grandoria aculeata</i> + [33, 84, 132]
<i>Dileptus similis</i> +++ [93]	<i>Grossglockneria acuta</i> +++ [84]
<i>Dileptus tenuis</i> ++ [39, 154, 179]	<i>Grossglockneria hyalina</i> +++ [84]
<i>Dileptus terrenus</i> +++ [64, 69]	<i>Halteria grandinella</i> +++ [114, 155, 227]
<i>Dileptus viesscheri</i> ++ [39]	<i>Halterioforma caudata</i> + [143]
<i>Dimacrocyton amphileptoides</i> +++ [39, 69, 145, 154, 240]	<i>Haplocaulus terrenus</i> +++ [63]
<i>Drepanomonas dentata</i> ++ [118, 154, 179, 186]	<i>Hastatella radians</i> +++ [57, 115, 157]
<i>Drepanomonas exigua</i> ++ [154, 179]	<i>Hausmanniella discoidea</i> +++ [69, 84, 129]
<i>Drepanomonas muscicola</i> +++ [77, 250]	<i>Hausmanniella patella</i> +++ [69, 84, 154]
<i>Drepanomonas pauciciliata</i> +++ [77]	

Table 3. continued.

Species	Species
<i>Hausmanniella quinquecirrata</i> + [84, 128]	<i>Kuklikophrya ougandae</i> +++ [42, 84]
<i>Hemiamphisiella granulifera</i> +++ [75, 78]	<i>Lacrymaria pulchra</i> ++ [240]
<i>Hemiamphisiella quadrinucleata</i> +++ [69, 78]	<i>Lagynophrya armata</i> + [157]
<i>Hemiamphisiella terricola</i> +++ [20, 44, 49, 69, 75, 209]	<i>Lagynophrya geleii</i> +++ [64]
<i>Hemiamphisiella wilberti</i> +++ [65, 78]	<i>Lagynophrya trichocystis</i> +++ [64]
<i>Hemisincirra gellerti</i> +++ [65, 69]	<i>Lamostyela abdita</i> +++ [97]
<i>Hemisincirra gracilis</i> +++ [65, 69]	<i>Lamostyela australis</i> +++ [20, 78, 183, 255]
<i>Hemisincirra heterocirrata</i> ++ [138]	<i>Lamostyela edaphoni</i> +++ [8, 9, 183]
<i>Hemisincirra inquieta</i> +++ [8, 11, 138]	<i>Lamostyela granulifera</i> +++ [97]
<i>Hemisincirra interrupta</i> +++ [65, 69]	<i>Lamostyela hyalina</i> +++ [9, 14]
<i>Hemisincirra kahli</i> + [26, 138]	<i>Lamostyela islandica</i> +++ [9]
<i>Hemisincirra muelleri</i> +++ [71]	<i>Lamostyela kirkeniensis</i> +++ [9]
<i>Hemisincirra octonucleata</i> ++ [138]	<i>Lamostyela lamottei</i> ++ [9, 27]
<i>Hemisincirra polynucleata</i> +++ [69]	<i>Lamostyela longa</i> ++ [9, 138]
<i>Hemisincirra pori</i> ++ [73, 244]	<i>Lamostyela perisincirra</i> +++ [9, 14, 138]
<i>Hemisincirra quadrinucleata</i> ++ [138]	<i>Lamostyela raptans</i> ++ [97, 138]
<i>Hemisincirra similis</i> +++ [65, 69]	<i>Laurentiella strenua</i> +++ [11, 38]
<i>Hemisincirra vermiculare</i> ++ [138]	<i>Leptopharynx costatus</i> +++ [4, 81, 116, 126, 154, 169, 175, 240]
<i>Hemisincirra wenzeli</i> +++ [75]	<i>Leptopharynx eurystoma</i> +++ [154, 175]
<i>Holophrya bimacronucleata</i> + [132, 133]	<i>Linostoma vorticella</i> +++ [115, 146]
<i>Holosticha adami</i> +++ [65]	<i>Litonotus crinitus</i> + [132]
<i>Holosticha australis</i> +++ [20]	<i>Litonotus digitatus</i> + [132]
<i>Holosticha bergeri</i> +++ [75]	<i>Litonotus muscorum</i> +++ [20, 154]
<i>Holosticha distyla</i> ++ [27]	<i>Maryna acuminata</i> ++ [84, 128]
<i>Holosticha islandica</i> +++ [11]	<i>Maryna antarctica</i> +++ [84]
<i>Holosticha longiseta</i> + [164]	<i>Maryna atra</i> +++ [84]
<i>Holosticha manca mononucleata</i> + [23, 128]	<i>Maryna lichenicola</i> +++ [84]
<i>Holosticha manca plurinucleata</i> ++ [23, 128]	<i>Maryna minima</i> ++ [84]
<i>Holosticha monilata</i> +++ [23, 102, 114, 151, 155]	<i>Maryna ovata</i> +++ [84]
<i>Holosticha multistilata</i> +++ [23, 27, 65, 114, 151, 155, 195]	<i>Maryna pinguis</i> + [38, 84]
<i>Holosticha muscicola</i> ++ [23, 129]	<i>Maryna rotunda</i> ++ [38, 84]
<i>Holosticha muscorum</i> +++ [23, 65, 155]	<i>Maryna socialis</i> + [84, 137]
<i>Holosticha sigmaoidea</i> +++ [23, 65, 69]	<i>Maryna umbrellata</i> ++ [84]
<i>Holosticha stueberi</i> +++ [74]	<i>Metacineta micraster</i> ++ [7, 166, 178]
<i>Holosticha sylvatica</i> +++ [11, 23, 65, 195]	<i>Metacineta mystacina</i> +++ [117, 166]
<i>Holosticha tetricirrata</i> +++ [20, 23, 28, 65]	<i>Metopus contractus</i> + [155, 179]
<i>Holostichides chardezi</i> +++ [75]	<i>Metopus es</i> +++ [115, 144, 155]
<i>Holostichides terricola</i> +++ [78]	<i>Metopus basei</i> +++ [63, 155, 201]
<i>Holostichides typicus</i> +++ [44, 208]	<i>Metopus ovalis</i> +++ [150, 155, this paper]
<i>Holostichides wilberti</i> +++ [44, 203]	<i>Metopus palaformis</i> ++ [144, 150, 155]
<i>Homalogastra setosa</i> +++ [27, 112, 132, 154, 185]	<i>Metopus rectus</i> ++ [61, 155]
<i>Idiocolpoda pelobia</i> +++ [85]	<i>Metopus setosus</i> ++ [144, 150, 155]
<i>Ilsiella palustris</i> +++ [84]	<i>Metopus setosus minor</i> ++ [61, 150, 155]
<i>Ilsiella venusta</i> +++ [75, 84]	<i>Microdiaphanosoma arcuatum</i> +++ [63, 84, 132, 240]
<i>Jaroschia sumptuosa</i> +++ [84]	<i>Microdiaphanosoma terricola</i> +++ [84]
<i>Kahlia acrobates</i> ++ [33, 140]	<i>Microthorax simulans</i> +++ [70, 148, 154]
<i>Kahlia simplex</i> +++ [8, 33, 142]	<i>Monodinium balbiani</i> +++ [117]
<i>Kablilembus attenuatus</i> +++ [112, 116, 136, 154]	<i>Mycterothrix tuamotuensis</i> ++ [84, 154]
<i>Kalometopia duplicata</i> +++ [84, 179]	<i>Mykophagophrys terricola</i> +++ [84, 93]
<i>Keronella gracilis</i> +++ [242]	<i>Nassula citrea</i> +++ [50, 154]
<i>Keronopsis algivora</i> ++ [126]	<i>Nassula picta</i> +++ [81, 116, 179, 230]
<i>Keronopsis dieckmanni</i> +++ (this paper)	<i>Nassula protectissima</i> + [154, 179]
<i>Keronopsis helluo</i> ++ [155, 179]	<i>Nassula terricola</i> +++ [81]
<i>Keronopsis tasmaniensis</i> +++ [20]	<i>Nassula tumida obscura</i> + [164]
<i>Keronopsis wetzeli</i> +++ [8, 240]	<i>Neogeneia hortualis</i> +++ [45]
<i>Keronopsis alpestris</i> + [155]	<i>Nivaliella plana</i> +++ [84]
<i>Krassniggia auxiliaris</i> +++ [75, 84]	<i>Notobrymena antarctica</i> +++ [95]
<i>Kreyella muscicola</i> + [84, 154]	<i>Notobrymena australis</i> +++ [20, 107]
<i>Kuehneltiella muscicola</i> +++ [84]	<i>Notobrymena rubescens</i> +++ [20, 253]
<i>Kuehneltiella terricola</i> +++ [84]	<i>Notobrymena selvatica</i> ++ [20, 138]

Table 3. continued.

Species	Species
<i>Notoxoma parabryophryides</i> +++ [84]	<i>Pelagotrichidium tisiae</i> +++ [42, 114, 146]
<i>Notoxoma sigmoides</i> +++ [84]	<i>Pentahymena corticicola</i> +++ [90]
<i>Obertrumia kahli</i> +++ [81]	<i>Periholosticha acuminata</i> ++ [138]
<i>Odontochlamys alpestris</i> +++ [21, 64, 184]	<i>Periholosticha lanceolata</i> ++ [138]
<i>Odontochlamys convexa</i> +++ [21, 154, 230]	<i>Phacodinium metchnicoffi</i> +++ [29, 40, 42, 54, 155, 196]
<i>Odontochlamys gouraudi</i> +++ [27, 29, 80, 154, 179, 240]	<i>Phialina binucleata</i> +++ [14]
<i>Odontochlamys wisconsinensis</i> +++ [154, 184]	<i>Phialina flagellifera</i> + [130]
<i>Onychodromopsis flexilis</i> +++ [183, 222]	<i>Phialina terricola</i> +++ [28, 69]
<i>Opercularia arenicola</i> + [134]	<i>Philanides australis</i> +++ [78]
<i>Opercularia asymmetrica</i> +++ [1, 18]	<i>Plagiocampa atra</i> + [132, 133]
<i>Opercularia curvicauda</i> +++ [19, 63, 157, 179]	<i>Plagiocampa caudata</i> + [4]
<i>Ophryoglena marginata</i> + [135]	<i>Plagiocampa difficilis</i> +++ [64]
<i>Opisthonecta minima</i> +++ [56]	<i>Plagiocampa rouxi</i> +++ [58, 148, 152]
<i>Opisthotricha elongata</i> + [132]	<i>Platycola longicollis</i> ++ [157, 158, 238]
<i>Opisthotricha procera</i> + [155]	<i>Platycola steineri</i> ++ [157, 177, 238]
<i>Opisthotricha terricola</i> + [130]	<i>Platyophrya binucleata</i> +++ [75, 84]
<i>Orthoamphisiella franzi</i> +++ [10, 45, 65]	<i>Platyophrya macrostoma</i> +++ [84]
<i>Orthoamphisiella grelli</i> +++ [48]	<i>Platyophrya paoletti</i> +++ [97]
<i>Orthoamphisiella stramenticola</i> +++ [46, 48]	<i>Platyophrya similis</i> +++ [84]
<i>Orthokreyella schiffmanni</i> +++ [69, 84]	<i>Platyophrya spumacola</i> +++ [27, 84, 143, 149, 152, 163]
<i>Oxytricha auripunctata</i> +++ [20]	<i>Platyophrya vorax</i> +++ [84, 148, 152]
<i>Oxytricha bimembranata</i> + [192]	<i>Platyophryides dragescoi</i> +++ [75, 84]
<i>Oxytricha gigantea</i> +++ [8, 10, 141]	<i>Platyophryides latus</i> +++ [41, 84, 128, 152]
<i>Oxytricha granulifera</i> +++ [101]	<i>Platyophryides magnus</i> +++ [84]
<i>Oxytricha granulifera quadricirrata</i> +++ [20]	<i>Pleuroplites australis</i> +++ [78]
<i>Oxytricha hengshanensis</i> ++ [252]	<i>Pleuroplitoïdes smithi</i> +++ [95]
<i>Oxytricha histrioides</i> + [130]	<i>Podophrya halophila</i> +++ [20, 156]
<i>Oxytricha islandica</i> +++ [11]	<i>Protocyclidium terrenum</i> ++ [4]
<i>Oxytricha lanceolata</i> +++ [8, 11, 95, 193]	<i>Protopathidium bonneti</i> +++ [27, 42, 64]
<i>Oxytricha longigranulosa</i> +++ [11]	<i>Protopathidium muscicola</i> +++ [14, 41, 184]
<i>Oxytricha nauplia</i> +++ [8]	<i>Protopathidium serpens</i> +++ [64, 95, 152, 184]
<i>Oxytricha opisthomuscorum</i> +++ [114, 155, 184]	<i>Protopathidium terricola</i> +++ (this paper and [184])
<i>Oxytricha ottowi</i> +++ [94]	<i>Pseudocarchesium claudicans</i> +++ [81, 179]
<i>Oxytricha proximata</i> + [193]	<i>Pseudochilonopsis mutabilis</i> +++ [64, 79, 126, 184]
<i>Oxytricha pseudosimilis</i> ++ [138]	<i>Pseudochilonopsis polyvacuolata</i> +++ [98, 102]
<i>Oxytricha quercineta</i> + [163]	<i>Pseudocohnilembus marinus</i> +++ [110, 228]
<i>Oxytricha rubripuncta</i> +++ [8]	<i>Pseudocohnilembus pusillus</i> +++ [5, 53, 110, 116, 185, 187]
<i>Oxytricha setigera</i> +++ [27, 42, 65, 114, 210]	<i>Pseudocohnilembus putrinus</i> +++ [110, 151, 154]
<i>Oxytricha siseris</i> +++ [65, 237]	<i>Pseudocyrtocephosis alpestris</i> +++ [84]
<i>Oxytricha tricirrata</i> ++ [27]	<i>Pseudocyrtocephosis terricola</i> +++ [84]
<i>Papillorhabdos multinucleatus</i> +++ [69]	<i>Pseudoglaucoma muscorum</i> + [84, 154, 240]
<i>Parabryophrya penardi</i> +++ [84, 154]	<i>Pseudoholophrya terricola</i> +++ [14]
<i>Paracineta lauterborni</i> +++ [93, 201]	<i>Pseudokreyella terricola</i> +++ [84]
<i>Paraenchelys terricola</i> +++ [69]	<i>Pseudokreyella australis</i> +++ [84]
<i>Paraenchelys wenzeli</i> +++ [69]	<i>Pseudomicrothorax agilis</i> +++ [32, 116, 169, 176, 213, 240]
<i>Parafurgasonia sorex</i> +++ [42, 99, 179]	<i>Pseudomicrothorax dubius</i> +++ [32, 116, 167, 176, 179]
<i>Paragastrostyla lanceolata</i> ++ [138]	<i>Pseudoplatyophrya nana</i> +++ [84, 148]
<i>Paraholosticha lichenicola</i> ++ [128]	<i>Pseudoplatyophrya saltans</i> +++ [78, 84]
<i>Paraholosticha muscicola</i> +++ [36, 37, 75, 155]	<i>Pseudothuricola dionysii</i> ++ [157, 179, 231]
<i>Paraholosticha nana</i> + [128]	<i>Pseudouroleptus buithkampi</i> +++ [8, 65]
<i>Parakahliella haideri</i> +++ [12]	<i>Pseudouroleptus procerus</i> +++ [8]
<i>Parakahliella macrostoma</i> +++ [15, 65]	<i>Pseudouroleptus terrestris</i> ++ [138]
<i>Parakahliella terricola</i> ++ [15, 26]	<i>Pseudostylophora franzii</i> +++ [75]
<i>Paramphisiella acuta</i> +++ [65, 78]	<i>Pseudovorticella mutans</i> +++ [60, 179]
<i>Paramphisiella cadata</i> +++ [49, 78, 138]	<i>Pseudovorticella sphagni</i> +++ [60, 93, 109]
<i>Paraurostylo granulifera</i> +++ [11]	<i>Psilotricha succisa</i> +++ [67, 173]
<i>Paraurostylo polynucleata</i> ++ [4]	<i>Pyxidium longicollum</i> + [19]
<i>Paraurostylo pulchra</i> ++ [26]	<i>Pyxidium tardigradum</i> ++ [233]
<i>Pattersoniella vitiphila</i> +++ [75]	<i>Reticulowoodruffia terricola</i> +++ [84]
<i>Pedohymena australiense</i> +++ [93]	<i>Rhabdostyla arborea</i> + [135]

Table 3. continued.

Species	Species
<i>Rhabdostyla muscorum</i> + [157]	<i>Telostomatella ferroii</i> + [84, 133]
<i>Rhabdotricha terricola</i> + [135]	<i>Telotrochidium cylindricum</i> +++ [14, 59]
<i>Rhopalophrya elegans</i> + [143]	<i>Terricirra livida</i> +++ [9, 11]
<i>Rhopalophrya pentacerca</i> + [132]	<i>Terricirra matsusakai</i> +++ [11]
<i>Rostrophrya camerounensis</i> +++ [42, 84, 98]	<i>Terricirra viridis</i> +++ [11, 65]
<i>Rostrophrya terricola</i> +++ [84]	<i>Territricha stramenticola</i> +++ [9]
<i>Rostrophryides africana</i> +++ [75, 84]	<i>Tetrahymena edaphoni</i> +++ [77]
<i>Rostrophryides australis</i> +++ [20, 84]	<i>Tetrahymena rostrata</i> +++ [4, 31, 77, 130, 148, 154]
<i>Sagittaria australis</i> +++ [84, 185]	<i>Thuricola innixa</i> ++ [157, 177, 218, 223, 231, 239]
<i>Sagittaria hyalina</i> +++ [84]	<i>Thuricola kellicottiana</i> ++ [115, 157, 223, 231]
<i>Sagittaria polygonalis</i> + [84, 132, 133]	<i>Thylakidium typicum</i> + [84, 128]
<i>Sathrophilus muscorum</i> +++ [27, 33, 112, 116, 154]	<i>Tokophrya muscicola</i> ++ [166, 178]
<i>Sathrophilus simonis</i> + [33, 163]	<i>Trachelochaeta gonostomoida</i> ++ [138]
<i>Semiplatyphrya foissneri</i> +++ [84, 244]	<i>Trachelophyllum apiculatum</i> +++ [68, 69, 117, 181, 206]
<i>Solenophrya flavescentia</i> ++ [166, 178]	<i>Tricoronella pulchra</i> +++ [20]
<i>Solenophrya massula</i> ++ [166, 178]	<i>Tribymena terricola</i> +++ [78, 84]
<i>Solenophrya sacculus</i> ++ [166, 178]	<i>Trithigmostoma bavariensis</i> +++ [73, 76, 79, 154, 230]
<i>Sorogena stoianovitchae</i> +++ [84]	<i>Urliella terricola</i> +++ [81]
<i>Spathidium alpinum</i> ++ [129]	<i>Uroleptoides kibni</i> + [22, 240]
<i>Spathidium anguilla</i> +++ [6, 69, 185, 235]	<i>Uroleptus humicola</i> ++ [129]
<i>Spathidium bavariense</i> ++ [152, 153, 230, 240]	<i>Uroleptus lepisma</i> +++ [11, 240]
<i>Spathidium claviforme</i> +++ [77, 152, 153]	<i>Uroleptus matthesi</i> + [240]
<i>Spathidium falciforme</i> + [152, 179]	<i>Uroleptus musculus</i> +++ [69, 114, 155]
<i>Spathidium furcatum</i> + [193]	<i>Uroleptus notabilis</i> +++ [8, 20, 65, 95]
<i>Spathidium geobium</i> + [164]	<i>Uronema nigricans</i> +++ [4, 116, 173]
<i>Spathidium holsatiae</i> ++ [152, 153]	<i>Urosoma acuminata</i> +++ [65, 155, 222]
<i>Spathidium lagyniforme</i> +++ [69, 152]	<i>Urosoma cienkowskii</i> +++ [65, 69, 155, 159]
<i>Spathidium longicaudatum</i> +++ [26, 28, 64]	<i>Urosoma karini</i> +++ [74]
<i>Spathidium multinucleatum</i> + [128]	<i>Urosoma macrostoma</i> ++ [130]
<i>Spathidium muscicola</i> ++ [4, 14, 27, 152, 153, 185, 206]	<i>Urosoma macrostyla</i> +++ [65, 66, 155, 248, 249]
<i>Spathidium procerum</i> +++ [4, 69, 152, 153]	<i>Urosoma octonucleata</i> +++ [11]
<i>Spathidium rusticum</i> ++ [64]	<i>Urosomoida agiliformis</i> +++ [65, 100, 120]
<i>Spathidium scalpriforme</i> ++ [152, 153]	<i>Urosomoida agilis</i> +++ [25, 51, 65, 138]
<i>Spathidium seppelti</i> +++ [184]	<i>Urosomoida antarctica</i> +++ [95]
<i>Spathidium spathula</i> +++ [69, 126, 152, 171, 172, 240]	<i>Urosomoida dorsiincisura</i> +++ [65]
<i>Spathidium spathuloides</i> + [128]	<i>Urosomoida granulifera</i> +++ [95]
<i>Spetazoon austriense</i> +++ [87]	<i>Urosomoida minima</i> ++ [138]
<i>Sphaerophrya parva</i> + [135, 166]	<i>Urostyla grandis</i> +++ [114, 213]
<i>Sphaerophrya terricola</i> ++ [71, 166]	<i>Urostyla muscorum</i> ++ [155]
<i>Spirofilopsis tubicola</i> + [38, 137]	<i>Urotricha atypica</i> ++ [4]
<i>Stammeridium kahli</i> +++ [70, 240]	<i>Urotricha mamilla</i> + [164]
<i>Steinia platystoma</i> ++ [27, 114]	<i>Urotricha terricola</i> ++ [5]
<i>Sterkiella cavicola</i> +++ [8, 27, 114, 157]	<i>Vaginicola chaperoni</i> ++ [157, 177, 179, 239]
<i>Sterkiella histriomuscorum</i> +++ [6, 15, 65, 114, 143, 155, 184]	<i>Vaginicola doliolum</i> ++ [157, 177, 179, 239]
<i>Sterkiella similis</i> f. <i>tricirrata</i> ++ [27]	<i>Vaginicola terricola</i> ++ [135, 177, 179]
<i>Sterkiella thompsoni</i> +++ [95]	<i>Vaginicola virgula</i> ++ [157, 177, 179, 239]
<i>Stichotricha aculeata</i> +++ [61, 114, 248]	<i>Vorticella astyliformis</i> +++ [63, 115]
<i>Stichotricha socialis</i> ++ [137]	<i>Vorticella coeni</i> + [163]
<i>Strongylidium californicum</i> + [155]	<i>Vorticella infusionum</i> +++ [115]
<i>Strongylidium muscorum</i> + [155]	<i>Vorticella lichenicola</i> ++ [135, 179]
<i>Styloynchia bifaria</i> +++ [222, 245]	<i>Vorticella microstoma</i> +++ [115]
<i>Styloynchia mytilus</i> +++ [65, 114, 172]	<i>Vorticella muralis</i> ++ [179]
<i>Styloynchia pustulata</i> +++ [114, 173]	<i>Vorticella operculariformis</i> +++ [60]
<i>Styloynchia quadrinucleata</i> ++ [5]	<i>Vorticella pileolata</i> + [163]
<i>Styloynchia stylomuscorum</i> ++ [114, 155]	<i>Vorticella similis</i> +++ [63, 115]
<i>Tachysoma granulifera</i> +++ [8]	<i>Wallackia bujoreani</i> +++ [11, 164]
<i>Tachysoma humicola</i> +++ [69, 130]	<i>Woodruffia australis</i> +++ [84]
<i>Tachysoma humicola longisetum</i> +++ (this paper)	<i>Woodruffia rostrata</i> +++ [75, 84, 154]
<i>Tachysoma terricola</i> ++ [138]	<i>Woodruffides metabolicus</i> +++ [84]
<i>Tectohymena terricola</i> +++ [84]	<i>Woodruffides terricola</i> +++ [75, 84]

Table 4. Statistics and ecological characteristics of the soil ciliate community (extracted from Tables 2 and 3).

Characteristics	According to Tables 2, 3		According to Tables 2, 3 plus 500 undescribed species ¹⁾	
	number of species	%	number of species	%
Total number of species described	643	100.0	—	—
Valid species	594	92.4	1092	100.0
Synonyms	49	7.6		not determined
Excellent described species	409	68.9		not determined
Sufficiently described species	107	17.9		not determined
Poorly described species	78	13.2	none	
Colpodids	129	21.8	146	13.4
Cyrtophorids	14	2.2	16	1.5
Gymnostomatids	105	17.8	289	26.5
Heterotrichs	24	4.1	35	3.2
Hymenostomes	23	4.0	35	3.2
Hypotrichs	214	36.0	429	39.2
Nassulids	26	4.4	50	4.6
Oligotrichs	1	0.1	3	0.3
Peritrichs	35	5.9	40	3.7
Prostomatids	10	1.7	22	2.0
Suctorians	12	2.0	26	2.4
Small species (biomass \leq 100mg/10 ⁶ cells)	416	70.9		not determined
Large species (biomass \geq 400mg/10 ⁶ cells)	48	8.2		not determined
Mean Biomass of 10 ⁶ specimens	241	—		not determined
Omnivores ²⁾	103	20.2		not determined
Mainly bacteriovorous	196	38.5		not determined
Mainly predaceous	172	34.1		not determined
Mainly (filamentous) cyanobacteria	18	3.6		not determined
Mainly mycophagous	8	1.6		not determined
Aerobics	582	98.3	1074	98.4
Anaerobics	10	1.7	18	1.6
Occurring only in terrestrial habitats (***)	133	22.5	397	36.4
Probably occurring only in terrestrial habitats (**)	310	52.2	519	47.6
Occurring in soil and freshwater (*)	151	25.3	174	16.0
Recorded from 1 geographical region only	265	44.5		not determined
Recorded from 2 geographical regions only	95	16.0		not determined
Recorded from 3 geographical regions only	64	10.8		not determined
Recorded from 4 geographical regions	103	17.4		not determined
Recorded from all (5) geographical regions	67	11.3		not determined
Species recorded from Holarctis	483	81.2		not determined
Species recorded from Palaeotropis	290	49.0		not determined
Species recorded from Australis	251	42.4		not determined
Species recorded from Neotropis	259	43.9		not determined
Species recorded from Archinotis	101	17.1		not determined

¹⁾ I have not yet described these species, which I found in about 1000 soil samples worldwide, but fully studied in live and silver slides. Many belong to new genera or to genera as yet known from soil only.²⁾ Feeding on \geq than three items.

the reliable biogeographic information is still much too incomplete and most of the rare species possibly have not yet been discovered due to methodological problems [96]. In spite of this, the present data set is probably the most complete and reliable available, not only for soil ciliates but also for ciliates in general. Thus, some preliminary conclusions should be possible.

Only about 30% of the soil ciliate species have been found in more than three out of five biogeographical regions, and 44% have been recorded from one region only (Tab. 4). These figures suggest a high degree of endemism. However, they are very likely strongly biased due to the very limited data available. To mention only two examples: When *Tribymena terricola* was discovered in African soils in 1988, I supposed that it could be endemic because of the unique organisation of its oral apparatus. Later it was found in many soils world-wide [16, 84; Tab. 2], indicating that we previously overlooked or misidentified this minute creature. Likewise, the bipolar biogeography of the genus *Colpoda*, suggested by Smith [197], did not withstand a more detailed analysis and was caused by methodological shortcomings [95].

Thus, it is probably more accurate to select the data very rigorously, that is to look at conspicuous species ("flagships" with large size or other very distinct characters) which are easily recognised and identified. Such species are, for example, *Apobryophyllum terricola*, *Australothrix* spp., *Bresslauides australis*, *Cosmocolpoda naschbergeri*, *Erniella filiformis*, *Jaroschia sumptuosa*, *Keronopsis tasmaniensis*, *Krassniggia auxiliaris*, *Periholosticha* spp. and *Tricoronella pulchra*, which have so far been found only in the Palaeotropis and/or Australis, whereas, e.g., *Amphisiella quadrinucleata*, *Bresslauides discoideus*, *Chilophrya terricola*, *Hemiamphisiella quadrinucleata*, *Kalometopia duplicata*, *Keronopsis wetzeli*, *Orthoamphisiella franzi* and *Wallowcia bujoreani* have been found only in the Holarctic. These data, especially the limited distribution of the two most conspicuous species, viz. *Krassniggia auxiliaris* and *Bresslauides discoideus*, both belonging to the order Colpodida, suggest that at least some species have a limited Gondwanan or Laurasian distribution. However, even this example is not entirely convincing because both species are very rare and the limited distribution could thus be caused by undersampling.

Description of new and insufficiently known species

Protospadidium terricola nov. spec. (Figs. 2–13, Tab. 5)

Diagnosis: Size in vivo about 90×25 µm. Elongate saccular with anterior portion usually slightly narrowed

and curved. Macronucleus elongate reniform (3.5:1). Extrusomes rod-shaped, about 5 µm long. On average 21 somatic kineties with 40 basal bodies each, 3 of them anteriorly differentiated to distinct brush on dorsal side.

Type location: Grassland soil from Mt. Kenya near the lodge "The Ark", Mount Kenya National Park, Kenya, equatorial Africa (about 37° E, 0° N).

Etymology: *terricola* because living in soil.


Description: Size in vivo 70–100×20–30 µm. Elongate saccular, usually looking like a swimming sausage, anterior portion frequently slightly curved, middle and posterior portion more or less distinctly inflated, depending on nutritional state (Figs. 2, 5). Unflattened, except for slightly compressed anterior third. Macronucleus in mid-body, elongate reniform, rarely slenderly ellipsoidal, contains many tiny nucleoli. Micro-nucleus globular, attached to macronucleus, but not in fixed position. Contractile vacuole in posterior end. Extrusomes mainly in oral bulge, rod-shaped, about 5 µm long (Fig. 3). Cortex flexible, contains about four rows of minute, colourless granules between each two kineties (Fig. 6). Cytoplasm usually packed with fat globules 0.5–3 µm across. Swims moderately fast by rotation about longitudinal body axis.

Cilia 10–12 µm long, rather unevenly spaced. Somatic kineties evenly spaced, extend meridionally, anterior ends conspicuously curved and composed of 3–6 distinct dikinetids associated with long, fine nematodes-mata. Dorsal brush of usual structure, i.e. composed of narrowly spaced dikinetids having short, slightly inflated cilia; at anterior end of each brush kinety 2–6 monokinetids (Figs. 7–11).

Oral bulge conspicuous, although indistinctly set off from neck, because rather high and refractile due to the extrusomes contained. Bulge centre depressed funnel-like. Circumoral kinety discontinuous because anterior dikinetidal fragments adhering to their respective somatic kineties separated by small gaps (Figs. 2, 7, 9–11).

Generic classification and comparison with related species: *Protospadidium terricola* has the typical generic character of *Protospadidium* [41], viz. a discontinuous circumoral kinety composed of small dikinetidal fragments. These fragments adhere to the anterior end of the somatic kineties and are separated from each other by a narrow gap (Figs. 9–11), unlike in *Spathidium* (Figs. 12, 13) and *Arcuospathidium* [69], where the fragments are fused to a single, continuous row [69].

Protospadidium terricola is a difficult species because it has few distinct characters and is thus, in life, easily confused with several congeners and with *Spathidium claviforme*, which have a similar size, shape and macronucleus; in fact, it was only a fortunate chance that I did not confuse it with *S. claviforme* during the routine inspection of the sample, i.e. recognised the discontinuous circumoral kinety! *Spathidium clavi-*



Figs. 2–13. *Protospathidium terricola* (2, 3, 5–11), *Spathidium vermiculus* (4, from [148]), and *S. claviforme* (12, 13, from [77]) from life (2–6) and after protargol impregnation (7–13). 2. Right lateral view of typical specimen packed with fat globules. 3. Extrosome (5 µm) at high magnification. 4. *Spathidium vermiculus*, length 40–50 µm. 5. Saccular shape variant of *P. terricola*. 6. Surface view showing cortical granules. 7, 8, 12. Infraciliature of right (7, 12) and left (8) side of *P. terricola* (7, 8) and *S. claviforme* (12). 9. Oral kinetofragments at high magnification. The fragments do not form a continuous circumoral kinety, as they do in *S. claviforme* (13). 10, 11. Infraciliature of anterior ventral and dorsal side. 13. Infraciliature of anterior ventral side of *S. claviforme*, where the oral kinetofragments form a continuous circumoral kinety. B = dorsal brush, CF = (circum)oral kinetofragments, CK = circumoral kinety, FG = fat globules, MI = micronucleus, N = nematodesmata, OB = oral bulge. Scale bar division 10 µm.

Table 5. Morphometric data from *Protospathidium terricola* (upper line) and *Apobryophyllum terricola* (lower line).

Character ¹⁾	\bar{x}	M	SD	SE	CV	Min	Max	n
Body, length	80.7	82.0	9.6	3.0	11.9	65.0	93.0	10
	154.0	155.0	22.1	7.0	14.3	125.0	190.0	10
Body, maximum width	22.5	22.0	3.6	1.1	16.1	16.0	28.0	10
	21.2	20.5	5.9	1.9	28.0	16.0	35.0	10
Oral bulge, length	14.3	15.0	1.7	0.6	11.6	12.0	17.0	10
	—	—	—	—	—	—	—	—
Macronuclear figure, length	28.1	26.5	4.2	1.3	15.0	22.0	35.0	10
	60.4	61.0	12.8	4.0	21.1	40.0	80.0	10
Macronucleus, width	7.6	8.0	0.8	0.3	11.1	6.0	9.0	10
	3.9	4.0	1.2	0.4	30.7	2.0	6.0	10
Macronuclei, number	1.0	1.0	0.0	0.0	0.0	1.0	1.0	10
	1.0	1.0	0.0	0.0	0.0	1.0	1.0	10
Micronucleus, largest diameter	3.5	3.5	—	—	—	3.0	4.0	10
	2.3	2.1	—	—	—	2.0	3.0	10
Micronuclei, number	1.0	1.0	0.0	0.0	0.0	1.0	1.0	10
	8.8	8.0	2.8	1.2	31.5	6.0	13.0	5
Somatic kineties, number in mid-body	21.3	21.0	1.6	0.5	7.7	19.0	25.0	10
	16.0	16.0	2.3	0.7	14.1	13.0	19.0	10
Basal bodies in right lateral kinety, number	42.2	39.5	7.3	2.3	17.3	35.0	55.0	10
	63.3	65.0	—	—	—	55.0	70.0	3
Dorsal brush rows, number	3.0	3.0	0.0	0.0	0.0	3.0	3.0	10
	4.2	4.0	—	—	—	4.0	5.0	10
Circumoral kinety to end of brush row 1, distance	15.0	15.0	2.6	0.9	17.0	11.0	18.0	9
	19.1	20.0	1.9	0.6	10.1	15.0	22.0	10
Circumoral kinety to end of brush row 2, distance	18.0	18.0	2.1	0.7	11.5	15.0	21.0	9
	35.8	35.0	5.0	1.6	13.9	30.0	48.0	10
Circumoral kinety to end of brush row 3, distance	10.8	10.0	1.7	0.6	15.9	8.0	14.0	9
	40.3	40.0	5.6	1.8	14.0	35.0	55.0	10
Circumoral kinety to end of brush row 4, distance	—	—	—	—	—	—	—	—
	39.8	39.0	5.9	1.9	14.8	35.0	55.0	10

¹⁾ Data based on protargol-impregnated and mounted specimens from field. Measurements in μm . Abbreviations: CV = coefficient of variation in %, M = median, Max = maximum, Min = minimum, n = number of individuals investigated, SD = standard deviation, SE = standard deviation of mean, \bar{x} = arithmetic mean.

forme has fewer ciliary rows than *P. terricola* (10–13 vs. 19–25) and is a “true” *Spathidium* because the anterior kinetofragments are fused to a continuous circumoral kinety (Figs. 12, 13 [77]). This was confirmed by a reinvestigation of the neotype slides.

As concerns the congeners, *P. serpens* has fewer ciliary rows (11–13 vs. 19–25) and its macronucleus is longer and usually distinctly coiled and/or nodulated [64, 95, 184]. *Protospathidium muscicola* [41] has many small macronuclear nodules and only 10–12 ciliary rows.

Protospathidium terricola also resembles *Spathidium vermiculus* [148] which, however, is smaller (40–60 μm) and has a more cylindroid shape and only about 10 somatic kineties (Fig. 4 [148, 152, 251]).

Apobryophyllum nov. gen.


Diagnosis: Spathidiidae with oral bulge extending to and around posterior end of organism. Dorsal brush on

anterior left side of cell, left brush kineties regular and dikinetidal, right brush kineties fragmented and very likely monokinetidal.

Type species: *Apobryophyllum terricola* nov. spec.

Etymology: Composite of *apo* (derived), *bryon* (moss), and *phyllon* (leaf). Neuter gender.

Comparison with related genera: Gymnostome ciliates have a fascinating variety of brush structures, often separating species at genus level which otherwise look very much alike [69]. *Apobryophyllum* certainly has a new type of brush, and is closely related to *Bryophyllum*, as indicated by the typical oral bulge and lateral flattening. *Bryophyllum tegularum* has, like *Spathidium* spp., three brush kineties of usual structure [69]. However, the *Bryophyllum* species described by Gelei [123] have, like *Apobryophyllum*, a more complicated, bipartited brush composed of many short kineties near the organism’s anterior end and three to five long rows in mid-body. Unfortunately, the details provided by Gelei [123] are insufficient to decide



Figs. 14–24. *Apobryophyllum terricola* from life (14–18, 23, 24) and after protargol impregnation (19–22). 14, 15. Left lateral and ventral view of typical specimen. 16. Broad shape variant showing arrangement of extrusomes. 17. Extrusome in surface and lateral view. 18. Optical section of cell periphery showing arrangement of cortical granules (cp. Fig. 24) and extrusomes, which form a fringe around the cell (cp. Fig. 16). 19. Frontal view of oral bulge showing orientation and ciliation of circumoral dikinetids. 20, 21. Infraciliature of right and left side and nuclear apparatus. 22. Infraciliature of anterior left side showing the genus-specific brush, which consists of four rows, in detail. 23. Posterior end of brush rows (numbers 1–4), which have a highly differentiated ciliation. 24. Surface view showing cortical granules. B = dorsal brush, C = normal somatic cilium, E = extrusomes, EP = pores of contractile vacuole, FG = fat globule, LC = left branch of circumoral kinety, MA = macronucleus, MI = micronucleus, N = nematodesmata, OB = oral bulge, RC = right branch of circumoral kinety. Scale bar division 20 µm.

whether his species have fragmented brush rows like *Apobryophyllum*.

The monokinetal (?) fragments in the brush of *Apobryophyllum* are reminiscent of those found in *Paraenchelys* which, however, has a very small, apical oral opening and lacks dikinetidal brush rows [69].

Apobryophyllum has, unlike all *Bryophyllum* species known, a slightly bipartite oral bulge and circumoral kinety, composed of a slightly concave and relatively short, cuneate, densely ciliated anterior part and a long, straight, loosely ciliated posterior portion. The anterior part is highly reminiscent of the oral bulge of *Arcuospadidium*, indicating that *Apobryophyllum*, *Bryophyllum* and *Arcuospadidium* might have a common ancestor.

Apobryophyllum terricola nov. spec. (Figs. 14–24, Tab. 5)

Diagnosis: Size in vivo about 160×25 µm. Slenderly knife-shaped to spatulate. Macronucleus filiform and tortuous. Extrusomes rod-shaped, about 5 µm long, form peripheral fringe. 16 somatic kineties on average, 4 of them differentiated to distinct brush on anterior left side.

Type location: Soil from Shetani volcano area, Tsavo National Park, Kenya, equatorial Africa (about 38° E, 2°55' N).

Etymology: *terricola* because living in soil.

Description: Size in vivo 130–200×20–55 µm, length: width ratio 5:1 to 9:1, usually near 7:1. Slenderly spatulate to knife-shaped, i.e. anterior ventral third rather distinctly curved and gradually narrowing to form bluntly pointed dorsal anterior end; blade leaf-like flattened and sometimes slightly broadened, handle evenly rounded posteriorly and flattened up to 2:1 (Figs. 14–16). Macronucleus in middle body portion, filiform and tortuous, with many small nucleoli. Micronuclei globular, near or attached to macronucleus. Contractile vacuole in posterior end, about 6 excretory pores on right posterior surface of cell (Fig. 20). Extrusomes rod-shaped, slightly curved, about 5 µm long, more narrowly spaced in anterior third than posteriorly, attached to oral bulge in two indistinct rows and to dorsal side, forming conspicuous peripheral fringe (Figs. 14–18). Cortex flexible, fragile, contains about five rows of colourless granules between each two ciliary rows (Fig. 24). Cytoplasm usually packed with colourless fat globules 2–6 µm across and many scattered extrusomes very similar to those found in oral bulge. Glides slowly on slide surface and soil particles.

Cilia rather loosely spaced, arranged in meridional rows distinctly separate from circumoral kinety. Dorsal brush at anterior end of first four kineties of left side, about as long as blade of knife; rows 1 and 2 composed

of many short, rather irregular fragments having about 4 µm long, distinctly inflated cilia; rows 3 and 4 regularly dikinetidal with anterior cilium slightly inflated and longer (3 µm) than posterior (2 µm); row 4 has a monokinetal tail with short bristles extending to level of posterior end of row 3 (Figs. 21–23).


Oral bulge very inconspicuous, hardly distinct from body proper and thus almost invisible in live specimens, extends from anterior dorsal end to posterior dorsal end, bordered by dikinetidal circumoral kinety associated with very fine nematodesmata; circumoral dikinetids on right side very likely ciliated and orientated transversely to longitudinal axis of cell, those on left side, having very likely only the posterior basal body ciliated, longitudinally to slightly obliquely arranged (Figs. 14, 19–22).

Comparison with related species: No species has been found in the literature which might be identical with *A. terricola*. It is easily distinguished from the *Bryophyllum* species known by its very slender shape and inconspicuous oral bulge. However, live specimens of *A. terricola* are easily confused with large *Arcuospadidium* species, especially *A. lionotiforme*, which has a very similar size, shape and nuclear apparatus [69]. The best in vivo character for separating these species is the arrangement of the extrusomes, which are restricted to the oral bulge in *Arcuospadidium*, and distributed in a peripheral girdle in *A. terricola*. Some *Prorodon* and *Cranotheridium* species, especially *P. armatus* [98], also look very much alike *A. terricola*. However, all *Prorodon* and *Cranotheridium* species known live in freshwater.

Metopus ovalis Kahl, 1927 (Figs. 25–30)

Redescription: Like Kahl [150, 155], I found only few specimens. Thus, the morphometric analysis is rather incomplete and summarised here: length 102–145 µm ($\bar{x} = 123.8$, SD = 16.9, n = 6), postoral width 42–57 µm ($\bar{x} = 47.6$, SD = 6.0, n = 5), distance anterior end to macronucleus 24–45 µm ($\bar{x} = 33.3$, SD = 8.2, n = 6), distance anterior end to proximal end of adoral zone 53–80 µm ($\bar{x} = 64.2$, SD = 10.2, n = 6), length of macronucleus 20–34 µm ($\bar{x} = 25.7$, SD = 5.6, n = 6), width of macronucleus 13–22 µm ($\bar{x} = 16.0$, SD = 3.1, n = 6), largest diameter of micronucleus 3–5 µm ($\bar{x} = 4.0$, SD = 0.6, n = 6), number of perizonal kineties 5 (n = 5), number of somatic kineties in postoral body portion 23–29 ($\bar{x} = 25.8$, SD = 2.5, n = 4), number of adoral membranelles 46–55 ($\bar{x} = 50.6$, SD = 3.8, n = 4).

Size in vivo about 120–170×50–60 µm. Cells reddish at low (< X100) magnification, as also mentioned by Kahl [150], possibly due to minute cytoplasmic granules and food vacuole contents. Preoral dome conspicuous, occupies about 1/3 of body length, projects



Figs. 25–30. *Metopus ovalis* from life (27–30) and after protargol impregnation (25, 26). 25, 26. Infraciliature of ventral and dorsal side of same specimen. 27. Ventral view of large specimen (150 µm). The cytoplasm of the Kenyan population is strongly vacuolated. 28. Ventral view of *M. ventrosus* Vuxanovici, 1962, very likely a junior synonym of *M. ovalis* (length 100 µm; from [236]). 29. *Metopus ovalis*, length 150 µm (from [155]). 30. Ventral and dorsal view of type populations of *M. ovalis*, length 100–150 µm (from [150]). G = violet granule accumulation, PM = paroral membrane, PZ = perizonal ciliary stripe. Scale bar division 20 µm.

sharply above ventral side and left margin when cell is viewed ventrally. Postoral portion elongate to slightly saccular, broadly rounded and slightly flattened posteriorly, wrinkled by longitudinal folds after release of contractile vacuole content (Figs. 25, 27). Macronucleus slightly ellipsoidal, usually close underneath preoral dome, contains many small nucleoli. Micronucleus globular, in small indentation of macronucleus (Figs. 26, 27). Contractile vacuole in posterior end. Cortex flexible, contains loosely arranged, pale disks about 1.5 µm across, possibly mitochondria and/or fat globules, and some 0.5 µm sized granules (extrusomes ?), which do not stain with methyl green-pyronin. Cytoplasm distinctly vacuolated, possibly due to the saline environment, contains large food vacuoles with bacterial residues; no anterior granule accumulation, but cytoplasmic vacuoles often contain some granular material. Moves slowly by rotation about longitudinal body axis.

Somatic infraciliature composed of dikinetids throughout, anterior cilium lacking in many postoral and dorsal kinetids. Cilia about 15 µm long, form about 5 oblique, sigmoidal rows on preoral dome and longitudinal rows on postoral trunk. Perizonal ciliary rows on proximal edge of preoral dome, exactly S-shaped like adoral zone of membranelles, separated from first dome kinety by a distance about twice as wide as between somatic kineties, narrowly spaced with dikinetids arranged in curious, arrow-like pattern (Figs. 25, 26). About 7 postperistomial kineties. No caudal cilia.

Adoral zone of membranelles exactly S-shaped, commences on left dorsal end of preoral dome and extends sigmoidally to right side, plunging into an inconspicuous cytopharynx supplied with a long, narrow bundle of fibres. Individual membranelles cuneiform, largest bases near proximal end of zone in vivo about 5 µm wide. Paroral membrane inconspicuous because short and near proximal end of membranellar zone, consists of narrowly spaced, zigzagging, ciliated dikinetids (Figs. 25–27).

Comparison with original description and related species: My population strongly resembles the original figures (Figs. 29, 30) and description [150]. There are, as usual, some small differences, viz. the strong vacuolation of the cytoplasm and the lack of an accumulation of violet granules in the preoral dome, which seems to be more distinct in my specimens than in that figured by Kahl (Figs. 27, 30). However, these differences are small compared to the conformities, and might be related to the rather saline environment my specimens lived in. Thus, I have few doubts about the identification, but suggest that further populations should be studied.

Metopus ovalis is easily identified by its rather large size (110–200 µm) and elliptical to slightly oval shape, the lack of caudal cilia and distinct extrusomes, the el-

lipsoidal macronucleus left of the cytostome, and the relatively short and little spiraled adoral zone of membranelles, which thus extends only to mid-body and terminates on the right side. *Metopus ventrosus* Vuxanovici [236] differs from *M. ovalis* only by its more distinct oval shape, resembling that of my specimens, and the greenish colour of the anterior granule accumulation (Fig. 28). Thus, I agree with Esteban et al. [52] that these species should be synonymized. *Metopus fuscus* Kahl [150, 155] also resembles *M. ovalis* in size (180–300×40–60 µm), shape, and absence of caudal cilia. However, it has 45–50 somatic kineties [144] and a thick (2 µm) ectoplasm containing blunt protrichocysts [150, 155]. *Brachonella elongata* Jankowski [144] is smaller (100×42 µm) than *M. ovalis* and its adoral zone of membranelles extends almost to the posterior body end.


Keronopsis dieckmanni nov. spec. (Figs. 31–34, Table 6)

Diagnosis: Size in vivo about 220×70 µm, elliptical with oral area usually slightly narrowed and set off head-like. 4 macronuclear nodules. 3 short and 2 long rows of frontoventral cirri. On average 52 adoral membranelles, 61 right marginal cirri, 60 left marginal cirri, 51 cirri in right long ventral row, 53 cirri in left long ventral row, 5 cirri in frontal corona, 6 buccal cirri, 5 transverse cirri, and 3 dorsal kineties. Buccal cavity deep and semicircularly curved anteriorly.

Type location: Saline soil at shore of Lake Baringo, Kenya, equatorial Africa (about 36° E, 0°45' N).

Dedication: Named in memory of my friend and colleague Josef Dieckmann (1948–1996).

Description: Size in vivo 180–260×50–100 µm. Slenderly to moderate broadly elliptical, rarely almost parallel-sided, oral portion slightly cephalized because narrowed and due to conspicuously short adoral zone of membranelles occupying only about 25% of body length (Figs. 31, 32). Flexible and dorsoventrally flattened up to 2:1. Usually four roughly ellipsoidal macronuclear nodules one after the other in middle portion of cell slightly left of midline, sometimes indistinctly grouped forming an anterior and posterior pair, contain many globular nucleoli. Micronuclei almost spherical, compact and thus well recognizable in live specimens, near or in small indentations of macronuclei (Figs. 31, 34). Contractile vacuole distinctly above mid-body. Cortex colourless, flexible, without special granules. Cytoplasm colourless, contains some 1–3 µm sized, yellowish fat globules and, usually, many food vacuoles with large heterotrophic flagellates (*Polytoma* sp.) and/or small ciliates (*Cyclidium*, *Halteria grandinella*). Movement without peculiarities, rather slowly gliding.



Figs. 31–34. *Keronopsis dieckmanni* from life (31, 32) and after protargol impregnation (33, 34). 31. Ventral view of typical specimen packed with food vacuoles containing small ciliates and heterotrophic flagellates. 32. Dorsal view of shape variant. 33, 34. Infraciliature of ventral and dorsal side. Arrows mark breaks in left long ventral row. CV = contractile vacuole, RM = right marginal cirral row extending on dorsal side. Scale bar division 20 μ m.

Cirri fine and very narrowly spaced, marginal rows almost confluent posteriorly, right row extends on dorsal side anteriorly. Frontal corona comparatively inconspicuous because composed of only five enlarged cirri, three rightmost cirri of corona at end of short frontal rows composed of cirri gradually decreasing in size posteriorly; rightmost row possibly belongs to long left ventral row. Long ventral rows extend from anterior end to transverse cirri; cirri of same size throughout, left row possibly composed of three fragments, as indicated by two small breaks in anterior half of most specimens. Buccal cirral row conspicuous. Transverse cirri inconspicuous, hardly projecting above posterior body margin. Dorsal bristles *in vivo* about 3 μ m long, arranged in three rows almost as long as body. No caudal cirri (Figs. 31, 33, 34).

Oral apparatus of usual structure, adoral zone of membranelles, however, conspicuously short, as described above. Buccal cavity rather narrow but deep, strongly curved anteriorly. Paroral membrane almost semicircular, consists of at least three rows of basal bodies, distinctly separate from endoral membrane obliquely crossing buccal cavity. Pharyngeal fibres inconspicuous (Figs. 31, 33).

Comparison with related species and generic classification: *K. dieckmanni* resembles *K. tasmaniensis* Blatterer & Foissner [20] which, however, has only 2 macronuclear nodules, 1–2 buccal cirri, and 2 short frontal cirral rows. *Keronopsis belluo* [179], type of genus, is also rather similar to *K. dieckmanni* because it has 5–6 macronuclear nodules, many buccal cirri, a similar size (250–300 μ m) and buccal field, and lives in

Table 6. Morphometric data from *Keronopsis dieckmanni*.

Character ¹⁾	\bar{x}	M	SD	SE	CV	Min	Max	n
Body, length	201.8	197.5	26.8	8.5	13.3	165.0	245.0	10
Body, width	67.8	68.5	11.8	3.7	17.4	48.0	91.0	10
Anterior somatic end to proximal end of adoral zone, distance	50.5	50.0	5.2	1.6	10.2	42.0	60.0	10
Anterior somatic end to posterior end of right long ventral row, distance	182.5	175.0	24.6	7.8	13.5	150.0	225.0	10
Anterior somatic end to posterior end of left long ventral row, distance	169.4	162.0	29.0	9.2	17.1	138.0	221.0	10
Anterior somatic end to transverse cirri, distance	181.5	176.0	23.3	7.4	12.8	150.0	225.0	10
Macronuclear nodules, length	19.9	19.0	4.3	1.4	21.8	15.0	26.0	10
Macronuclear nodules, width	12.4	12.5	1.7	0.5	13.8	10.0	16.0	10
Macronuclear figure, length	103.5	96.0	18.8	5.9	18.2	80.0	138.0	10
Macronuclear nodules, number	3.9	4.0	0.9	0.3	22.5	2.0	5.0	10
Micronuclei, length	4.2	4.0	0.6	0.2	14.0	3.5	5.0	10
Micronuclei, width	3.6	3.5	—	—	—	3.0	4.0	10
Micronuclei, number	3.1	3.0	1.1	0.4	33.9	1.0	4.0	9
Adoral membranelles, number	50.7	52.5	6.1	1.9	12.0	40.0	60.0	10
Right marginal cirri, number	64.2	61.0	7.0	2.3	10.9	55.0	75.0	9
Left marginal cirri, number	62.3	60.0	8.7	2.9	14.0	51.0	74.0	9
Anterior frontal cirri, number	5.0	5.0	0.0	0.0	0.0	5.0	5.0	10
Buccal cirri, number	5.6	6.0	—	—	—	5.0	6.0	8
Cirri in right ventral row, number	50.3	51.0	5.6	1.9	11.2	42.0	58.0	9
Cirri in left ventral row, number ²⁾	54.0	53.0	9.4	3.3	17.4	41.0	68.0	8
Transverse cirri, number	5.2	5.0	1.0	0.3	19.9	4.0	7.0	10
Dorsal kineties, number	3.0	3.0	0.0	0.0	0.0	3.0	3.0	9

¹⁾ Data based on protargol-impregnated and mounted specimens from field. Measurements in μm . Abbreviations: CV = coefficient of variation in %, M = median, Max = maximum, Min = minimum, n = number of individuals investigated, SD = standard deviation, SE = standard deviation of mean, \bar{x} = arithmetic mean.

²⁾ Including rightmost short row of frontal cirri.

mosses. However, *K. belluo* lacks short frontal rows and has at least 10 cirri in the frontal cirral corona.

Both *Keronopsis* and *Paraholosticha* have a frontal cirral corona and a very similar general organization. However, I agree with Dieckmann [36] that the presence (*Keronopsis*), respectively, absence (*Paraholosticha*) of transverse cirri is sufficient to maintain both genera. The frontal cirral corona of *K. dieckmanni* and *K. tasmaniensis* is rather inconspicuous because it consists of only 5–7 cirri, whereas it is composed of 10–20 cirri in the other members of the genus [8, 36, 37, 75]. Thus, the generic classification of *K. dieckmanni* and *K. tasmaniensis* might be questioned. However, both have, like the other *Keronopsis* and *Paraholosticha* species, 3 dorsal kineties and lack caudal cirri.


Tachysoma humicola longisetum nov. subspec. (Figs. 35–38, Table 7)

Diagnosis: Like *T. humicola* Gellért, 1957, but posteriormost bristles of dorsal kineties 1–3 distinctly elongated (10 μm).

Type location: Soil from Shetani volcano area, Tsavo National Park, Kenya, equatorial Africa (about 38° E, 2°55' N).

Etymology: *longisetum* (long bristles) refers to the main subspecies character, viz. the elongated cilia of the posteriormost dorsal kinetids.

Description: Size *in vivo* 40–55×20–30 μm , usually about 45×25 μm . Shape conspicuous because of its regularity, i.e. slenderly to broadly elliptical (Figs. 35, 36), often fusiform after protargol impregnation (Figs. 37). Rather inflexible like, e.g. *Sterkiella histriomuscorum*, and dorsoventrally flattened up to 2:1. Macronuclear nodules slightly ellipsoidal, in middle third of cell to left of midline, contain many granular nucleoli. Micronucleus globular, invariably between macronuclear nodules (Figs. 35, 38). Contractile vacuole in mid-body near left margin, without distinct collecting canals. Cortex colourless, rather inflexible, without special granules. Cells usually very transparent, contain 10–20 curved crystals (3 μm) and some 3–4 μm sized food vacuoles with bacterial residues; posterior end often darkly granulated after protargol impregnation (Fig. 38).



Figs. 35–38. *Tachysoma humicola longisetum* from life (35, 36) and after protargol impregnation (37, 38). 35. Ventral view of typical specimen. 36. Dorsal view of slender shape variant. 37, 38. Infraciliature of ventral and dorsal side. Arrowheads mark two dorsal bristles between which the distance is slightly increased. AM = distalmost adoral membranelle, DB = elongated dorsal cilia, FC = rightmost frontal cirrus, MC = last marginal cirrus of right row, PC = pretransverse cirrus. Scale bar division 10 µm.

Movement conspicuous due to long periods of gliding interrupted by short, fast jumps.

Cirri conspicuously thick compared to size of cell, about 10 µm long; transverse cirri, however, about 15 µm long and distinctly projecting above posterior body margin, conspicuously enlarged and slightly sigmoidal. Marginal rows widely open at both ends, posterior gap occupied by transverse cirri; remarkably short because commencing at level of proximal end of adoral zone of membranelles and terminating at level of uppermost transverse cirri. Right pretransverse cirrus difficult to identify because usually almost in line with right marginal row; likewise, rightmost anterior frontal cirrus difficult to recognize because in line with distalmost adoral membranelles (Figs. 35, 37). Dorsal cilia *in vivo* about 3 µm long, those of posteriormost kinetid in kinetics 1–3, however, elongated to 10 µm; arranged in four rows with row 4 terminating in mid-body; one kinetid frequently lacking near posterior end of kinety 3 (Figs. 36, 38).

Adoral zone of usual structure, distalmost three to four membranelles, however, slightly separate from rest

of zone and with cilia more distinctly curved rightwards than those of other anterior membranelles; occupies about 34% of body length, bases of largest membranelles about 5 µm wide. Buccal cavity narrow and flat. Undulating membranes almost straight and side by side, paroral shorter than endoral, forms thick line along its posterior half; endoral composed of dikanetids separated slightly Y-like in anterior third of membrane (Figs. 35, 37).

Comparison with related species: *T. humicola longisetum* highly resembles *T. humicola* Gellér [130] as redescribed by Foissner [69], both in general morphology and morphometry (Tab. 7). For instance, both have the right pretransverse cirrus close to the last cirrus of the right marginal cirral row. The only significant difference concerns the posteriormost dorsal bristles, which are distinctly elongated in the African population. Admittedly, this is a fairly indistinct character. However, I found such populations at several sites in Africa and South America, indicating that they are stable modifications. Thus, separation at subspecies rank seems appropriate.

Table 7. Morphometric data from *Tachysoma humicola longisetum* (upper line) and *T. humicola humicola* (lower line; from [69]).

Character ¹⁾	\bar{x}	M	SD	SE	CV	Min	Max	n
Body, length	44.0	45.0	3.9	1.1	8.8	37.0	50.0	13
	42.4	41.0	4.3	1.3	10.0	38.0	50.0	11
Body, width	21.2	21.0	2.4	0.7	11.2	18.0	27.0	13
	21.0	20.0	2.6	0.8	12.4	18.0	27.0	11
Anterior somatic end to proximal end of adoral zone, distance	14.6	15.0	0.7	0.2	5.3	13.0	15.0	13
	14.0	14.0	0.9	0.3	6.4	13.0	16.0	11
Anterior somatic end to anterior end of right marginal row, distance	14.2	14.0	1.2	0.3	8.2	12.0	16.0	13
	14.0	14.0	0.6	0.2	4.5	13.0	15.0	11
Anterior somatic end to anterior end of left marginal row, distance	14.2	14.0	1.2	0.3	8.6	12.0	16.0	13
	13.1	13.0	1.0	0.3	8.0	11.0	15.0	11
Macronuclear nodules, length	7.2	7.0	1.3	0.4	18.8	6.0	10.0	13
	7.8	7.0	1.8	0.5	22.9	5.6	10.0	11
Macronuclear nodules, width	5.4	5.0	0.7	0.2	14.3	4.0	7.0	13
	5.7	5.6	0.4	0.1	6.3	4.8	6.0	11
Micronucleus, diameter	2.3	2.2	0.3	0.1	11.8	2.0	3.0	13
	2.3	2.4	0.4	0.1	15.1	1.6	2.8	11
Macronuclear nodules, number	2.0	2.0	0.0	0.0	0.0	2.0	2.0	13
	2.0	2.0	0.0	0.0	0.0	2.0	2.0	11
Micronuclei, number	1.0	1.0	0.0	0.0	0.0	1.0	1.0	13
	1.1	1.0	—	—	—	1.0	2.0	11
Adoral membranelles, number	15.0	15.0	0.0	0.0	0.0	15.0	15.0	13
	14.8	15.0	0.6	0.2	4.1	14.0	16.0	11
Right marginal cirri, number	6.0	6.0	0.0	0.0	0.0	6.0	6.0	13
	6.4	6.0	0.7	0.2	10.7	6.0	8.0	11
Left marginal cirri, number	7.1	7.0	—	—	—	7.0	8.0	13
	7.2	7.0	—	—	—	7.0	8.0	11
Anterior frontal cirri, number	3.0	3.0	0.0	0.0	0.0	3.0	3.0	13
	3.0	3.0	0.0	0.0	0.0	3.0	3.0	11
Posterior frontal cirri, number	4.0	4.0	0.0	0.0	0.0	4.0	4.0	13
	4.0	4.0	0.0	0.0	0.0	4.0	4.0	11
Buccal cirri, number	1.0	1.0	0.0	0.0	0.0	1.0	1.0	13
	1.0	1.0	0.0	0.0	0.0	1.0	1.0	11
Postoral cirri, number	3.0	3.0	0.0	0.0	0.0	3.0	3.0	13
	3.0	3.0	0.0	0.0	0.0	3.0	3.0	11
Pretransverse cirri, number	2.0	2.0	0.0	0.0	0.0	2.0	2.0	13
	2.0	2.0	0.0	0.0	0.0	2.0	2.0	11
Transverse cirri, number	5.0	5.0	0.0	0.0	0.0	5.0	5.0	13
	5.0	5.0	0.0	0.0	0.0	5.0	5.0	11
Dorsal kinetics, number	4.0	4.0	0.0	0.0	0.0	4.0	4.0	13
	4.0	4.0	0.0	0.0	0.0	4.0	4.0	11

¹⁾ Data based on protargol-impregnated and mounted specimens from field. Measurements in μm . Abbreviations: CV = coefficient of variation in %, M = median, Max = maximum, Min = minimum, n = number of individuals investigated, SD = standard deviation, SE = standard deviation of mean, \bar{x} = arithmetic mean.

Acknowledgements: Supported by the Austrian Science Foundation (FWF project P10264-BIO). The technical assistance of Dr. Eva Herzog and Mag. Eric Strobl is greatly acknowledged. Special thanks to my wife, Prof. Dr. Ilse Foissner, for assisting me in sampling.

References

- 1 Aesch E. and Foissner W. (1992): Biology of a high-rate activated sludge plant of a pharmaceutical company. Arch. Hydrobiol. Suppl., 90, 207–251.
- 2 Aesch E. and Foissner W. (1993): Effects of organically enriched magnesite fertilizers on the soil ciliates of a spruce forest. Pedobiologia, 38, 321–335.
- 3 Alekperov I. Kh. (1991): Two new species *Thylakidium* (Bursariidae, Ciliophora) from the soils of the Apsheron peninsula. Zool. Zh., 70, 127–129 (in Russian with English summary).
- 4 Alekperov I. Kh. (1993): Free-living ciliates in the soils of St. Petersburg parks (Protozoa). Zoosyst. Rossica, 2, 13–28.
- 5 Alekperov I. Kh. and Musayev M. A. (1988): New and rare free-living infusoria from fresh waters and soil of the Apsheron peninsula. Zool. Zh., 67, 1904–1909 (in Russian with English summary).

- 6 Augustin H. and Foissner W. (1992): Morphologie und Ökologie einiger Ciliaten (Protozoa: Ciliophora) aus dem Belebtschlamm. Arch. Protistenk., 141, 243–283.
- 7 Batisse A. (1967): Ultrastructure de la coque et du disque chez le genre *Metacineta* Butschli. Protistologica, 3, 395–411.
- 8 Berger H. and Foissner W. (1987): Morphology and biometry of some soil hypotrichs (Protozoa, Ciliophora). Zool. Jb. Syst., 114, 193–239.
- 9 Berger H. and Foissner W. (1988a): Revision of *Lamnostyla* Buitkamp, 1977 and description of *Territricha* nov. gen. (Ciliophora: Hypotrichida). Zool. Anz., 220, 113–134.
- 10 Berger H. and Foissner W. (1988b): The morphogenesis of *Kahlialiella franzii* (Foissner, 1982) nov. comb. and *Oxytricha gigantea* Horváth, 1933 (Ciliophora, Hypotrichida). Arch. Protistenk., 136, 65–77.
- 11 Berger H. and Foissner W. (1989a): Morphology and biometry of some soil hypotrichs (Protozoa, Ciliophora) from Europe and Japan. Bull. Br. Mus. nat. Hist. (Zool.), 55, 19–46.
- 12 Berger H. and Foissner W. (1989b): Morphology and morphogenesis of *Parakahlialiella baideri* nov. spec. (Ciliophora, Hypotrichida). Bull. Br. Mus. nat. Hist. (Zool.), 55, 11–17.
- 13 Berger H., Foissner W., and Adam H. (1983): Morphology and morphogenesis of *Fuscheria terricola* n. sp. and *Spathidium muscorum* (Ciliophora: Kinetofragminophora). J. Protozool., 30, 529–535.
- 14 Berger H., Foissner W., and Adam H. (1984): Taxonomie, Biometrie und Morphogenese einiger terricoler Ciliaten (Protozoa: Ciliophora). Zool. Jb. Syst., 111, 339–367.
- 15 Berger H., Foissner W., and Adam H. (1985): Morphological variation and comparative analysis of morphogenesis in *Parakahlialiella macrostoma* (Foissner, 1982) nov. gen. and *Histrichus muscorum* (Kahl, 1932), (Ciliophora, Hypotrichida). Protistologica, 21, 295–311.
- 16 Berthold A. (1994): Zooökologische Untersuchungen der Wimpertierchen (Ciliophora, Protozoa) in der Umgebung von Brixlegg mit einer allgemeinen Einführung und einem Überblick über Schwermetalleffekte und Toxizitätstests. Reports Federal Environment Agency, Austria, UBA-94-99b, 1–69 (each contribution paginated separately!).
- 17 Bick H. and Buitkamp U. (1976): Ciliated protozoa from Canadian grassland soils. Trans. Am. microsc. Soc., 95, 490–491.
- 18 Biczok F. (1956): Morphologische und physiologische Untersuchungen an einer neuen *Pyxidium*-Art. Acta biol., Szeged., 2, 155–165.
- 19 Biegel M. (1954): Beitrag zur Peritrichenfauna der Umgebung Erlangens. Arch. Protistenk., 100, 153–182.
- 20 Blatterer H. and Foissner W. (1988): Beitrag zur terricolen Ciliatenfauna (Protozoa, Ciliophora) Australiens. Staphia (Linz), 17, 1–84.
- 21 Blatterer H. and Foissner W. (1992): Morphology and infraciliature of some cyrtophorid ciliates (Protozoa, Ciliophora) from freshwater and soil. Arch. Protistenk., 142, 101–118.
- 22 Borror A. C. and Evans F. R. (1979): *Cladotricha* and phylogeny in the suborder Stichotrichina (Ciliophora, Hypotrichida). J. Protozool., 26, 51–55.
- 23 Borror A. C. and Wicklow B. J. (1983): The suborder Urostylina Jankowski (Ciliophora, Hypotrichida): morphology, systematics and identification of species. Acta Protozool., 22, 97–126.
- 24 Bowers N. J. and Pratt J. R. (1995): Estimation of genetic variation among soil isolates of *Colpoda inflata* (Stokes) (Protozoa: Ciliophora) using the polymerase chain reaction and restriction fragment length polymorphism analysis. Arch. Protistenk., 145, 29–36.
- 25 Buitkamp U. (1975): The morphogenesis of *Oxytricha agilis* Engelmann (Ciliata, Hypotrichida). Acta Protozool., 14, 67–74.
- 26 Buitkamp U. (1977a): Über die Ciliatenfauna zweier mitteleuropäischer Bodenstandorte (Protozoa; Ciliata). Decheniana (Bonn), 130, 114–126.
- 27 Buitkamp U. (1977b): Die Ciliatenfauna der Savanne von Lamto (Elfenbeinküste). Acta Protozool., 16, 249–276.
- 28 Buitkamp U. and Wilbert N. (1974): Morphologie und Taxonomie einiger Ciliaten eines kanadischen Präriebodens. Acta Protozool., 13, 201–210.
- 29 Certes A. (1891): Note sur deux infusoires nouveaux des environs de Paris. Mém. Soc. zool. Fr., 4, 536–541.
- 30 Chardez D. (1967): Infusoires ciliés terricoles (Protozoa, Infusoria Ciliata). Revue Écol. Biol. Sol., 4, 289–298.
- 31 Corliss J. O. (1952): Le cycle autogamique de *Tetrahymena rostrata*. C. r. hebd. Séanc. Acad. Sci., Paris, 235, 399–402.
- 32 Corliss J. O. (1958): The systematic position of *Pseudomicrothorax dubius*, ciliate with a unique combination of anatomical features. J. Protozool., 5, 184–193.
- 33 Corliss J. O. (1960): The problem of homonyms among generic names of ciliated protozoa, with proposal of several new names. J. Protozool., 7, 269–278.
- 34 Corliss J. O. (1979): The ciliated protozoa. Characterization, classification and guide to the literature. 2nd ed. Pergamon Press, Oxford, New York.
- 35 Detcheva R. B. (1992): Protozoa, Ciliophora. Catalogi Faunae Bulgaricae, 1, 1–136.
- 36 Dieckmann J. (1988): Infraciliature and morphogenesis of *Parabolosticha sterkii* (Garnjobst, 1934) n. comb. (Ciliophora, Hypotrichida). Europ. J. Protistol., 23, 218–228.
- 37 Dieckmann J. (1989): Neubeschreibung der Morphogenese von *Parabolosticha muscicola* Kahl, 1932 (Ciliophora, Hypotrichida). Arch. Protistenk., 137, 143–156.
- 38 Dingfelder J. H. (1962): Die Ciliaten vorübergehender Gewässer. Arch. Protistenk., 105, 509–658.
- 39 Dragesco J. (1963): Révision du genre *Dileptus* Dujardin 1841 (Ciliata Holotricha) (systématique, cytologie, biologie). Bull. biol. Fr. Belg., 97, 103–145.
- 40 Dragesco J. (1970): Ciliés libres du Cameroun. Annls. Fac. Sci. Univ. féd. Cameroun (Numéro Hors-série), year 1970, 1–141.
- 41 Dragesco J. and Dragesco-Kerneis A. (1979): Ciliés muscicoles nouveaux ou peu connus. Acta Protozool., 18, 401–416.
- 42 Dragesco J. and Dragesco-Kerneis A. (1986): Ciliés libres de l'Afrique intertropicale. Faune tropicale, 26, 1–559.
- 43 Dragesco J. and Njiné T. (1971): Compléments à la connaissance des ciliés libres du Cameroun. Annls. Fac. Sci. Univ. féd. Cameroun, 7–8, 97–140.
- 44 Eigner P. (1994): Divisional morphogenesis and reorganization in *Eschaneustyla brachytoma* Stokes, 1886 and revision of the Bakuellinae (Ciliophora, Hypotrichida). Europ. J. Protistol., 30, 462–475.

- 45 Eigner P. (1995): Divisional morphogenesis in *Deviata abbrevescens* nov. gen., nov. spec., *Neogeneia hortualis* nov. gen., nov. spec., and *Kahliella simplex* (Horváth) Corliss and redefinition of the Kahliellidae (Ciliophora, Hypotrichida). *Europ. J. Protistol.*, 31, 341–366.
- 46 Eigner P. and Foissner W. (1991): *Orthoamphisiella stramenticola* nov. gen., nov. spec., a new hypotrichous ciliate (Ciliophora: Hypotrichida) occurring in walnut leaf litter. *Acta Protozool.*, 30, 129–133.
- 47 Eigner P. and Foissner W. (1992): Divisional morphogenesis in *Bakuella pampinaria* nov. spec. and reevaluation of the classification of the urostylids (Ciliophora, Hypotrichida). *Europ. J. Protistol.*, 28, 460–470.
- 48 Eigner P. and Foissner W. (1993) Divisional morphogenesis in *Orthoamphisiella stramenticola* Eigner & Foissner, 1991 and *O. grelli* nov. spec. (Ciliophora, Hypotrichida). *Arch. Protistenk.*, 143, 337–345.
- 49 Eigner P. and Foissner W. (1994): Divisional morphogenesis in *Amphisiellides illuvialis* n. sp., *Paramphisiella caudata* (Hemberger) and *Hemiamphisiella terricola* Foissner, and redefinition of the Amphisiellidae (Ciliophora, Hypotrichida). *J. Euk. Microbiol.*, 41, 243–261.
- 50 Eisler K. and Bardele C. F. (1986): Cortical morphology and morphogenesis of the nassulid ciliates *Furgasonia blochmanni* Fauré-Fremiet, 1967 and *Nassula citrea* Kahl, 1930. *Protistologica*, 22, 461–476.
- 51 Engelmann T. W. (1862): Zur Naturgeschichte der Infusorienstiere. *Z. wiss. Zool.*, 11, 347–393.
- 52 Esteban G., Fenchel T., and Finlay B. (1995): Diversity of free-living morphospecies in the ciliate genus *Metopus*. *Arch. Protistenk.*, 146, 137–164.
- 53 Evans F. R. and Thompson J. C., Jr. (1964): Pseudocohnilembidae n. fam., a hymenostome ciliate family containing one genus, *Pseudocohnilembus* n. g., with three new species. *J. Protozool.*, 11, 344–352.
- 54 Fernández-Galiano D. and Calvo P. (1992): Redescription of *Phacodinium metchnikoffi* (Ciliophora, Hypotrichida): general morphology and taxonomic position. *J. Protozool.*, 39, 443–448.
- 55 Finlay B. J., Corliss J. O., Esteban G., and Fenchel T. (1996): Biodiversity at the microbial level: the number of free-living ciliates in the biosphere. *Q. Rev. Biol.*, 71, 221–237.
- 56 Foissner W. (1975): Opisthonectidae (Ciliata, Peritrichida) nov. fam. und Revision der Genera *Telotrochidium* (Kent) und *Opisthonecta* (Fauré-Fremiet). *Protistologica*, 11, 395–414.
- 57 Foissner W. (1977): Revision der genera *Astylozoon* (Engelmann) und *Hastatella* (Erlanger) (Ciliata Natantina). *Protistologica*, 13, 353–379.
- 58 Foissner W. (1978a): Morphologie, Infraciliatur und Silberliniensystem von *Plagiocampa rouxi* Kahl, 1926 (Prostomatida, Plagiocampidae) und *Balanonema sapropelica* nov. spec. (Philasterina, Loxocephalidae). *Protistologica*, 14, 381–389.
- 59 Foissner W. (1978b): *Opisthonecta bivacuolata* nov. spec., *Telotrochidium cylindricum* nov. spec. und *Epistylis alpestris* nov. spec., drei neue peritrichie Ciliaten aus dem Hochgebirge (Hohe Tauern, Österreich). *Annl. naturh. Mus. Wien*, 81, 549–565.
- 60 Foissner W. (1979): Peritrichie Ciliaten (Protozoa: Ciliophora) aus alpinen Kleingewässern. *Zool. Jb. Syst.*, 106, 529–558.
- 61 Foissner W. (1980): Taxonomische Studien über die Ciliaten des Grossglocknergebietes (Hohe Tauern, Österreich). IX. Ordnungen Heterotrichida und Hypotrichida. *Ber. Nat.-Med. Ver. Salzburg*, 5, 71–117.
- 62 Foissner W. (1981a): Die Gemeinschaftsstruktur der Ciliatenzönose in alpinen Böden (Hohe Tauern, Österreich) und Grundlagen für eine Synökologie der terrestrischen Ciliaten (Protozoa, Ciliophora). *Veröff. Österr. MaB-Programms*, 4, 7–52.
- 63 Foissner W. (1981b): Morphologie und Taxonomie einiger heterotricher und peritricher Ciliaten (Protozoa: Ciliophora) aus alpinen Böden. *Protistologica*, 17, 29–43.
- 64 Foissner W. (1981c): Morphologie und Taxonomie einiger neuer und wenig bekannter kinetofragminophorer Ciliaten (Protozoa: Ciliophora) aus alpinen Böden. *Zool. Jb. Syst.*, 108, 264–297.
- 65 Foissner W. (1982): Ökologie und Taxonomie der Hypotrichida (Protozoa: Ciliophora) einiger österreichischer Böden. *Arch. Protistenk.*, 126, 19–143.
- 66 Foissner W. (1983a): Die Morphogenese von *Urosoma macrostyla* (Wrześniowski, 1870) (Ciliophora: Oxytrichidae). *Arch. Protistenk.*, 127, 413–428.
- 67 Foissner W. (1983b): Morphologie und Morphogenese von *Psilotricha succisa* (O. F. Müller, 1786) nov. comb. (Ciliophora, Hypotrichida). *Protistologica*, 19, 479–493.
- 68 Foissner W. (1983c): Taxonomische Studien über die Ciliaten des Großglocknergebietes (Hohe Tauern, Österreich) I. Familien Holophryidae, Prorodontidae, Plagiocampidae, Colepidae, Enchelyidae und Lacrymariaidae nov. fam. *Annl. naturh. Mus. Wien*, 84B, 49–85.
- 69 Foissner W. (1984): Infraciliatur, Silberliniensystem und Biometrie einiger neuer und wenig bekannter terrestrischer, limnischer und mariner Ciliaten (Protozoa: Ciliophora) aus den Klassen Kinetofragminophora, Colpodea und Polyhymenophora. *Stapfia* (Linz) 12, 1–165.
- 70 Foissner W. (1985): Morphologie und Infraciliatur der Genera *Microthorax* und *Stammeridium* und Klassifikation der Microthoracina Jankowski, 1967 (Protozoa: Ciliophora). *Zool. Anz.*, 214, 33–53.
- 71 Foissner W. (1986a): Beitrag zur Kenntnis der Bodenciliaten (Protozoa: Ciliophora) des Himalaja. *Zool. Jb. Syst.*, 113, 45–53.
- 72 Foissner W. (1986b): Wimpertiere (Protozoa: Ciliophora) in Flechten. Kataloge des Oberösterreichischen Landesmuseums Linz (N.F.), 5, 43–46.
- 73 Foissner W. (1987a): Soil protozoa: fundamental problems, ecological significance, adaptations in ciliates and testaceans, bioindicators, and guide to the literature. *Progr. Protistol.*, 2, 69–212.
- 74 Foissner W. (1987b): Faunistische und taxonomische Notizen über die Protozoen des Fuscher Tales (Salzburg, Österreich). *Jb. Haus Nat. Salzburg*, 10, 56–68.
- 75 Foissner W. (1987c): Neue und wenig bekannte hypotrichie und colpodide Ciliaten (Protozoa: Ciliophora) aus Böden und Moosen. *Zool. Beitr. (N. F.)*, 31, 187–282.
- 76 Foissner W. (1987d): Miscellanea nomenclatorica ciliata (Protozoa: Ciliophora). *Arch. Protistenk.*, 133, 219–235.
- 77 Foissner W. (1987e): Neue terrestrische und limnische Ciliaten (Protozoa, Ciliophora) aus Österreich und Deutschland. *Sber. Akad. Wiss. Wien*, 195, 217–268.
- 78 Foissner W. (1988a): Gemeinsame Arten in der terrestrischen Ciliatenfauna (Protozoa: Ciliophora) von Australien und Afrika. *Stapfia* (Linz), 17, 85–133.

- 79 Foissner W. (1988b): Taxonomie und Ökologie einiger Ciliaten (Protozoa, Ciliophora) des Saprobiensystems. II. Familie Chilodonellidae. *Hydrobiologia*, 162, 21–45.
- 80 Foissner W. (1988c): Taxonomic and nomenclatural revision of Sládeček's list of ciliates (Protozoa: Ciliophora) as indicators of water quality. *Hydrobiologia*, 166, 1–64.
- 81 Foissner W. (1989): Morphologie und Infraciliatur einiger neuer und wenig bekannter terrestrischer und limnischer Ciliaten (Protozoa, Ciliophora). *Sber. Akad. Wiss. Wien*, 196, 173–247.
- 82 Foissner W. (1991): Basic light and scanning electron microscopic methods for taxonomic studies of ciliated protozoa. *Europ. J. Protistol.*, 27, 313–330.
- 83 Foissner W. (1992): Estimating the species richness of soil protozoa using the "non-flooded petri dish method". In: Lee J. J. and Soldo A. T. (eds.): *Protocols in Protozoology*. Allen Press, Lawrence, Kansas. B-10.1–B-10.2.
- 84 Foissner W. (1993a): Colpodea (Ciliophora). *Protozoenfauna*, 4/1, I–X + 1–798.
- 85 Foissner W. (1993b): *Idiocolpoda pelobia* gen. n., sp. n., a new colpodid ciliate (Protozoa, Ciliophora) from an ephemeral stream in Hawaii. *Acta Protozool.*, 32, 175–182.
- 86 Foissner W. (1993c): *Corticocolpoda kaneshiroae* n. g., n. sp., a new colpodid ciliate (Protozoa, Ciliophora) from the bark of Ohia trees in Hawaii. *J. Euk. Microbiol.*, 40, 764–775.
- 87 Foissner W. (1994a): *Spetazoon australiense* nov. gen., nov. spec., ein neues Wimpertier (Protozoa, Ciliophora) von Australien. *Kataloge des Oberösterreichischen Landesmuseums Linz* (N.F.), 71, 267–278.
- 88 Foissner W. (1994b): Morphology and morphogenesis of *Circinella arenicola* nov. gen., nov. spec., a cephalized hypotrich (Ciliophora, Hypotrichida) from sand dunes in Utah, USA. *Europ. J. Protistol.*, 30, 156–170.
- 89 Foissner W. (1994c): *Bryometopus hawaiiensis* sp. n., a new colpodid ciliate from a terrestrial biotope of the Hawaiian Archipelago (Protozoa: Ciliophora). *Annl. naturh. Mus. Wien*, 96B, 19–27.
- 90 Foissner W. (1994d): *Pentabyxena corticicola* nov. gen., nov. spec., a new colpodid ciliate (Protozoa, Ciliophora) from bark of *Acacia* trees in Costa Rica. *Arch. Protistenk.*, 144, 289–295.
- 91 Foissner W. (1994e): Soil protozoa as bioindicators in ecosystems under human influence. In: Derbyshire J. F. (ed.): *Soil Protozoa*, pp. 147–193. CAB International, Wallingford, Oxon, England.
- 92 Foissner W. (1994f): Die Urtiere (Protozoen) des Bodens. *Kataloge des Oberösterreichischen Landesmuseums Linz* (N.F.), 71, 169–218.
- 93 Foissner W. (1995): Tropical protozoan diversity: 80 ciliate species (Protozoa, Ciliophora) in a soil sample from a tropical dry forest of Costa Rica, with descriptions of four new genera and seven new species. *Arch. Protistenk.*, 145, 37–79.
- 94 Foissner W. (1996a): Terrestrial ciliates (Protozoa, Ciliophora) from two islands (Gough, Marion) in the southern oceans, with description of two new species, *Arcuospathidium cooperi* and *Oxytricha ottowi*. *Biol. Fertil. Soils*, 23, 282–291.
- 95 Foissner W. (1996b): Faunistics, taxonomy and ecology of moss and soil ciliates (Protozoa, Ciliophora) from Antarctica, with description of new species, including *Pleuroplitoides smithi* gen. n., sp. n. *Acta Protozool.*, 35, 95–123.
- 96 Foissner W. (1997a): Global soil ciliate (Protozoa, Ciliophora) diversity: a probability-based approach using large sample collections from Africa, Australia and Antarctica. *Biodiv. Conserv.*, 6, 1627–1638.
- 97 Foissner W. (1997b): Soil ciliates (Protozoa: Ciliophora) from evergreen rain forests of Australia, South America and Costa Rica: diversity and description of new species. *Biol. Fertil. Soils*, 25, 317–339.
- 98 Foissner W. (1997c): Faunistic and taxonomic studies on ciliates (Protozoa, Ciliophora) from clean rivers in Bavaria (Germany), with descriptions of new species and ecological notes. *Limnologica (Berlin)*, 27, 179–238.
- 99 Foissner W. and Adam H. (1981): Morphologie und Infraciliatur von *Parafurgasonia sorex* (Penard, 1922) nov. gen. und *Obertrumia georgiana* (Dragesco, 1972) nov. gen. (Protozoa: Ciliophora). *Zool. Anz.*, 207, 303–319.
- 100 Foissner W. and Adam H. (1983a): Die Morphogenese von *Urosomoida agiliformis* Foissner, 1982 (Ciliophora, Oxytrichidae). *Zool. Anz.*, 211, 161–176.
- 101 Foissner W. and Adam H. (1983b): Morphologie und Morphogenese des Bodenciliaten *Oxytricha granulifera* sp. n. (Ciliophora, Oxytrichidae). *Zool. Scr.*, 12, 1–11.
- 102 Foissner W. and Didier P. (1981): Morphologie und Infraciliatur einiger kinetofragminophorer und hypotricher Ciliaten aus den Fließgewässern von Besse-en-Chandesse (Frankreich). *Annls. Stn. limnol. Besse*, 15, 254–275.
- 103 Foissner W. and Foissner I. (1985): Oral monokinetids in the free-living haptorid ciliate *Enchelydium polynucleatum* (Ciliophora, Enchelydiidae): ultrastructural evidence and phylogenetic implications. *J. Protozool.*, 32, 712–722.
- 104 Foissner W. and Foissner I. (1988): The fine structure of *Fuscheria terricola* Berger et al., 1983 and a proposed new classification of the subclass Haptoria Corliss, 1974 (Ciliophora, Litostomatea). *Arch. Protistenk.*, 135, 213–235.
- 105 Foissner W. and Foissner I. (1994): Fine structure of *Cosmocolpoda naschbergeri* (Ciliophora, Colpodida). *Arch. Protistenk.*, 144, 129–136.
- 106 Foissner W. and Foissner I. (1995): Fine structure and systematic position of *Enchelyomorpha vermicularis* (Smith, 1899) Kahl, 1930, an anaerobic ciliate (Protozoa, Ciliophora) from domestic sewage. *Acta Protozool.*, 34, 21–34.
- 107 Foissner W. and O'Donoghue P. J. (1990): Morphology and infraciliature of some freshwater ciliates (Protozoa: Ciliophora) from western and south Australia. *Invertebr. Taxon.*, 3, 661–696.
- 108 Foissner W. and Peer T. (1985): Protozoologische Untersuchungen an Almböden im Gasteiner Tal (Zentralalpen, Österreich). I. Charakteristik der Taxotope, Faunistik und Autökologie der Testacea und Ciliophora. *Veröff. Österr. MaB-Programms*, 9, 27–50.
- 109 Foissner W. and Schiffmann H. (1974): Vergleichende Studien an argyrophilen Strukturen von vierzehn peritrichen Ciliaten. *Protistologica*, 10, 489–508.
- 110 Foissner W. and Wilbert N. (1981): A comparative study of the infraciliature and silverline system of the fresh-water scuticociliates *Pseudocohnilembus putrinus* (Kahl, 1928) nov. comb., *P. pusillus* (Quennerstedt, 1869) nov. comb., and the marine form *P. marinus* Thompson, 1966. *J. Protozool.*, 28, 291–297.

- 111 Foissner W., Adam H., and Foissner I. (1982a): Morphologie und Infraciliatur von *Bryometopus pseudochilodon* Kahl, 1932, *Balantidiooides dragescoi* nov. spec. und *Kahliella marina* nov. spec. und Revision des Genus *Balantidiooides* Penard, 1930 (Protozoa, Ciliophora). *Protistologica*, 18, 211–225.
- 112 Foissner W., Adam H., and Foissner I. (1982b): Morphologie, Infraciliatur und Silberliniensystem einiger wenig bekannter Scuticociliatida (Protozoa: Ciliophora). *Zool. Jb. Syst.*, 109, 443–468.
- 113 Foissner W., Peer T., and Adam H. (1985): Pedologische und protozoologische Untersuchung einiger Böden des Tullnerfeldes (Niederösterreich). *Mitt. öst. bodenk. Ges.*, 30, 77–117.
- 114 Foissner W., Blatterer H., Berger H., and Kohmann F. (1991): Taxonomische und ökologische Revision der Ciliaten des Saprobiensystems. Band I: Cyrtophorida, Oligotrichida, Hypotrichia, Colpodea. *Informationsberichte des Bayer. Landesamtes für Wasserwirtschaft*, 1/91, 1–478.
- 115 Foissner W., Berger H. und Kohmann F. (1992): Taxonomische und ökologische Revision der Ciliaten des Saprobiensystems. Band II: Peritrichia, Heterotrichida, Odontostomatida. *Informationsberichte des Bayer. Landesamtes für Wasserwirtschaft*, 5/92, 1–502.
- 116 Foissner W., Berger H., and Kohmann F. (1994): Taxonomische und ökologische Revision der Ciliaten des Saprobiensystems. Band III: Hymenostomata, Prostomatida, Nassulida. *Informationsberichte des Bayer. Landesamtes für Wasserwirtschaft*, 1/94, 1–548.
- 117 Foissner W., Berger H., Blatterer H., and Kohmann F. (1995): Taxonomische und ökologische Revision der Ciliaten des Saprobiensystems. Band IV: Gymnostomata, *Loxodes*, Suctoria. *Informationsberichte des Bayer. Landesamtes für Wasserwirtschaft*, 1/95, 1–540.
- 118 Fresenius G. (1858): Beiträge zur Kenntnis mikroskopischer Organismen. Abh. senckenb. naturforsch. Ges., 2, 211–242.
- 119 Fryd-Versavel G. and Tuffrau M. (1978): Compléments a la connaissance du genre *Transitella* Gellert 1950. *Protistologica*, 14, 91–98.
- 120 Ganner B., Foissner W., and Adam H. (1987): Morphometric and biometric comparison of four populations of *Urosoomoida agiliformis* (Ciliophora, Hypotrichida). *Annls Sci. nat. (Zool.)*, 8 (years 1986–87), 199–207.
- 121 Gaston K. J. (1991): The magnitude of global insect species richness. *Conserv. Biol.*, 5, 283–296.
- 122 Gaston K. J. and Mound L. A. (1993): Taxonomy, hypothesis testing and the biodiversity crisis. *Proc. R. Soc. London B*, 251, 139–142.
- 123 Gelei J. v. (1934): Beiträge zur Ciliatenfauna der Umgebung von Szeged. II. Vier *Bryophyllum*-Arten. *Arch. Protistenk.*, 81, 201–230.
- 124 Gelei J. (1939): Adatok Szeged környékének ázalékkállatka világához. III. Néhány *Blepharisma* Szeged környékéről [Beiträge zur Ciliatenfauna der Umgebung von Szeged. II. Einige Blepharismen]. *Acta Univ. szeged.*, 5, 169–194 (in Hungarian with German summary).
- 125 Gelei J. v. (1954): Über die Lebensgemeinschaft einiger temporärer Tümpel auf einer Bergwiese im Börzsönygebirge (Oberungarn) III. Ciliaten. *Acta biol. hung.*, 5, 259–343.
- 126 Gellért J. (1942): Életegyüttes a fákéreg zöldporos bevonatában. *Acta Sci. math.-nat. Univ. Kolozsvár (N.F.)*, 8, 1–36 (in Hungarian).
- 127 Gellért J. (1950): A legkisebb vízi életter átmeneti végény formái (Espèces transitoires d'infusoires du minimum de biotop aquatique). *Hidrol. Közl.*, 30, 121–126 (in Hungarian with French summary on page 158).
- 128 Gellért J. (1955): Die Ciliaten des sich unter der Flechte *Parmelia saxatilis* Mass. gebildeten Humus. *Acta biol. hung.*, 6, 77–111.
- 129 Gellért J. (1956): Ciliaten des sich unter dem Moosrasen auf Felsen gebildeten Humus. *Acta biol. hung.*, 6, 337–359.
- 130 Gellért J. (1957): Nébány hazai lomblevelű és tülelevelű erdő talajának ciliáta faújá (Ciliatenfauna im Humus einiger ungarischen Laub- und Nadelholzwälder). *Annls Inst. Biol. porest hung.*, 24, 11–34 (in Hungarian with German summary).
- 131 Gellért J. and Tamás G. (1958): Detritusz-turzások kovámoszatainak és csíllósainak ökológiai vizsgálata a tihanyi-félsziget keleti partján (Ökologische Untersuchungen an Diatomeen und Ciliaten der Detritus-Drifte am Ostufer der Halbinsel Tihany). *Annls Inst. biol. Tihany*, 25, 217–240 (in Hungarian with German summary).
- 132 Grandori R. and Grandori L. (1934): Studi sui protozoi del terreno. *Boll. Lab. Zool. agr. Bachic. R. Ist. sup. agr. Milano*, 5, 1–341.
- 133 Grandori R. and Grandori L. (1935): Nuovi protozoi del terreno agrario (nota preliminare). *Boll. Lab. Zool. agr. Bachic. R. Ist. sup. agr. Milano*, 4, 64–80.
- 134 Greeff R. (1873): Vorkommen von Vorticellen in der Erde. *Sber. Ges. Beförd. ges. Naturw. Marburg*, 3, 23–24.
- 135 Greeff R. (1888): Land-Protozoen. *Sber. Ges. Beförd. ges. Naturw. Marburg*, 3, 90–158.
- 136 Groliere C. A. and Couteaux M. M. (1984): Morphologie et infraciliature de *Kahlilembus fusiformis* (Kahl, 1926) gen. nov., scuticociliate dé sol. *Acta Protozool.*, 23, 77–83.
- 137 Gruber A. (1879): Neue Infusorien. *Z. wiss. Zool.*, 33 (year 1880), 439–466.
- 138 Hemberger H. (1985): Neue Gattungen und Arten hypotricher Ciliaten. *Arch. Protistenk.*, 130, 397–417.
- 139 Hirshfield H. I., Isquith I. R., and Bhandary A. V. (1965): A proposed organization of the genus *Blepharisma* Perty and description of four new species. *J. Protozool.*, 12, 136–144.
- 140 Horváth J. v. (1932): Ein neues hypotriches Infusor, *Kahlia acrobates* nov. gen., nov. sp. *Arch. Protistenk.*, 77, 424–433.
- 141 Horváth J. v. (1933): Beiträge zur hypotrichen Fauna der Umgebung von Szeged. I. *Arch. Protistenk.*, 80, 281–302.
- 142 Horváth J. v. (1934): *Kahlia simplex* nov. sp. alkata, élettani megvilágításban. *Acta Litt. Scient. R. Univ. hung. Francisco-Josephina*, 3, 60–76 (in Hungarian).
- 143 Horváth J. (1956): Beiträge zur Kenntnis einiger neuer Bodenciliaten. *Arch. Protistenk.*, 101, 269–276.
- 144 Jankowski A. W. (1964): Morphology and evolution of Ciliophora. III. Diagnoses and phylogeny of 53 sapropelebiants, mainly of the order Heterotrichida. *Arch. Protistenk.*, 107, 185–294.
- 145 Jankowski A. W. (1967): New genera and subgenera of classes Gymnostomea and Ciliostomea. Materials of the V Conference of Young Scientists of Moldavia, year 1967, p. 36 (in Russian).
- 146 Jankowski A. W. (1978): Systematische Revision der Klasse Polyhymenophora (Spirotricha). *Trudy zool. Inst., Leningr.*, year 1978, 39–40 (in Russian).

- 147 Jankowski A. W. (1992): New generic combinations for some species of Ciliophora. *Cytologia*, 34, p. 171 (in Russian with English title translation).
- 148 Kahl A. (1926): Neue und wenig bekannte Formen der holotrichen und heterotrichen Ciliaten. *Arch. Protistenk.*, 55, 197–438.
- 149 Kahl A. (1927a): Neue und ergänzende Beobachtungen holotricher Ciliaten. I. *Arch. Protistenk.*, 60, 34–129.
- 150 Kahl A. (1927b): Neue und ergänzende Beobachtungen heterotricher Ciliaten. *Arch. Protistenk.*, 57, 121–203.
- 151 Kahl A. (1928): Die Infusorien (Ciliata) der Oldesloer Salzwasserstellen. *Arch. Hydrobiol.*, 19, 50–123.
- 152 Kahl A. (1930a): Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria) 1. Allgemeiner Teil und Prostomata. *Tierwelt Dtl.*, 18, 1–180.
- 153 Kahl A. (1930b): Neue und ergänzende Beobachtungen holotricher Infusorien. II. *Arch. Protistenk.*, 70, 313–416.
- 154 Kahl A. (1931): Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria) 2. Holotrichia außer den im 1. Teil behandelten Prostomata. *Tierwelt Dtl.*, 21, 181–398.
- 155 Kahl A. (1932): Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria) 3. Spirotricha. *Tierwelt Dtl.*, 25, 399–650.
- 156 Kahl A. (1934): Suctoria. *Tierwelt N.- und Ostsee*, 1 (II c5), 184–226.
- 157 Kahl A. (1935): Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria) 4. Peritricha und Chonotricha. *Tierwelt Dtl.*, 30, 651–886.
- 158 Kent W. S. (1880–1882): A manual of the infusoria: including a description of all known flagellate, ciliate, and tentaculiferous protozoa British and foreign, and an account of the organization and affinities of the sponges. Vols. I–III. David Bogue, London. 913 pp. (Vol. I 1880: 1–432; Vol. II 1881: 433–720; 1882: 721–913; Vol. III 1882: Plates).
- 159 Kowalewski M. (1882): Przyczynek do historyi naturalnej oxytrichów. Pam. fizyogr., 2, 395–413 (in Polish with French figure explanation; Polish spelling of name: Kowalewskiego).
- 160 Larsen H. F. (1983): Observations on the morphology and ecology of *Blepharisma lateritium* (Ehrenberg, 1831) Kahl, 1932. *Arch. Protistenk.*, 127, 65–80.
- 161 Lehle E. (1989): Beiträge zur Fauna der Ulmer Region II. Ciliaten (Protozoa: Ciliophora) Bioindikatoren in Waldböden. *Mitt. Ver. Naturwiss. Math. Ulm (Donau)*, 35, 131–156.
- 162 Lehle E. (1994): Die Auswirkungen von Düngung und Kalkung auf die Bodenciliaten (Protozoa: Ciliophora) eines Fichtenbestandes im Schwarzwald (Süddeutschland). *Arch. Protistenk.*, 144, 113–125.
- 163 Lepsi I. (1948): Protozoen aus Boden und Laubstreu eines Eichenwaldes. *Notat. biol.*, Buc., 6, 149–159.
- 164 Lepsi I. (1951): Modificarea faunei de protozoare tericole, prin irrigatii agricole. *Bul. Stiint. Sect. Stiinte Biol. Agron. Geol. Geog.*, 3, 513–523.
- 165 Maeda M. and Carey P. G. (1984): A revision of the genera *Trachelostyla* and *Gonostomum* (Ciliophora, Hypotrichida), including redescriptions of *T. pediculiformis* (Cohn, 1866) Kahl, 1932 and *T. caudata* Kahl, 1932. *Bull. Br. Mus. nat. Hist. (Zool.)*, 47, 1–17.
- 166 Matthes D. (1988): Suctoria (Sauginfusorien). *Protozoenfauna*, 7/1, I–XIII, 1–226 and 307–309.
- 167 Maupas E. (1883): Contribution a l'étude morphologique et anatomique des infusoires ciliés. *Archs Zool. exp. gén.*, 1, 427–664.
- 168 May R. M. and Nee S. (1995): The species alias problem. *Nature*, 378, 447–448.
- 169 Mermod G. (1914): Recherches sur la faune infusorienne des tourbières et des eaux voisines de Sainte-Croix (Jura vaudois). *Revue suisse Zool.*, 22, 31–114.
- 170 Mirabdullayev I. M. (1986): *Foissneria viridis* gen. et sp. n. (Synhymenida, Furgasoniidae), a new planktonic infusorian from Uzbekistan piscicultural ponds. *Zool. Zh.*, 65, 928–929 (in Russian with English summary).
- 171 Moody J. E. (1912): Observations on the life-history of two rare ciliates, *Spathidium spathula* and *Actinobolus radians*. *J. Morph.*, 23, 349–407.
- 172 Müller O. F. (1773): *Vermium terrestrium et fluviatilium, seu animalium infusorum, helminthicorum et testaceorum, non marinorum, succincta historia*. Heineck & Faber, Havniae et Lipsiae.
- 173 Müller O. F. (1786): *Animalcula infusoria fluviatilia et marina, quae detexit, systematice descripsit et ad vivum delineari curavit*. N. Mölleri, Hauniae.
- 174 Müller P. (1981): *Arealsysteme und Biogeographie*. Ulmer, Stuttgart.
- 175 Njine T. (1979): Structure et morphogenèse buccales du cilié *Leptopharynx* (Mermod, 1914). *Protistologica*, 15, 459–465.
- 176 Peck R. K. (1974): Morphology and morphogenesis of *Pseudomicrothorax*, *Glaucoma* and *Dexiotricha*, with emphasis on the types of stomatogenesis in holotrichous ciliates. *Protistologica*, 10, 333–369.
- 177 Penard E. (1914a): Les cothurnidés muscicoles. *Mém. Soc. Phys. Hist. nat. Genève*, 38, 19–66.
- 178 Penard E. (1914b): Sur quelques tentaculifères muscicoles. *Arch. Protistenk.*, 34, 277–294.
- 179 Penard E. (1922): Études sur les infusoires d'eau douce. Georg & Cie, Genève.
- 180 Perty M. (1849): Mikroskopische Organismen der Alpen und der italienischen Schweiz. *Mitt. naturf. Ges. Bern*, year 1849, 153–176.
- 181 Perty M. (1852): Zur Kenntniss kleinster Lebensformen nach Bau, Funktionen, Systematik, mit Specialverzeichniss der in der Schweiz beobachteten. Jent & Reinert, Bern.
- 182 Petz W. (1997): Ecology of the active soil microfauna (Protozoa, Metazoa) of Wilkes Land, East Antarctica. *Polar Biol.*, 18, 33–44.
- 183 Petz W. and Foissner W. (1996): Morphology and morphogenesis of *Lamnostyla edaphoni* Berger and Foissner and *Onychodromopsis flexilis* Stokes, two hypotrichs (Protozoa: Ciliophora) from Antarctic soils. *Acta Protzool.*, 35, 257–280.
- 184 Petz W. and Foissner W. (1997): Morphology and infraciliature of some soil ciliates (Protozoa, Ciliophora) from continental Antarctica, with notes on the morphogenesis of *Sterkiella histiomuscorum*. *Polar Record*, 33, 307–326.
- 185 Pomp R. and Wilbert N. (1988): Taxonomic and ecological studies of ciliates from Australian saline soils: colpodids and hymenostomate ciliates. *Aust. J. mar. Freshwat. Res.*, 39, 479–495.
- 186 Prelle A. (1968): Ultrastructures corticales du cilié holotriché *Drepanomonas dentata* Fresenius, 1858. *J. Protzool.*, 15, 517–520.
- 187 Quennerstedt A. (1869): Bidrag till sveriges infusoriefauna. *Acta Univ. Lund.*, 6, 1–35 (in Swedish).

- 188 Ruinen J. (1938): Notizen über Ciliaten aus konzentrierten Salzwässern. Zool. Med. Leiden, 20, 243–256.
- 189 Ryan P. G., Watkins B. P., Lewis Smith R. I., Dastych H., Eicker A., Foissner W., Heatwole H., Miller W. R., and Thompson G. (1989): Biological survey of Roberts-kollen, western Dronning Maud Land: area description and preliminary species list. S. Afr. T. Antarkt., 19, 10–20.
- 190 Sandon H. (1927): The composition and distribution of the protozoan fauna of the soil. Oliver and Boyd, Edinburgh and London.
- 191 Serrano S., Martín-González A., and Fernández-Galiano D. (1990): General morphology and cytological events during the conjugation process in *Acaryophrya collaris* (Kahl, 1926) (Ciliophora, Haptorina). Arch. Protistenk., 138, 65–74.
- 192 Shibuya M. (1929): Notes on two new hypotrichous ciliates from the soil. Proc. imp. Acad. Japan, 5, 155–156.
- 193 Shibuya M. (1930): Ciliates found in soils from some parts of Japan. J. imp. agric. Exp. Stn Tokyo, 1, 200, 212–214 (résumé of an article written in Japanese).
- 194 Shibuya M. (1931): Notes on two ciliates, *Cyrtolophosis mucicola* Stokes and *Gastrostyla philippinensis* sp. nov., found in the soil of the Philippines. Proc. imp. Acad. Japan, 7, 124–127.
- 195 Shin M. K. and Kim W. (1993): Redescription of two holostichid species of genus *Holosticha* Wrzesiowski 1877 (Ciliophora, Hypotrichida, Holostichidae) from Seoul, Korea. Korean J. Syst. Zool., 1, 251–259.
- 196 Silva da Neto I. D. (1993): Structural and ultrastructural observations of the ciliate *Phacodinium metchnicoffi* Certes, 1891 (Heterotrichaea, Phacodiniida). Europ. J. Protistol., 29, 209–218.
- 197 Smith H. G. (1973): The temperature relations and bi-polar biogeography of the ciliate genus *Colpoda*. Br. Antarct. Surv. Bull., 37, 7–13.
- 198 Smith H. G. (1978): The distribution and ecology of terrestrial protozoa of sub-Antarctic and maritime Antarctic islands. Br. Antarct. Surv. Sci. Rep., 95, 1–104.
- 199 Smith H. G. (1981): A comparative study of the terrestrial protzoa inhabiting moss turf peat on Iles Crozet South Georgia and the South Orkney Islands. Colloque sur les Ecosystèmes Subantarctiques (Paimpont, C.N.R.S.A.), 51, 137–145.
- 200 Smith J. C. (1899): Notices of some undescribed infusoria, from the infusorial fauna of Louisiana. Trans. Am. microsc. Soc., 20, 51–56.
- 201 Sondheim M. (1929): Protozoen aus der Ausbeute der Voeltzkowschen Reisen in Madagaskar und Ostafrika. Abh. senckenb. naturforsch. Ges., 41, 283–313.
- 202 Song W. (1990a): A comparative analysis of the morphology and morphogenesis of *Gonostomum strenua* (Engelmann, 1862) (Ciliophora, Hypotrichida) and related species. J. Protozool., 37, 249–257.
- 203 Song W. (1990b): Morphologie und Morphogenese des Bodenciliaten *Periholosticha wilberti* nov. spec. (Ciliophora, Hypotrichida). Arch. Protistenk., 138, 221–231.
- 204 Song W. (1991): The infraciliature and silverline system of *Euplates novemcarinata* Wang, 1930 (Ciliophora, Hypotrichida). J. Ocean Univ. Qingdao, 21, 76–84 (in Chinese with English summary).
- 205 Song W. (1994a): On two soil ciliates of the genus *Dileptus* (Ciliophora: Haptorida). Acta Zootax. Sin., 19, 385–391 (in Chinese with English summary).
- 206 Song W. (1994b): Faunistical studies on some soil ciliates from Qingdao. – I. Kinetofragminophora, Oligohymenophora, Colpodea. J. Ocean Univ. Qingdao, 24, 15–23 (in Chinese with English summary).
- 207 Song W. and Gao L. (1994): Observations on the soil ciliate *Acineria uncinata* with light and SEM methods (Ciliophora, Pleurostomatida). Zool. Res., 15, 10–14 (in Chinese with English summary).
- 208 Song W. and Wilbert N. (1988): *Parabakuella typica* nov. gen., nov. spec. (Ciliata, Hypotrichida) aus dem Edaphon eines Standortes in Qingdao, China. Arch. Protistenk., 135, 319–325.
- 209 Song W. and Wilbert N. (1989a): Morphology and infraciliature of *Uroleptoides qingdaensis* sp. nov. (Ciliophora; Hypotrichida; Amphisiellidae). Acta Zootax. Sin., 14, 390–395 (in Chinese with English summary).
- 210 Song W. and Wilbert N. (1989b): Taxonomische Untersuchungen an Aufwuchsciliaten (Protozoa, Ciliophora) im Poppelsdorfer Weiher, Bonn. Lauterbornia, 3, 2–221.
- 211 Song W., Packroff G., and Wilbert N. (1988): Morphologie und Infraciliatur von *Dileptus orientalis* sp. n., einem Bodenciliaten aus Qingdao, China. Acta Protozool., 27, 271–277.
- 212 Song W., Wilbert N., and Berger H. (1992): Morphology and morphogenesis of the soil ciliate *Bakuella edaphoni* nov. spec. and revision of the genus *Bakuella* Agamaliev & Alekperov, 1976 (Ciliophora, Hypotrichida). Bull. Br. Mus. nat. Hist. (Zool.), 58, 133–148.
- 213 Šrámek-Hušek R. (1952): O několika zajímavých nálevnících z českých mechů (Über einige bemerkenswerte Ciliaten aus böhmischen Moosen). Čslka. Biol., Praha, 1, 175–184 (in Czech).
- 214 Šrámek-Hušek R. (1957): K poznání nálevníků ostravského kraje [Zur Kenntnis der Ciliaten des Ostrauer-Gebietes (Tschechoslowakei)]. Věst. čsl. zool. Spol., 21, 1–24 (in Czech with German summary).
- 215 Stein F. (1859): Der Organismus der Infusionsthiere nach eigenen Forschungen in systematischer Reihenfolge bearbeitet. I. Abtheilung. Allgemeiner Theil und Naturgeschichte der hypotrichen Infusionsthiere. W. Engelmann, Leipzig.
- 216 Stein F. (1867): Der Organismus der Infusionsthiere nach eigenen Forschungen in systematischer Reihenfolge bearbeitet. II. Abtheilung. 1) Darstellung der neuesten Forschungsergebnisse über Bau, Fortpflanzung und Entwicklung der Infusionsthiere. 2) Naturgeschichte der heterotrichen Infusorien. W. Engelmann, Leipzig.
- 217 Sterki V. (1878): Beiträge zur Morphologie der Oxytrichinen. Z. wiss. Zool., 31, 29–58.
- 218 Stokes A. C. (1882): A new *Thuricola*. Am. mon. microsc. J., 3, 181–183.
- 219 Stokes A. C. (1884): Notes on some apparently undescribed forms of fresh-water infusoria. Am. J. Sci., 28, 38–49.
- 220 Stokes A. C. (1886a): Some new hypotrichous infusoria. Proc. Am. phil. Soc., 23, 21–30.
- 221 Stokes A. C. (1886b): Notices of new fresh-water infusoria. Proc. Am. phil. Soc., 23, 562–568.
- 222 Stokes A. C. (1887a): Some new hypotrichous infusoria from American fresh waters. Ann. Mag. nat. Hist., 20, 104–114.
- 223 Stokes A. C. (1887b): Notices of new fresh-water infusoria. Proc. Am. phil. Soc., 24, 244–255.
- 224 Stokes A. C. (1893): Notices of some undescribed infu-

- soria from brackish waters of the eastern United States. Jl R. microsc. Soc., year 1893, 298–302.
- 225 Stokes A. C. (1894): Notices of presumably undescribed infusoria. Proc. Am. phil. Soc., 33, 338–344.
- 226 Suzuki S. (1954): Taxonomic studies on *Blepharisma undulans* Stein with special reference to the macronuclear variation. J. Sci. Hiroshima Univ., 15, 205–220.
- 227 Szabó M. (1934): Beiträge zur Kenntnis der Gattung *Halteria* (Protozoa, Ciliata). Arb. ung. biol. ForschInst., 7, 95–106 (in German and Hungarian).
- 228 Thompson J. C., Jr. (1966): *Pseudocochilembus marinus* n. sp., a hymenostome ciliate from the Virginia coast. J. Protozool., 13, 463–465.
- 229 Tirjaková E. (1988): Structures and dynamics of communities of ciliated protozoa (Ciliophora) in field communities. I. Species composition, group dominance, communities. Biológia, Bratisl., 43, 497–503.
- 230 Tirjaková E. and Matis D. (1987): Ciliates of dry mosses in Bratislava in relation to air pollution. Acta Fac. Rerum nat. Univ. comen., Bratisl. Series Zoologia, 29, 17–31.
- 231 Trueba F. J. (1980): A taxonomic revision of the peritrich ciliate genera *Thuricola* and *Pseudothuricola*. Beaufortia, 30, 125–138.
- 232 Tucolesco J. (1962): Études protozoologiques sur les eaux Roumaines I. Espèces nouvelles d'infusoires de la mer noire et des bassins salés paramarins. Arch. Protistenk., 106, 1–36.
- 233 Van der Land J. (1964): A new peritrichous ciliate as a symphoriont on a tardigrade. Zool. Med. Leiden, 39, 85–88.
- 234 Villeneuve-Brachon S. (1940): Recherches sur les ciliés hétérotriches: cinétoème, argyrome, myonèmes, formes nouvelles ou peu connues. Archs Zool. exp. gén., 82, 1–180.
- 235 Vuxanovici A. (1962a): Contributii la sistematica ciliatelor (Nota I). Studii Cerc. Biol. (Biol. Anim.), 14, 197–215 (in Roumanian with Russian and French summaries).
- 236 Vuxanovici A. (1962b): Contributii la sistematica ciliatelor (Nota III). Studii Cerc. Biol. (Biol. Anim.), 14, 549–573 (in Roumanian with Russian and French summaries).
- 237 Vuxanovici A. (1963): Contributii la studiul speciilor din subordinul Hypotricha (Ciliata) (Nota I). Studii Cerc. Biol. (Biol. Anim.), 15, 199–222 (in Roumanian with Russian and French summaries).
- 238 Warren A. (1982): A taxonomic revision of the genus *Platycola* (Ciliophora: Peritrichida). Bull. Br. Mus. nat. Hist. (Zool.), 43, 95–108.
- 239 Warren A. and Paynter J. (1991): A revision of *Cothurnia* (Ciliophora: Peritrichida) and its morphological relatives. Bull. Br. Mus. nat. Hist. (Zool.), 57, 17–59.
- 240 Wenzel F. (1953): Die Ciliaten der Moosrasen trockner Standorte. Arch. Protistenk., 99, 70–141.
- 241 Wenzel F. (1955): Über eine Artentstehung innerhalb der Gattung *Spathidium* (Holotricha, Ciliata). [S. *ascendens* n. sp. und S. *polymorphum* n. sp.]. Arch. Protistenk., 100, 514–540.
- 242 Wiackowski K. (1985): The morphology and morphogenesis of *Keronella gracilis* n. gen., n. spec. (Hypotrichida, Ciliophora). Protistologica, 21, 81–91.
- 243 Wilbert N. (1982): Ein neuer colpodider Ciliat aus einer Grassteppe in Ningerhar, Afghanistan: *Colpodidium caudatum* nov. gen., nov. spec. Arch. Protistenk., 125, 291–296.
- 244 Wilbert N. and Kahan D. (1986): *Semiplatyophrya foissneri* nov. gen., nov. spec. und *Perisincirra pori* nov. spec., Ciliaten aus einem Salzboden des Sinai. Arch. Protistenk., 131, 129–138.
- 245 Wirnsberger E., Foissner W., and Adam H. (1985): Morphological, biometric, and morphogenetic comparison of two closely related species, *Stylonychia vorax* and *S. pustulata* (Ciliophora: Oxytrichidae). J. Protozool., 32, 261–268.
- 246 Wirnsberger-Aesch E., Foissner W., and Foissner I. (1989): Morphogenesis and ultrastructure of the soil ciliate *Engelmanniella mobilis* (Ciliophora, Hypotrichida). Europ. J. Protistol., 24, 354–368.
- 247 Wirnsberger-Aesch E., Foissner W., and Foissner I. (1990): Natural and cultured variability of *Engelmanniella mobilis* (Ciliophora, Hypotrichida); with notes on the ultrastructure of its resting cyst. Arch. Protistenk., 138, 29–49.
- 248 Wrześniowski A. (1866): Spis wymoczków spostrzeganych w Warszawie i jej okolicach wlatach 1861 do 1865 (Verzeichnis der Infusorien, welche in Warschau und seiner Umgebungen von 1861–65 gesammelt wurden). Wykaz Szkoty Głównej Warszawskiej, 5, 15–28 (in Polish with Latin diagnoses of new taxa; Polish spelling of name: Wrzesińskiego).
- 249 Wrześniowski A. (1870): Beobachtungen über Infusorien aus der Umgebung von Warschau. Z. wiss. Zool., 20, 467–511.

Additional References

- 250 Foissner W. (1979): Taxonomische Studien über die Ciliaten des Großglocknergebietes (Hohe Tauern, Österreich). Familien Microthoracidae, Chilonellidae und Furgasoniidae. Sber. Akad. Wiss. Wien, 188, 27–43.
- 251 Foissner W. (1980): Taxonomische Studien über die Ciliaten des Großglocknergebietes (Hohe Tauern, Österreich) IV. Familien Spathidiidae, Podophryidae und Urnulidae. Verh. zool.-bot. Ges. Wien, 118/119, 97–112.
- 252 Shen Y., Liu J., Song B., and Gu M. (1992): Protozoa. In: Yin W. et al. (eds.): Subtropical Soil Animals of China, pp. 97–156. Science Press, Beijing (in Chinese).
- 253 Voss H.-J. (1991a): Die Morphogenese von *Notohyphmena rubescens* Blatterer & Foissner, 1988 (Ciliophora, Oxytrichidae). Arch. Protistenk., 140, 219–236.
- 254 Voss H.-J. (1991b): Die Morphogenese von *Cyrtophysa muscorum* (Kahl, 1932) Foissner, 1989 (Ciliophora, Oxytrichidae). Arch. Protistenk., 140, 67–81.
- 255 Voß H.-J. (1992): Morphogenesis in *Amphisialla australis* Blatterer & Foissner, 1988 (Ciliophora, Hypotrichida). Europ. J. Protistol., 28, 405–414.
- 256 Borror A. C. and Hill B. F. (1995): The order Euplotida (Ciliophora): taxonomy, with division of *Euplotes* into several genera. J. Euk. Microbiol., 42, 457–466.
- 257 Voss H.-J. (1989): Vergleichende morphogenetische Untersuchung von 13 Arten der Gattung *Euplotes* (Ciliophora, Hypotrichida). Arch. Protistenkd., 137, 331–344.

Address for correspondence: Wilhelm Foissner, Universität Salzburg, Institut für Zoologie, Hellbrunnerstrasse 34, A - 5020 Salzburg, Austria (Europe).